


POLICY PLATFORM

Challenges and opportunities for control and elimination of soil-transmitted helminth infection beyond 2020

Matthew C. Freeman^{1*}, **Oladele Akogun**², **Vicente Belizario, Jr.**³, **Simon J. Brooker**⁴, **Theresa W. Gyorkos**⁵, **Rubina Imtiaz**⁶, **Alejandro Krolewiecki**⁷, **Seung Lee**⁸, **Sultani H. Matendechero**⁹, **Rachel L. Pullan**¹⁰, **Jürg Utzinger**^{11,12}

1 Department of Environmental Health, Emory University, Atlanta, Georgia, United States of America, **2** Modibbo Adama University of Technology, Yola, Nigeria, **3** College of Public Health, University of the Philippines Manila, Manila, the Philippines, **4** Global Health, Bill & Melinda Gates Foundation, Seattle, Washington, United States of America, **5** Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Quebec, Canada, **6** Children Without Worms, The Task Force for Global Health, Decatur, Georgia, United States of America, **7** Instituto de Investigaciones en Enfermedades Tropicales, Universidad Nacional de Salta, Oran, Argentina, **8** Save the Children, Washington, DC, United States of America, **9** Ministry of Health, Nairobi, Kenya, **10** London School of Hygiene & Tropical Medicine, London, United Kingdom, **11** Swiss Tropical and Public Health Institute, Basel, Switzerland, **12** University of Basel, Basel, Switzerland

* matthew.freeman@emory.edu



OPEN ACCESS

Citation: Freeman MC, Akogun O, Belizario V, Jr., Brooker SJ, Gyorkos TW, Imtiaz R, et al. (2019) Challenges and opportunities for control and elimination of soil-transmitted helminth infection beyond 2020. *PLoS Negl Trop Dis* 13(4): e0007201. <https://doi.org/10.1371/journal.pntd.0007201>

Editor: Joanne P. Webster, Imperial College London, UNITED KINGDOM

Published: April 11, 2019

Copyright: © 2019 Freeman et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

More than half of the world’s population lives in places endemic for soil-transmitted helminths (STHs), and an estimated 1.45 billion people are infected [1,2]. In 2017, the global burden of STH infection (*Ascaris lumbricoides*, hookworm, and *Trichuris trichiura*) was estimated at 1.9 million disability-adjusted life years (DALYs) [3]. Moderate and heavy infection intensity and chronic STH infection are associated with anemia, malnutrition, educational loss, and cognitive deficits, but recent systematic reviews and meta-analyses produced conflicting results on the impact of preventive chemotherapy (PC) [4–6].

The [Soil-Transmitted Helminthiasis Advisory Committee](#) (hereafter called “the Committee”) is a group of independent experts with a broad range of expertise. It is convened annually by Children Without Worms (CWW), an organization whose purpose is to utilize available evidence to identify best practices and opportunities for the prevention and control of STH infection [7]. On November 1 and 2, 2017, the Committee met in Baltimore, Maryland, United States of America, in order to discuss the critical need to develop a data-driven guide to the STH endgame on late-stage program functioning, processes, and surveillance. The focus was on research and field experiences from countries approaching the “elimination of STH infection as a public health problem” after consecutive years of PC and countries that are now considering scaling down their PC frequency but may be concerned about infection rebound. Emphasis was placed on interim recommendations for monitoring and decision-making for national program managers desiring to achieve the World Health Organization (WHO) goal of eliminating STH infection as a public health problem by 2020, particularly related to STH infections in risk groups other than school-age children (SAC), namely preschool-age children (PSAC) and women of reproductive age (WRA) [8]. The following is the Committee’s recommendations stemming from the Baltimore meeting in November 2017. It complements and updates the publication derived by the Committee’s meeting a year earlier in Basel, Switzerland

[7], and was instrumental in shaping the agenda for the October 2018 meeting, convened jointly by CWW and WHO, with recommendations to be reported elsewhere.

The 2020 roadmap and beyond

In its roadmap for implementation for 2020, WHO set a goal to achieve at least 75% coverage of PC—either annual or biannual—of SAC and PSAC [9–11]. As we approach 2020, it is imperative that we not only accelerate what has worked for the control of STH infection-related morbidity but that we look beyond 2020 and better understand what more is required to eliminate STH infection as a public health problem.

Fig 1 summarizes progress made in terms of both coverage and impact using 2016 country data. Despite considerable gains in SAC coverage, less than half of the at-risk countries are treating PSAC, a number that has not changed much in recent years, yielding a total coverage level of approximately 50% but with considerable variability of coverage from year to year and across countries (Fig 2). If the current trend in PC coverage persists, PSAC and the combined group of children aged 1–14 years will not reach the goal by 2020. Accelerating PSAC PC coverage might avoid that failure, and hence, needs a clear, strong global policy now. While the initial focus on SAC coverage has enabled the mobilization of resources, other priorities have emerged, both in places and populations in which the target was not achieved and within areas that are now shifting priority from scaling up PC to eliminating STH infection as a public health problem—defined by the WHO as when less than 1% of the at-risk population has moderate or heavy infection (MHI) [8]—and potentially interrupting the transmission of STH infection [12]. WHO recommends stopping PC once less than 1% of the at-risk population has MHI infections. Surveillance will, however, need to continue in order to pick up potential recurrence of infection and to plan further intervention, if warranted. Hence, as we turn our sights towards and beyond 2020, the Committee recognized it as timely to review and assess the successes and challenges of progress made to date. Indeed, it is hoped that the Baltimore 2017 meeting deliberations will inform future control strategies and targets.

There have been many important developments by the time the Committee convened in November 2017, but six are particularly noteworthy:

- updated PC guidelines published by WHO for all at-risk populations [13];
- the Bellagio Declaration focusing on girls and WRA [14];
- widening of WHO’s engagement with experts around the globe [15];
- success of the Global Program to Eliminate Lymphatic Filariasis (GPELF) [16];
- increasing importance for deworming programs to measure impact [17]; and
- the launch by WHO-AFRO of the Expanded Special Project on the Elimination of Neglected Tropical Diseases (ESPEN) portal (<http://espen.afro.who.int>), which provides—for the first time—subnational data on disease endemicity and PC coverage for each of the five PC-NTDs, including STH infection.

In this Policy Platform, we lay out critical challenges in seven key areas that need attention, discuss progress until November 2017, and put forward recommendations for immediate action. Our recommendations arose at a critical juncture for STH control efforts, as current global policies, goals, and related strategies and resources are revitalized through 2020, a year which is upon us.

PC STH not started (n=8)	PC STH coverage < 75% (n=31)		PC STH coverage ≥75% (less than 5 years) (n=37)		PC STH coverage ≥75% (5 years and more) (n=20)	MHI < 1% (n=7)
Antigua and Barbuda	Angola	Micronesia	Afghanistan	Philippines	Bangladesh	Benin*
Bahamas	Armenia	(Federated States)	Azerbaijan	South Africa	Belize	Bhutan*
Botswana	Bolivia	Namibia	Cabo Verde	Swaziland	Burundi	Burkina Faso*
Cuba	Brazil	Pakistan	Côte d'Ivoire	Timor-Leste	Cambodia	Mali*
Dominica	Central African Republic	Papua New Guinea	Ecuador	Togo*	Cameroon*	Niger*
Nauru	Chad	Sao Tome and Principe	El Salvador	Tonga	Dominican Republic	Paraguay*
Saint Lucia	China	Solomon Islands	Fiji	Uganda	Democratic People's Republic of Korea	Senegal*
Suriname	Colombia	Somalia	Gambia	United Republic of Tanzania	Republic of Korea	
	Comoros	South Sudan	Guatemala	Uzbekistan	Ghana	
	Congo	Sudan	Guinea	Vanuatu	Haiti	
	Dem. Republic of the Congo	Trinidad and Tobago	Guyana	Venezuela	Kiribati	
	Djibouti		Honduras	Viet Nam	Kyrgyzstan	
	Equatorial Guinea		India	Yemen	Lao People's	
	Ethiopia		Iraq	Zambia	Malawi	
	Gabon		Liberia	Zimbabwe	Mexico*	
	Georgia		Madagascar		Myanmar	
	Indonesia		Marshall Islands		Nicaragua	
	Jamaica		Mozambique		Rwanda*	
	Kenya		Nepal		Sierra Leone*	
	Lesotho		Nigeria		Tajikistan	
			Panama		Tuvalu	
			Peru			

Fig 1. Progress for PC coverage (by country) in SAC and where STH infection is no longer considered a public health problem (MHI <1%). Source: 2016 PC data from WHO. * Country with detailed epidemiologic information available. MHI, moderate or heavy infection; PC, preventive chemotherapy; SAC, school-age children; STH, soil-transmitted helminth.

<https://doi.org/10.1371/journal.pntd.0007201.g001>




Fig 2. Global PSAC treatment and coverage, 2006–2016. Source: WHO PC Databank, PSAC PC coverage data from 2006–2016; http://www.who.int/neglected_diseases/preventive_chemotherapy/lf/en/. * Coverage is calculated by dividing the number of children requiring PC and treated by the total number of children in need of PC. PC, preventive chemotherapy; PSAC, preschool children; WHO, World Health Organization.

<https://doi.org/10.1371/journal.pntd.0007201.g002>

Challenges and recommendations

Challenge #1: Incomplete and inconsistent monitoring of program impact

Recommendation: Define standard impact goals and targets post-2020. To date, programs mainly focused on reporting on PC coverage, as guided by a simplified coverage target specified by the Roadmap; yet there are few standardized data on program impact. It is noteworthy that country programs are increasingly interested in quantifying the impact of deworming on health outcomes [17–19]. Current approaches to measuring impact vary across countries and across deworming program implementers, limiting comparability and the possibility to appreciate changes over time and across countries and regions. Moreover, impact data are not readily available, with most of the evaluations conducted by research groups. For the Africa region, there has been recent progress in data sharing and transparency for neglected tropical diseases (NTDs), through the work of ESPEN and their data portal [20]. Currently, there is availability of data on endemicity and coverage at the subnational level for the five PC-NTDs, including STH infection, for 47 of 49 countries in the AFRO region. This successful approach to reporting sub-national data should be expanded to the other WHO regions where STHs are endemic.

The ESPEN portal is starting to include impact assessment data for lymphatic filariasis and onchocerciasis and there is an opportunity to include comparable data for STH infection. However, this will require a standardized and comprehensive monitoring and evaluation (M&E) framework that is tied to clearly defined, quantitative goals. For example, globally, a reported 69.5% of SAC and 50.8% of PSAC requiring PC reportedly received PC in 2016 [21]. It is conceivable that the observed scale-up of PC targeting STH over the past decade was a major contributor to the decline in the global burden of STH infection [22–25]; yet the precise extent to which this coverage has reduced prevalence, intensity, and burden of STH infection is not known. Country programs, in particular, require a comprehensive, standardized, yet flexible approach to measure progress toward morbidity-related goals. Such an approach would capture essential programmatic elements, and be used by each country to map their needs, commitment, and resources. This will generate realistic timelines and planning processes as well as alerting WHO, partners, and donors to better assess the resource and technical capacity needs of each program. Subnational data with standardized indicators for anthelmintic drug availability and coverage of the target populations are critical to track progress at the subnational level where PC program implementation may not be uniform. Data can also be disaggregated (e.g., by district, sex, and age categories, and any other useful determinant) to better understand the equity of program access and delivery for the three target risk groups (i.e., PSAC, SAC, and WRA). PC needs for refugees and migrants are also increasingly being recognized, and hence, WHO is considering to add these populations to the at-risk groups, while some countries (especially those in AFRO and EMRO regions) are actively engaged in estimating access, burden, and resource issues for migrants.

Challenge #2: Reaching at-risk groups other than SAC

Recommendation: Identify new PC strategies, platforms, and reporting mechanisms. Deworming of SAC has been shown to reduce disease burden, especially reducing high-burden infections in a cost-effective manner, but the empirical evidence from both multiyear deworming programs and modeling studies suggest that targeting SAC alone for PC is insufficient for sustained control and elimination of STH infection [17,26,27]. In particular, this is the case for *T. trichiura* due to poor drug efficacy [28,29] and for hookworm due to the age

distribution of infection. Indeed, in many settings, adults are at particularly high risk of hookworm infection and thus contribute substantially to transmission [30]. These issues are in contrast to countries where school-based deworming has been coupled with community-based programs in both experimental studies and as part of lymphatic filariasis control programs that target entire communities [31]. Using an exclusive school-based intervention platform potentially excludes 12.8% SAC who are out of school, with some countries like South Sudan having 66% of their SAC out of school [32]. With that and the availability of a pediatric mebendazole preparation as a donation for PSAC, out-of-school children and PSAC can be immediately prioritized to help achieve the overall children's coverage goal. Recent progress has been made on a more comprehensive preventive approach to include all risk groups, as demonstrated by the Kenya "Breaking Transmission Strategy" in addition to community-wide PC coverage through the lymphatic filariasis elimination program through USAID. Successful PC interventions targeting SAC by most countries are generating requests for specific guidance on the next operational phase: how to efficiently implement sentinel surveillance and which indicators to measure in order to predict, detect, and treat widely dispersed and persistent pockets of transmission. There is a need to consider how to scale up approaches to reach PSAC, given the current drug availability and added cost of this approach. Consistent with these needs is a recognized gap in our current knowledge of disease transmission at low prevalence and persistent environmental factors that facilitate transmission. Thus, there is a renewed need to clearly identify research gaps and questions that would facilitate implementation in these settings.

A recent report has provided guidance and recommendations for WRA [33]. Building on this report, WHO needs to develop implementation guidelines linked to clearly defined targets, both for WRA and PSAC as critical populations at risk of high STH burden [7,14]. Operational research is also needed to define platforms, partners, and recording and reporting tools to monitor progress of control programs targeting PSAC and WRA. In addition, there remains the challenge of providing additional anthelmintic drugs necessary to treat WRA (not targeted by the current donations). There is optimism that there will be the possible donation of chewable Vermox from Johnson & Johnson (J&J) for PSAC, but the quantities donated may not be enough to cover the total numbers at risk. Consideration of new drug options for WRA needs immediate discussion at national, regional, and global fora. Data from the GPELF and other community-based PC programs may provide insight into efficacy and safety issues through birth cohort studies.

Challenge #3: The risk of anthelmintic drug resistance

Recommendation: Develop standardized indicators to detect emerging resistance.

Experience from the veterinary sector demonstrated that anthelmintic drug resistance developed after years of large-scale monotherapy [34]. We suspect that if we wait until resistance is clinically detected in humans, it will be too late to respond [35]. While progress is being made, there are currently no routine, field-applicable diagnostics that can effectively identify and monitor signs of emerging resistance, so research on developing such tests urgently needs financial support. Indeed, we need to better identify resistant genes and to track refugia (i.e., that proportion of the worm population that remains susceptible to anthelmintic drugs). Human populations with a long history of single-drug deworming that have low worm burdens but have not reached transmission break points are likely to be at the greatest risk for the development of drug-resistant parasites. For diagnostic approaches such as quantitative polymerase chain reaction (qPCR) to be used in an STH-programmatic setting, we need to take molecular diagnostics to a level at which we can use its full potential by standardizing analysis and reporting and including appropriate quality control measures [36]. A

standardized approach to the monitoring of potential emerging resistance needs to be established, especially in those countries that have mature PC programs (e.g., Mexico and Togo, among others), to track both drug efficacy and mutations known to be associated with resistance [37].

Challenge #4: Poor diagnostics to assess program needs by implementation stage

Recommendation: Employ validated program stage-specific diagnostic techniques.

There is a need for new diagnostics that are appropriate for informing key decision points for national STH control programs [38,39]. The Kato–Katz technique, while relatively inexpensive, widely used, and sensitive in detecting MHI, will have lower positive predictive values in low-prevalence (and low-intensity) settings [40,41]. In addition, some preliminary analysis from microscopy and PCR has suggested that hookworm infection may be misidentified. New diagnostic tests have been validated [7], yet there is a need for novel, highly sensitive tests that can be employed in settings that move from STH control to elimination. However, their specific role and use in the context of national program implementation needs to be assessed [41]. Furthermore, capacity strengthening for national programs will be needed to fully take advantage of any new diagnostics. Field and laboratory protocols need to be standardized, reference laboratories established, and training developed for different contexts and languages. Gaps in the current diagnostic landscape have been identified, and formative research is underway. The next steps are to finalize and disseminate results and to identify additional resources for gap areas (e.g., sustaining animal models, conducting field studies to calculate test performance in various settings, etc.). Guidelines will then need to be developed to inform country programs on how to incorporate these tests for improved assessment of program impact and further planning.

Challenge #5: Limited efficacy of current drugs and gaps in drug availability

Recommendation: Promote research into combination therapies and fast track pre-qualification processes. The existing anthelmintic drugs have variable efficacies against different STH species, with particularly low efficacies against *T. trichiura* when using single-dose treatments [29]. New efforts must be undertaken to identify and provide guidelines for use cases for combination therapies in general and specifically in which *T. trichiura* is the predominant species [42]. In 2018, the WHO Essential Medicines Committee approved the inclusion of ivermectin for both STH and *Strongyloides stercoralis* to the WHO Model List of Essential Medicines [43]. However, there are implications and potential challenges in adding ivermectin to albendazole and mebendazole in STH deworming programs. Merck provides ivermectin dedicated for the control of lymphatic filariasis and onchocerciasis, and new manufacturers will need to become prequalified to meet the growing demand from STH control programs. Additionally, bioequivalence studies and other considerations will be needed to make the drug available at low cost. This will take time and requires innovative financing mechanisms.

Challenge #6: Limited coordination with the water, sanitation, and hygiene (WASH) sector

Recommendation: Identify WASH indicator(s) to be included in routine STH M&E.

As articulated in an editorial put forth in *The Lancet* [44], the 2020 WHO roadmap identified the critical role of WASH in the control of STH infection but did not set actionable targets or

strategies [11]. The STH community has largely avoided establishing an approach for addressing WASH or establishing a structure for engagement with the WASH sector. However, the recent development of the WASH–NTD joint strategy provides an entry point and guidance for improved communication, coordination, and collaboration [45]. Ample observational evidence [46,47], some recent randomized trials, biological plausibility, and history suggest that improved WASH is critical in the control and elimination of STH infection. Alignment with the WASH sector, specifically including WASH indicators as part of STH M&E [48] (and vice versa) by using data to advocate for WASH activities in STH-endemic areas, employing program monitoring, and conducting operational research and advocacy to ensure normative inclusion of STH-related WASH behaviors (e.g., shoe-wearing [49]), would be valuable contributions in STH control programming. Better quantification of the specific mechanisms and use of consistent indicators across programs would provide support to the WASH sector on gaps in typical WASH programming (e.g., type of water sources or food hygiene).

Challenge #7: New targets needed for post-2020

Recommendation: Develop clearly defined, quantitative program targets for all at-risk groups and move beyond PC coverage estimates. One of the most important next steps as we move toward 2020 and beyond is the need to establish clearly defined, quantitative program goals and targets post-2020. The current targets of achieving at least 75% coverage for deworming for SAC (and PSAC) is in reach, but we know that even meeting this target is insufficient to achieve elimination of STH as a public health problem. We must also critically evaluate whether the threshold of 1% prevalence of MHI is useful for M&E of STH morbidity control and moving toward elimination. There is a need to look beyond simple PC coverage measures and include impact targets for PSAC and WRA, benchmarks for WASH that will encourage investment in WASH in STH-endemic areas, and estimates of PC uptake along the distribution chain that would give accurate estimates for not only availability of the drug, but population compliance. Addressing Challenges 1–6, articulated above, will support this critical effort.

Outlook

There has been substantial progress in increasing coverage of PC for the control of STH infection, particularly among SAC and, to a lesser extent, among PSAC. As we approach 2020, work remains to accelerate action to achieve these targets in many places. At the same time, we need to think critically about what is needed beyond the 2020 roadmap and increase efforts in the seven areas discussed in this Policy Platform. This needs to be achieved through active collaboration and coordination by pertinent government ministries, researchers, donors, WHO, drug manufacturers, and multisectoral collaboration [50]. In doing so, it will help ensure progress toward eliminating STH infection—and other NTDs—as a public health problem, and it will yield more efficient allocation of resources and greater sustained impact, driven by targets and thresholds based on scientific evidence.

References

1. Pullan RL, Brooker SJ (2012) The global limits and population at risk of soil-transmitted helminth infections in 2010. *Parasit Vectors* 5: 81. <https://doi.org/10.1186/1756-3305-5-81> PMID: 22537799
2. Pullan RL, Smith JL, Jasrasaria R, Brooker SJ (2014) Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit Vectors* 7: 37. <https://doi.org/10.1186/1756-3305-7-37> PMID: 24447578
3. GBD 2017 DALYs and HALE Collaborators (2018) Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and

- territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 392: 1859–1922. [https://doi.org/10.1016/S0140-6736\(18\)32335-3](https://doi.org/10.1016/S0140-6736(18)32335-3) PMID: 30415748
4. Owada K, Nielsen M, Lau CL, Clements ACA, Yakob L, et al. (2017) Measuring the effect of soil-transmitted helminth infections on cognitive function in children: systematic review and critical appraisal of evidence. *Adv Parasitol* 98: 1–37. <https://doi.org/10.1016/bs.apar.2017.05.002> PMID: 28942767
 5. Welch VA, Ghogomu E, Hossain A, Awasthi S, Bhutta ZA, et al. (2017) Mass deworming to improve developmental health and wellbeing of children in low-income and middle-income countries: a systematic review and network meta-analysis. *Lancet Glob Health* 5: e40–e50. PMID: 27955788
 6. Pabalan N, Singian E, Tabangay L, Jarjanazi H, Boivin MJ, et al. (2018) Soil-transmitted helminth infection, loss of education and cognitive impairment in school-aged children: a systematic review and meta-analysis. *PLoS Negl Trop Dis* 12: e0005523. <https://doi.org/10.1371/journal.pntd.0005523> PMID: 29329288
 7. Becker SL, Liwanag HJ, Snyder JS, Akogun O, Belizario V Jr., et al. (2018) Toward the 2020 goal of soil-transmitted helminthiasis control and elimination. *PLoS Negl Trop Dis* 12: e0006606. <https://doi.org/10.1371/journal.pntd.0006606> PMID: 30106975
 8. WHO (2012) Soil-transmitted helminthiasis: eliminating as public health problem soil-transmitted helminthiasis in children: progress report 2001–2010 and strategic plan 2011–2020. Geneva: World Health Organization.
 9. WHO (2012) Accelerating work to overcome the global impact of neglected tropical diseases: a road-map for implementation. Geneva: World Health Organization.
 10. Brooker SJ, Nikolay B, Balabanova D, Pullan RL (2015) Global feasibility assessment of interrupting the transmission of soil-transmitted helminths: a statistical modelling study. *Lancet Infect Dis* 15: 941–950. PMID: 25886799
 11. WHO (2017) Integrating neglected tropical diseases in global health and development: fourth WHO report on neglected tropical diseases. Geneva: World Health Organization.
 12. Ásbjörnsdóttir KH, Ajjampur SSR, Anderson RM, Bailey R, Gardiner I, et al. (2018). Assessing the feasibility of interrupting the transmission of soil-transmitted helminths through mass drug administration: the DeWorm3 cluster randomized trial protocol. *PLoS Negl Trop Dis* 12: e0006166. <https://doi.org/10.1371/journal.pntd.0006166> PMID: 29346377
 13. WHO (2017) Preventive chemotherapy to control soil-transmitted helminth infections in at-risk population groups. Geneva: World Health Organization.
 14. WHO (2018) Reaching girls and women of reproductive age with deworming. Report of the advisory group on deworming in girls and women of reproductive age. Geneva: World Health Organization.
 15. Uniting to Combat Neglected Tropical Diseases (2018) Reaching a billion: ending neglected tropical diseases: a gateway to universal health coverage. Fifth progress report on the London Declaration on NTDs. <https://unitingtocombatntds.org/reports/5th-report/>
 16. Turner HC, Bettis AA, Chu BK, McFarland DA, Hooper PJ, et al. (2016) Investment success in public health: an analysis of the cost-effectiveness and cost-benefit of the global programme to eliminate lymphatic filariasis. *Clin Infect Dis* 64: 728–735.
 17. Okoyo C, Nikolay B, Kihara J, Simiyu E, Garn JV, et al. (2016) Monitoring the impact of a national school based deworming programme on soil-transmitted helminths in Kenya: the first three years, 2012–2014. *Parasit Vectors* 9: 408. <https://doi.org/10.1186/s13071-016-1679-y> PMID: 27457129
 18. Ortu G, Assoum M, Wittmann U, Knowles S, Clements M, et al. (2016) The impact of an 8-year mass drug administration programme on prevalence, intensity and co-infections of soil-transmitted helminthiasis in Burundi. *Parasit Vectors* 9: 513. <https://doi.org/10.1186/s13071-016-1794-9> PMID: 27660114
 19. Adriko M, Tinkitina B, Arinaitwe M, Kabatereine NB, Nanyunja M, et al. (2018) Impact of a national deworming campaign on the prevalence of soil-transmitted helminthiasis in Uganda (2004–2016): implications for national control programs. *PLoS Negl Trop Dis* 12: e0006520. <https://doi.org/10.1371/journal.pntd.0006520> PMID: 29975696
 20. World Health Organization: Regional Office for Africa (2018) Expanded Special Project for Elimination of Neglected tropical diseases (ESPEN). <http://espen.afro.who.int/>
 21. WHO (2017) Schistosomiasis and soil-transmitted helminthiasis: number of people treated in 2016. *Wkly Epidemiol Rec* 92: 749–760. PMID: 29218962
 22. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, et al. (2012) Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380: 2197–2223. [https://doi.org/10.1016/S0140-6736\(12\)61689-4](https://doi.org/10.1016/S0140-6736(12)61689-4) PMID: 23245608
 23. GBD 2016 DALYs and HALE Collaborators (2017) Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and

- territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390: 1260–1344. [https://doi.org/10.1016/S0140-6736\(17\)32130-X](https://doi.org/10.1016/S0140-6736(17)32130-X) PMID: 28919118
24. Marocco C, Bangert M, Joseph SA, Fitzpatrick C, Montresor A (2017) Preventive chemotherapy in one year reduces by over 80% the number of individuals with soil-transmitted helminthiases causing morbidity: results from meta-analysis. *Trans R Soc Trop Med Hyg* 111: 12–17. <https://doi.org/10.1093/trstmh/trx011> PMID: 28340144
 25. Montresor A, Trouleau W, Mupfasoni D, Bangert M, Joseph SA, et al. (2017) Preventive chemotherapy to control soil-transmitted helminthiasis averted more than 500 000 DALYs in 2015. *Trans R Soc Trop Med Hyg* 111: 457–463. <https://doi.org/10.1093/trstmh/trx082> PMID: 29346640
 26. Truscott JE, Turner HC, Farrell SH, Anderson RM (2016) Soil-transmitted helminths: mathematical models of transmission, the impact of mass drug administration and transmission elimination criteria. *Adv Parasitol* 94: 133–198. <https://doi.org/10.1016/bs.apar.2016.08.002> PMID: 27756454
 27. Coffeng LE, Nery SV, Gray DJ, Bakker R, de Vlas SJ, et al. (2018) Predicted short and long-term impact of deworming and water, hygiene, and sanitation on transmission of soil-transmitted helminths. *PLoS Negl Trop Dis* 12: e0006758. <https://doi.org/10.1371/journal.pntd.0006758> PMID: 30522129
 28. Anderson R, Truscott J, Hollingsworth TD (2014) The coverage and frequency of mass drug administration required to eliminate persistent transmission of soil-transmitted helminths. *Philos Trans R Soc Lond B Biol Sci* 369: 20130435. <https://doi.org/10.1098/rstb.2013.0435> PMID: 24821921
 29. Moser W, Schindler C, Keiser J (2017) Efficacy of recommended drugs against soil transmitted helminths: systematic review and network meta-analysis. *BMJ* 358: j4307. <https://doi.org/10.1136/bmj.j4307> PMID: 28947636
 30. Bethony J, Chen J, Lin S, Xiao S, Zhan B, et al. (2002) Emerging patterns of hookworm infection: influence of aging on the intensity of *Necator* infection in Hainan province, People's Republic of China. *Clin Infect Dis* 35: 1336–1344. <https://doi.org/10.1086/344268> PMID: 12439796
 31. Taylor MJ, Hoerauf A, Bockarie M (2010) Lymphatic filariasis and onchocerciasis. *Lancet* 376: 1175–1185. [https://doi.org/10.1016/S0140-6736\(10\)60586-7](https://doi.org/10.1016/S0140-6736(10)60586-7) PMID: 20739055
 32. UNESCO Institute for Statistics (2018) Out-of-school children of primary school age. In: UNESCO, editor. <http://data.uis.unesco.org/>.
 33. WHO (2017) Report of the WHO advisory group on deworming in girls and women of reproductive age. Rockefeller Foundation Bellagio Center, Bellagio, Italy, 28–30 June 2017. World Health Organization.
 34. Geerts S, Gryseels B (2000) Drug resistance in human helminths: current situation and lessons from livestock. *Clin Microbiol Rev* 13: 207–222. PMID: 10755998
 35. Vercruyse J, Levecke B, Prichard R (2012) Human soil-transmitted helminths: implications of mass drug administration. *Curr Opin Infect Dis* 25: 703–708. <https://doi.org/10.1097/QCO.0b013e328358993a> PMID: 22964945
 36. Papaïakovou M, Pilotte N, Williams SA. Recent advances in developing an optimized qPCR system for STH detection in stool. Soil-Transmitted Helminthiasis Advisory Committee Meeting, 2017; Baltimore, MD.
 37. Vlamincq J, Cools P, Albonico M, Ame S, Ayana M, et al. (2018) Comprehensive evaluation of stool-based diagnostic methods and benzimidazole resistance markers to assess drug efficacy and detect the emergence of anthelmintic resistance: a Starworms study protocol. *PLoS Negl Trop Dis* 12: e0006912. <https://doi.org/10.1371/journal.pntd.0006912> PMID: 30388108
 38. Bergquist R, Johansen MV, Utzinger J (2009) Diagnostic dilemmas in helminthology: what tools to use and when? *Trends Parasitol* 25: 151–156. <https://doi.org/10.1016/j.pt.2009.01.004> PMID: 19269899
 39. Utzinger J, Becker SL, van Lieshout L, van Dam GJ, Knopp S (2015) New diagnostic tools in schistosomiasis. *Clin Microbiol Infect* 21: 529–542. <https://doi.org/10.1016/j.cmi.2015.03.014> PMID: 25843503
 40. Bärenbold O, Raso G, Coulibaly JT, N'Goran EK, Utzinger J, et al. (2017) Estimating sensitivity of the Kato-Katz technique for the diagnosis of *Schistosoma mansoni* and hookworm in relation to infection intensity. *PLoS Negl Trop Dis* 11: e0005953. <https://doi.org/10.1371/journal.pntd.0005953> PMID: 28976979
 41. Lim MD, Brooker SJ, Belizario VY Jr, Gay-Andrieu F, Gilleard J, et al. (2018) Diagnostic tools for soil-transmitted helminths control and elimination programs: a pathway for diagnostic product development. *PLoS Negl Trop Dis* 12: e0006213. <https://doi.org/10.1371/journal.pntd.0006213> PMID: 29494581
 42. Palmeirim MS, Hürlimann E, Knopp S, Speich B, Belizario V Jr, et al. (2018) Efficacy and safety of co-administered ivermectin plus albendazole for treating soil-transmitted helminths: a systematic review meta-analysis and individual patient data analysis. *PLoS Negl Trop Dis* 12: e0006458. <https://doi.org/10.1371/journal.pntd.0006458> PMID: 29702653

43. Montresor A (2016) 21st WHO expert committee on the selection and use of essential medicines: application for inclusion of ivermectin on the WHO model list of essential medicines (EML) and model list of essential medicines for children (EMLE). Geneva: World Health Organization.
44. Lancet Editorial Board (2012) Progress in sanitation needed for neglected tropical diseases. *Lancet* 379: 978.
45. WHO (2015) Water sanitation & hygiene for accelerating and sustaining progress on neglected tropical diseases: a global strategy 2015–2020. Geneva: World Health Organization.
46. Ziegelbauer K, Speich B, Mäusezahl D, Bos R, Keiser J, et al. (2012) Effect of sanitation on soil-transmitted helminth infection: systematic review and meta-analysis. *PLoS Med* 9: e1001162. <https://doi.org/10.1371/journal.pmed.1001162> PMID: 22291577
47. Strunz EC, Addiss DG, Stocks ME, Ogden S, Utzinger J, et al. (2014) Water, sanitation, hygiene, and soil-transmitted helminth infection: a systematic review and meta-analysis. *PLoS Med* 11: e1001620. <https://doi.org/10.1371/journal.pmed.1001620> PMID: 24667810
48. Waite RC, Woods G, Velleman Y, Freeman MC (2017) Collaborating to develop joint water, sanitation and hygiene (WASH) and neglected tropical disease (NTD) sector monitoring: an expert consultation. *Int Health* 9: 215–225. <https://doi.org/10.1093/inthealth/ihx008> PMID: 28407112
49. Tomczyk S, Deribe K, Brooker SJ, Clark H, Rafique K, et al. (2014) Association between footwear use and neglected tropical diseases: a systematic review and meta-analysis. *PLoS Negl Trop Dis* 8: e3285. <https://doi.org/10.1371/journal.pntd.0003285> PMID: 25393620
50. Nakagawa J, Ehrenberg JP, Nealon J, Fürst T, Aratchige P, et al. (2015) Towards effective prevention and control of helminth neglected tropical diseases in the Western Pacific Region through multi-disease and multi-sectoral interventions. *Acta Trop* 141: 407–418. <https://doi.org/10.1016/j.actatropica.2013.05.010> PMID: 23792012