

Kuche Desalegn (Orcid ID: 0000-0001-7118-8352)

Factors associated with dietary diversity and length-for-age z-score in rural Ethiopian children aged 6-23 months: a novel approach to the analysis of baseline data from the Sustainable Undernutrition Reduction in Ethiopia evaluation

Desalegn Kuche\*, Cami Moss†, Solomon Eshetu\*, Girmay Ayana\*, Mihretab Salasibew†, Alan D. Dangour†, Elizabeth Allen†

\*Food Science and Nutrition Research Directorate, Ethiopian Public Health Institute, Addis Ababa, Ethiopia. †London School of Hygiene and Tropical Medicine, London, UK.

### **Acknowledgements**

We gratefully acknowledge the Tigray, Amhara, Oromiya and SNNP regional, zonal, district health bureaus and offices for facilitating household data collection. We thank the supervisors, enumerators and EPHI staff who collected data, and we also extend our gratitude to Mrs. Alemnesh Petros for supporting data cleaning and Mrs. Beza Teshome for managing and cleaning data. Finally, but most importantly, we would like to thank all the mothers and children who participated in the study.

### **Source of funding**

This study was supported by the Children's Investment Fund Foundation (grant: G-1610-01691). The funders did not participate in the study design, data collection or analysis, review of the manuscript or decision to publish.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/mcn.12852

## **Conflict of interest**

The authors declare that they have no conflict of interest.

## **Contributors**

DK, CM, MS, AD and EA conceived the idea and designed the study. DK, SE, GA, MS and CM coordinated and supervised training and data collection. DK, GA, SE, CM, MS and EA analysed the data and interpreted. DK and CM wrote the first draft of the manuscript. AD, CM, MS and EA critically reviewed the manuscript. EA provided technical support in analysis. All authors reviewed and approved the manuscript.

The word count of abstract is 248.

The word count of main body of the text is 3649.

A total of 54 references were included

One figure and 3 tables were included

## Abstract

Infants and young children need diets high in nutrient density and diversity to meet the requirements of rapid growth and development. Our aim was to evaluate sociodemographic, agricultural diversity, and women's empowerment factors associated with child dietary diversity and length-for-age z-score (LAZ) in children 6-23 months using data collected as part of the Sustainable Undernutrition Reduction in Ethiopia (SURE) evaluation study baseline survey in May-June 2016. We here present a novel analysis using Directed Acyclic Graphs (DAGs) to represent our assumptions about the causal influences between the factors of interest and the outcomes. The causal diagrams enabled the identification of variables to be included in multivariable analysis to estimate the total effects of factors of interest using ordinal logistic/linear regression models. We found that child dietary diversity was positively associated with LAZ with children consuming 4 or more food groups having on average an LAZ score 0.42 [95% CI 0.08, 0.77] higher than those consuming no complementary foods. Household production of fruits and vegetables was associated with both increased child dietary diversity (adjusted OR 1.16 95% CI [1.09, 1.24]) and LAZ (adjusted mean difference 0.05 [95% CI 0.005, 0.10]). Other factors positively associated with child dietary diversity included age in months, socioeconomic status, maternal education, women's empowerment and dietary diversity, paternal childcare support, household food security, fruit and vegetable cultivation, and land ownership. LAZ was positively associated with age, socioeconomic status, maternal education, fruit and vegetable production, and land ownership.

*Key words:* Nutrition, dietary diversity, length-for-age, causal diagram

## Introduction

Infants and young children require diets high in nutrient density and variety to meet the requirements of rapid growth and development (Black et al., 2013). Growth failure is most pronounced between ages 6 to 23 months of age (Shrimpton et al., 2001) and stunting – or low length-for-age (LAZ) – is associated with poorer cognitive development, weakened immune systems and increased risk of chronic disease later in life (Dewey & Begum, 2011). Globally only 29% of children 6-23 months consume the minimum dietary diversity (defined by WHO as  $\geq 4$  of 7 food groups) (UNICEF, 2016) and an estimated 23% of children under five are stunted (UNICEF, 2017).

Dietary diversity is associated with stunting in multiple low-income settings characterised by high prevalence of undernutrition (Arimond & Ruel, 2004). In Ethiopia, children's diets are among the least diverse in the world (12.5% of children 6-23 months consume the minimum dietary diversity) (UNICEF, 2018) and 37% of children under five are stunted (Central Statistical Agency, 2016). The Federal Government of Ethiopia has committed to improving the nutritional status of children, developing the third National Nutrition Programme (2016-2020) to drive policy actions across multiple key sectors including health, agriculture, education and business (Federal Democratic Republic of Ethiopia, 2016). Infant and young child feeding (IYCF) practices are highlighted and the plan calls for the cross-sectoral delivery of nutrition-sensitive interventions to improve child dietary diversity and, consequently, reduce stunting.

In support of these objectives, the Government designed the Sustainable Undernutrition Reduction in Ethiopia (SURE) programme, a multi-sectoral intervention that aims to integrate the work of the health and agriculture sectors to improve child feeding and reduce stunting. Household counselling and participatory community events are delivered by health and

agriculture extension workers to improve infant and young child feeding (IYCF) and nutrition-sensitive agriculture practices; and multi-sectoral governance structures are supported at district and *kebele* (sub-district) levels. Since June 2017, SURE has been implemented in 50 districts in the four agrarian regions of Ethiopia (Amhara, Oromia, Tigray and SNNP) and reaches approximately 800,000 children aged 6 to 23 months.

Risk factors for low LAZ or stunting have been studied in settings around the world including Ethiopia (Danaei et al., 2016), but context-specific risk factors for dietary diversity are less understood as is the contribution of dietary diversity to linear growth. In particular, agricultural diversity and women's empowerment – factors central to the SURE programme evaluation – and their respective relationships to child diets and growth have been less researched and thus comprise a particular focus of this research. Using data from the baseline survey, this study uses novel methods – construction of directed acyclic graphs to diagram causal relationships – to evaluate the association between child dietary diversity and LAZ in agrarian regions of Ethiopia and to identify risk factors for each of the two nutrition-orientated outcomes.

## **Methods**

### **Study design**

Evaluation of the SURE intervention uses a quasi-experimental study design to determine the impact on child minimum acceptable diet (MAD) and stunting. This study analyses data from the SURE baseline survey completed in May-July 2016 and comprising 1,848 children 6-23 months of age in 36 intervention districts (Oromia: n=18; Amhara: n=8; Tigray: n=4; and SNNP: n=6) and 36 comparison districts selected by region in equal proportion. Complete details are available in the SURE evaluation study protocol (Moss et al., 2018).

## Sampling

At baseline 4,980 children 0-47 months (761 children 0-5 months; 1,848 children 6-23 months; 2,371 children 24-47 months) were selected from 4,299 households. Sample size calculations were based on detecting a change at endline in length/height-for-age (LAZ) score and MAD attributable to the intervention. Detectable differences of 0.15 HAZ and of 4% MAD were calculated for intra-cluster correlation coefficients (ICC) of 0.03 with 80% power with a significance level of 5%; and differences of 0.21 HAZ and 6% MAD were calculated for ICC of 0.08 with 80% power with a significance level of 5%.

*Kebeles* were selected at study outset using probability proportional to size (PPS) sampling from lists and population data provided by district officials. *Gotes* (sub-*kebeles*) were selected by simple random sampling (paper in hat) during data collection. A complete listing of all households with children under 47 months in the *gote* was conducted and 15 were selected using systematic random sampling. Resident children 0-47 months within a selected household were listed in the following age groups: 0-5 months, 6-23 months and 24-47 months. Where only one child for any/all age categories was present, all eligible children were selected (up to three children). Where multiple children from a single age category were present, one child was randomly selected per age group by the computer-assisted personal interview (CAPI) programme. This study uses data from children sampled who were between 6-23 months of age only as this is the age group for which WHO complementary feeding indicators apply and for whom dietary data were collected.

## **Questionnaire and anthropometry**

The household questionnaire (see Supplementary File 1) comprised modules on child feeding and care practices, child anthropometry and Haemoglobin (Hb), household characteristics including food security, mother's dietary diversity, agricultural practices including diversity of food production, and women's empowerment. Child anthropometric measurements comprising length were taken using a portable measuring board (UNICEF Supply Division, Copenhagen, 2016). Data collectors were trained in anthropometric measurement for 5 days and completed a standardisation exercise prior to survey deployment. Survey training on the questionnaire, operating procedures and piloting was completed in 12 days for a total of 17 days of training.

## **Data management**

### *Length-for-age*

Child length-for-age z-scores were generated using the WHO growth standards (WHO Multicentre Growth Reference Study Group & de Onis, 2006). Scores of  $<-6$  or  $>6$  LAZ were excluded for biological implausibility as per WHO guidelines (World Health Organisation, 2006).

### *Dietary data and food insecurity*

Dietary diversity was first generated for children as a score ranging from 0-7 food groups as defined by WHO indicators for infant and young child feeding (WHO et al., 2010). We then combined 4-7 food groups consumed into a single category to create a 5-category child dietary diversity outcome variable: 0, 1, 2, 3, and 4-7 food groups consumed. Women's minimum dietary diversity (MDD-W) was generated as a scale variable ranging from 0-10 food groups as defined by FAO/FANTA (Food and Agriculture Organisation & USAID's Food and Nutrition Technical Assistance III Project (FANTA), 2016). A scale variable from 0-27 based on the Household Food Insecurity Access Score (HFIAS) was also used (Coates, Swindale, &

Bilinsky, 2007). All dietary data were cleaned by comparing 24-hour recall foods first entered in CAPI questionnaire as free text with final food group assignments made by data collectors.

#### *Socio-demographic variables*

A household wealth index was created using principle components analysis (PCA) applied to proxy indicators of household socio-economic status, namely, ownership of consumer goods, electricity, livestock (non-food producing), source of water, type of toilet, and type of materials used for floor, roof and walls). We created tertiles and checked internal validity by assessing ownership of consumer goods and housing characteristics by SES tertile. Maternal education, land ownership, and ownership of livestock producing animal sources foods such as cows or sheep were excluded from the index due to known effects on nutrition outcomes that we wished to explore independently.

#### *Agricultural production*

Variables for household food production were constructed from crops grown, animals reared, and resulting food types produced by the household within the past major and minor growing seasons (one-year reference period). We constructed a variable for fruit and vegetable production as the sum of all distinct types of fruit and vegetable crops grown in the past year. The score ranged from 0 (i.e. no fruit and vegetable crops grown) to a maximum of 21 as reported by the household. A variable for animal source food production was constructed as the sum of all such food types produced on a scale of 0-10 including eggs, meats, milks and other dairy, and fish.

#### **Statistical analyses**

Descriptive statistics were generated for child, mother, and household risk factors and outcomes. Continuous variables were summarised using means and standard deviations or

medians and inter-quartile ranges for non-normally distributed variables. Categorical variables were summarised using numbers and percentages.

We hypothesised pathways of impact between risk factors and dietary diversity and linear growth in children. To represent these pathways, a directed acyclic graph (DAG) was developed using DAGitty software (Textor, van der Zander, Gilthorpe, Liškiewicz, & Ellison, 2016) (see **Figure 1**). DAGs represent a set of assumptions about the causal relationships between variables that are made by researchers based on evidence and logical reasoning and provide a basis on which to reduce bias in statistical modelling (Sauer & J Vanderweele, 2013; Shrier & Platt, 2008). Recent evidence on agricultural production and women's empowerment factors supported construction of our diagram (Cunningham, Ruel, Ferguson, & Uauy, 2015; Hirvonen & Hoddinott, 2016; Ruel & Alderman, 2013). Occurrence of fever and diarrhoea in the past two weeks were hypothesised to influence child feeding practices but not growth due to the limited reference period.

Based on ancestor variables for each individual association, we used the DAGitty software to identify the unique set of covariates to be included in a multivariate model of that association. Unlike stepwise regression, covariates were not forced into a single multivariate model thus avoiding the table 2 fallacy (i.e. presentation of multiple adjusted effect estimates from a single model in a single table, often misunderstood to represent effects of the same direct causal type and therefore misinterpreted) (Westreich & Greenland, 2013). We also decided *a priori* to add age and sex into all multivariate models, and age squared into multivariate model for LAZ. Associations between factors and child dietary diversity were investigated using univariate and multivariate ordinal logistic regression modelling. Associations between factors and LAZ were investigated using univariate and multivariate linear regression modelling. Robust standard errors were used to account for clustering at the *kebele* level. All analyses were conducted using Stata/IC (version 15).

## Ethics

The baseline protocol was approved by the Scientific and Ethical Review Committee (SERC) at EPHI (Ref number: SERO-54-3-2016) and by the LSHTM ethics committee (Ref number: 10937) prior to data collection. By approval of both institutional review boards, informed written consent was given by caregivers of young children and witnessed by the local health extension worker.

## Results

A total of 1,848 children 6-23 months and their respective caregivers participated in the study. Of children 6-23 months of age, 6.0% consumed no complementary foods (breastfed only), 16.6% ate 1 food group, 31.1% ate 2 groups, 26.9% ate 3 groups, and 19.5% ate  $\geq 4$  groups.

The number of food groups consumed increased with child age; the mean (SD) age of those children consuming no complementary foods was 11.1 (4.5) months, and 15.7 (4.7) months among those consuming  $\geq 4$  food groups. Boys had lower LAZ (-1.23; SD 1.6) compared to girls (-1.05; SD 1.5), as did those of lower socioeconomic status (-1.29; SD 1.6) compared to those of higher status (-0.99; SD 1.5). Median maternal education was 3 years [IQR: 0, 7] for those children who consumed  $\geq 4$  food groups in comparison to 0 years of education [IQR: 0, 2] among those who consumed 0 food groups. Children with fathers who supported childcare daily or weekly consumed a higher number of food groups to those with less frequent paternal support. Complete socioeconomic and demographic characteristics of the participants are described in **Table 1**.

### *Dietary diversity*

In adjusted models, children from the highest socioeconomic group had increased odds (adjusted OR 1.50; 95% CI [1.16, 1.93];  $p=0.004$ ) of high dietary diversity compared to those in the lowest group (see **Table 2**). Other demographic factors positively associated with child

dietary diversity included increase in age in months and land ownership. Cultivation of each additional type of fruit or vegetable within the household was associated with 16% higher odds of achieving high dietary diversity (adjusted OR 1.16 [1.09, 1.25],  $p < 0.001$ ), and 6% higher odds for production of each animal source food (adjusted OR 1.06 [1.00, 1.13];  $p = 0.06$ ). Positive associations were also identified between dietary diversity and gender-related variables including years of maternal education, women's empowerment (major decision-making power), and paternal childcare support.

### *Length-for-age*

Child dietary diversity was positively associated with LAZ: children who consumed the highest number of food groups (between 4-7) had 0.42 higher mean LAZ (95% CI [0.08, 0.77];  $p = 0.006$ ) compared to those children who consumed no complementary foods (see **Table 3**).

Girls had better growth compared to boys and other demographic predictors of LAZ included age, socioeconomic status, and land ownership. Household cultivation of fruit and vegetables was positively associated with LAZ (adjusted coefficient 0.05 [0.01, 0.10],  $p = 0.03$ ). Weak evidence suggests that animal source foods may be negatively associated with growth. Maternal education was linked to higher LAZ, but no evidence of association was identified between LAZ and women's empowerment or father's support for childcare.

### **Discussion**

In this study, we used novel methods – construction of directed acyclic graphs to diagram causal relationships – to more precisely identify and estimate factors associated with child dietary diversity and LAZ among children 6-23 months old in rural agrarian regions of Ethiopia. Child dietary diversity was itself associated with increased LAZ. Socioeconomic status, maternal education, land ownership, and cultivation of fruits and vegetables were also positively associated with LAZ, whilst weak evidence suggested that animal source food

production at household level may be a risk factor for decreased LAZ. Factors positively associated with child dietary diversity included age, socioeconomic status, land ownership, maternal education, women's empowerment, partner support for childcare, food security, household fruit and vegetable production and women's dietary diversity.

The association between dietary diversity and linear growth in rural agrarian settings is well-established (Dickson A Amugsi, Dimbuene, Kimani-Murage, Mberu, & Ezeh, 2017; Borkotoky, Unisa, & Gupta, 2018; Busert et al., 2016) and consumption of at least four of seven food groups (WHO/UNICEF) is an indicator of micronutrient adequacy (WHO et al., 2010). In this study we found that growth was improved among children consuming two food groups or more compared with 0-1 food groups. The benefits from improved dietary diversity even at the very low levels may be partly explained by widespread challenges in Ethiopia with delayed initiation of complementary feeding, as observed in this study and in national statistics (51% of children 6-8 months do not yet consume complementary foods) (Central Statistical Agency, 2016), and by high prevalence of religious fasting which prohibits adherents from eating animal source foods for more than 200 days per year (Desalegn, Lambert, Riedel, Negese, & Biesalski, 2018). Diversity of the diet also increased with child age in months, indicating that children of complementary feeding age were slow to begin consuming a wider range of food groups.

Access to a variety of foods is a prerequisite to good child dietary diversity. Our findings also confirm evidence from multiple studies showing that quality of children's diets worsens as food insecurity becomes more severe (Rodriguez, Mundo-Rosas, Mendez-Gomez-Humaran, Perez-Escamilla, & Shamah-Levy, 2017). In the Ethiopian context, in which meals are largely shared, maternal dietary diversity may also partly serve as a proxy indicator of household access to diverse foods. A study conducted in Ghana showed that as the number of food groups

consumed by mother increased, so too did the number consumed by their children (Dickson Abanami Amugsi, Mittelmark, & Oduro, 2015). Low socioeconomic status has been identified as a risk factor for low child dietary diversity in a wide range of contexts (Karwa, Godhia, & Jadhav, 2016; Rakotonirainy et al., 2018) and associations with stunted growth are well-established (Adekanmbi, Kayode, & Uthman, 2013; Devakumar et al., 2018; Leroy, Habicht, Gonzalez de Cossio, & Ruel, 2014; Poda, Hsu, & Chao, 2017). Similarly, land ownership was also an important protective factor for child feeding practices in this study and others (Devakumar et al., 2018; Hailemariam, Girmay, & Girmay, 2018), perhaps due to the central role of agriculture in ensuring food security via food provision and/or income.

Agriculture-related characteristics and practices may increase access to and use of nutrient-rich food. We found strong evidence of an association between the cultivation of fruits and vegetables within the household and increases in both child dietary diversity and LAZ. Consumption of fruits and vegetables improves micronutrient status and immune function and may support child growth (Aguayo, Nair, Badgaiyan, & Krishna, 2016). A previous study in Ethiopia also reported that higher vegetable and fruit dietary consumption is associated with increased LAZ and reduced risk of stunting (Melaku et al., 2018), consistent with findings from Nepal and northern Ghana (Mulmi et al., 2017; Saaka, Osman, & Hoeschle-Zeledon, 2017).

By contrast, we found weak suggestion of an inverse association between LAZ and animal source food production at household level. Poor environmental hygiene related to livestock husbandry has been documented in Ethiopia and in other developing contexts (Mosites et al., 2015). Keeping poultry in the household dwelling overnight, a common local practice, has been found to be associated with decreased child growth (Headey & Hirvonen, 2016). Despite some evidence from our study that production of animal source foods increases child dietary diversity, interventions that promote livestock rearing may need to include strong animal hygiene education and child feeding counselling to ensure that – in the Ethiopian social context

– animal source foods are given with sufficient consistency and quantity to support child growth.

We investigated a range of gender-related risk factors. Maternal education is a well-established predictor of child feeding practices and nutritional status (Boyle et al., 2006; Wachs, Creed-Kanashiro, Cueto, & Jacoby, 2005; Wang et al., 2017). A study conducted in Addis Ababa, Ethiopia reported that mother's attainment of secondary education and above was associated with higher odds of achieving minimum dietary diversity among children 6–23 months old (Solomon, Aderaw, & Tegegne, 2017). Women's empowerment – measured in this study by women's power to decide major household expenses – was associated with child dietary diversity but not LAZ, consistent with other findings linking women's participation in financial decisions and quality of child diets (Dickson A Amugsi, Lartey, Kimani-Murage, & Mberu, 2016; Beyene, Worku, & Wassie, 2015). We also found that paternal childcare support predicted improved child diet quality. Mechanisms may include increased time for mothers to prepare food or increased prioritisation of children's needs when drawing on household resources. Ethiopian fathers' knowledge and support for child feeding practices has been shown to be positively associated with child dietary diversity (Bilal et al., 2016).

### *Strengths and limitations*

In this study, we have extended investigation of factors beyond those commonly known or hypothesised to be associated with child diets and nutritional status to include some less-researched factors: household agricultural diversity and gender-related norms. Our analyses were based on a large dataset collected as primary data, and we used causal diagrams to represent our assumptions about the causal influences between the factors of interest and the outcomes. We then used these diagrams to identify variables to include in multivariable analysis. This approach allowed for improved estimates of the multiple primary associations

between individual factors of interest and outcomes; we identified and adjusted for the unique set of covariates exerting direct or indirect effects on an individual association as per the causal relationships specified. This is an advance on ‘single model single table’ multivariate modelling which displays – as a result of modelling a single primary association – figures representing a mixture of different types of causal effect estimates for covariates (secondary, direct and indirect) without explicit distinction or comparability between them (Westreich & Greenland, 2013).

Data was obtained from a single cross-sectional study and therefore evidence from our analyses is limited by underlying assumptions of temporality. Dietary data may have been subject to recall bias and may not have shown usual intake as it was based on one 24-hour recall. The data were also collected over a period inclusive both of the Lent fasting period before Easter and the meat-heavy period directly following Easter. Other seasonal variation in the diet was not captured. Certain factors known to be associated with stunting such as child birthweight and household sanitation were not included in this study.

#### *Policy implications and further research*

We showed that child diets in rural, agrarian regions of Ethiopia may be improved with increased household production of fruits and vegetables and of animal source foods. Interventions that seek to diversify farms such as agriculture extension advising, homestead gardening promotion and/or provision of seed or animal inputs may be effective (Hirvonen & Headey, 2018; Sibhatu, Krishna, & Qaim, 2015). Other means of increasing access to and use of diverse foods might include market-based initiatives or voucher/cash interventions combined with nutrition education and promotion (Ruel & Alderman, 2013). Interventions to increase livestock rearing at household level may help to improve dietary diversity. Further

research is needed on the complex relationships between livestock rearing, hygiene practices, consumption of animal source foods and child growth (Mosites et al., 2015).

Child feeding was also linked to paternal childcare support and women's empowerment, suggesting that inclusion of husbands and other community members in nutrition education initiatives – as part of a broader social and behaviour change communication strategy – may be effective to improve child diets (Nguyen et al., 2018). However, strategies to effectively engage men and community influencers are not yet well researched (Lutter et al., 2013). Education of girls remains essential to sustained improvement in child growth in Ethiopia (Lailulo, Sathiya Susuman, & Blignaut, 2015).

## **Conclusions**

Young children in Ethiopia are slow to begin consuming a range of diverse complementary foods. Household cultivation of fruits and vegetables in rural agrarian settings can improve both child feeding practice and linear growth. Gender-sensitive interventions to increase maternal education, women's empowerment and paternal support for childcare may support improved diversity of child feeding.

## **Key messages**

- Young children in Ethiopia are slow to begin consuming a range of diverse complementary foods.
- Child dietary diversity is positively associated with linear growth; and household production of fruits and vegetables is positively associated with child dietary diversity and length-for-age-z score.
- Gender-sensitive interventions to increase maternal education, women's empowerment and paternal support for childcare may support improved diversity of child feeding.

## References

- Adekanmbi, V. T., Kayode, G. A., & Uthman, O. A. (2013). Individual and contextual factors associated with childhood stunting in Nigeria: a multilevel analysis. *Matern Child Nutr*, 9(2), 244-259.
- Aguayo, V. M., Nair, R., Badgaiyan, N., & Krishna, V. (2016). Determinants of stunting and poor linear growth in children under 2 years of age in India: an in-depth analysis of Maharashtra's comprehensive nutrition survey. *Matern Child Nutr*, 12, 121-140.
- Amugsi, D. A., Dimbuene, Z. T., Kimani-Murage, E. W., Mberu, B., & Ezeh, A. C. (2017). Differential effects of dietary diversity and maternal characteristics on linear growth of children aged 6–59 months in sub-Saharan Africa: a multi-country analysis. *Public health nutrition*, 20(6), 1029-1045.
- Amugsi, D. A., Lartey, A., Kimani-Murage, E., & Mberu, B. U. (2016). Women's participation in household decision-making and higher dietary diversity: findings from nationally representative data from Ghana. *Journal of Health, Population and Nutrition*, 35(1), 16.
- Amugsi, D. A., Mittelmark, M. B., & Oduro, A. (2015). Association between maternal and child dietary diversity: an analysis of the Ghana demographic and health survey. *PloS one*, 10(8), e0136748.
- Arimond, M., & Ruel, M. T. (2004). Dietary Diversity Is Associated with Child Nutritional Status: Evidence from 11 Demographic and Health Surveys. *The Journal of Nutrition*, 134(10), 2579-2585. doi: 10.1093/jn/134.10.2579
- Beyene, M., Worku, A. G., & Wassie, M. M. (2015). Dietary diversity, meal frequency and associated factors among infant and young children in Northwest Ethiopia: a cross-sectional study. *BMC public health*, 15(1), 1007.

- Bilal, S. M., Dinant, G., Blanco, R., Crutzen, R., Mulugeta, A., & Spigt, M. (2016). The influence of father's child feeding knowledge and practices on children's dietary diversity: a study in urban and rural districts of Northern Ethiopia, 2013. *Matern Child Nutr*, 12(3), 473-483.
- Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., De Onis, M., . . . Martorell, R. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries. *The lancet*, 382(9890), 427-451.
- Borkotoky, K., Unisa, S., & Gupta, A. K. (2018). State-level dietary diversity as a contextual determinant of nutritional status of children in India: A multilevel approach. *Journal of biosocial science*, 50(1), 26-52.
- Boyle, M. H., Racine, Y., Georgiades, K., Snelling, D., Hong, S., Omariba, W., . . . Rao-Melacini, P. (2006). The influence of economic development level, household wealth and maternal education on child health in the developing world. *Social science & medicine*, 63(8), 2242-2254.
- Busert, L. K., Neuman, M., Rehfuess, E. A., Dulal, S., Harthan, J., Chaube, S. S., . . . Manandhar, D. S. (2016). Dietary Diversity Is Positively Associated with Deviation from Expected Height in Rural Nepal. *J Nutr*, 146(7), 1387-1393.
- Central Statistical Agency. (2016). Ethiopia Demographic and Health Survey 2016. Addis Ababa: Federal Democratic Republic of Ethiopia.
- Coates, J., Swindale, A., & Bilinsky, P. (2007). Household Food Insecurity Access Scale (HFIAS) for Measurement of Food Access: Indicator Guide. Version 3. Washington, DC: Food and Nutrition Technical Assistance III Project (FANTA).
- Cunningham, K., Ruel, M., Ferguson, E., & Uauy, R. (2015). Women's empowerment and child nutritional status in South Asia: a synthesis of the literature. *Maternal & Child Nutrition*, 11(1), 1-19. doi: 10.1111/mcn.12125

- Danaei, G., Andrews, K. G., Sudfeld, C. R., Fink, G., McCoy, D. C., Peet, E., . . . Fawzi, W. W. (2016). Risk Factors for Childhood Stunting in 137 Developing Countries: A Comparative Risk Assessment Analysis at Global, Regional, and Country Levels. *PLOS Medicine*, *13*(11), e1002164. doi: 10.1371/journal.pmed.1002164
- Desalegn, B. B., Lambert, C., Riedel, S., Negese, T., & Biesalski, H. K. (2018). Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia. *International Journal of Environmental Research and Public Health*, *15*(8), 1767. doi: 10.3390/ijerph15081767
- Devakumar, D., Kular, D., Shrestha, B. P., Grijalva, E., Eternod, C., Daniel, R. M., Saville, N. M., . . . Wells, J. C. (2018). Socioeconomic determinants of growth in a longitudinal study in Nepal. *Matern Child Nutr*, *14*(1), e12462.
- Dewey, K. G., & Begum, K. (2011). Long-term consequences of stunting in early life. *Matern Child Nutr*, *7 Suppl 3*, 5-18. doi: 10.1111/j.1740-8709.2011.00349.x
- Federal Democratic Republic of Ethiopia. (2016). National Nutrition Programme 2016-2020. Addis Ababa.
- Food and Agriculture Organisation, & USAID's Food and Nutrition Technical Assistance III Project (FANTA). (2016). Minimum Dietary Diversity for Women: A Guide to Measurement. Rome, Italy: Food and Agriculture Organisation (FAO).
- Hailemariam, T., Girmay, T., & Girmay, G. (2018). Determinants of individual dietary diversity score of children less than five years old in the southern zone of Tigray, Ethiopia. *African Journal of Food, Agriculture, Nutrition and Development*, *18*(1).
- Headey, D., & Hirvonen, K. (2016). Is Exposure to Poultry Harmful to Child Nutrition? An Observational Analysis for Rural Ethiopia. *PLoS One*, *11*(8), e0160590. doi: 10.1371/journal.pone.0160590

- Hirvonen, K., & Headey, D. (2018). Can governments promote homestead gardening at scale? Evidence from Ethiopia. *Global Food Security*, 19, 40-47. doi: <https://doi.org/10.1016/j.gfs.2018.09.001>
- Hirvonen, K., & Hoddinott, J. (2016). Agricultural production and children's diets: evidence from rural Ethiopia. *Agricultural Economics*. doi: 10.1111/agec.12348
- Karwa, K., Godhia, M. L., & Jadhav, A. R. (2016). Infant and Young Child-Feeding Practices, Indicators and Index, and Role of Socio-economic Status. *International Journal of Advanced Nutritional and Health Science*, 4(1), pp. 212-218.
- Lailulo, Y. A., Sathiya Susuman, A., & Blignaut, R. (2015). Correlates of gender characteristics, health and empowerment of women in Ethiopia. *BMC Women's Health*, 15, 116. doi: 10.1186/s12905-015-0273-3
- Leroy, J. L., Habicht, J. P., Gonzalez de Cossio, T., & Ruel, M. T. (2014). Maternal education mitigates the negative effects of higher income on the double burden of child stunting and maternal overweight in rural Mexico. *J Nutr*, 144(5), 765-770. doi: 10.3945/jn.113.188474
- Lutter, C. K., Iannotti, L., Creed-Kanashiro, H., Guyon, A., Daelmans, B., Robert, R., & Haider, R. (2013). Key principles to improve programmes and interventions in complementary feeding. *Maternal & Child Nutrition*, 9, 101-115. doi: 10.1111/mcn.12087
- Melaku, Y. A., Gill, T. K., Taylor, A. W., Adams, R., Shi, Z., & Worku, A. (2018). Associations of childhood, maternal and household dietary patterns with childhood stunting in Ethiopia: proposing an alternative and plausible dietary analysis method to dietary diversity scores. *Nutrition journal*, 17(1), 14.
- Mosites, E. M., Rabinowitz, P. M., Thumbi, S. M., Montgomery, J. M., Palmer, G. H., May, S., . . . Walson, J. L. (2015). The relationship between livestock ownership and child

stunting in three countries in Eastern Africa using national survey data. *PloS one*, *10*(9), e0136686.

Moss, C., Bekele, T. H., Salasibew, M. M., Sturgess, J., Ayana, G., Kuche, D., . . . Dangour, A. D. (2018). Sustainable Undernutrition Reduction in Ethiopia (SURE) evaluation study: a protocol to evaluate impact, process and context of a large-scale integrated health and agriculture programme to improve complementary feeding in Ethiopia. *BMJ Open*, *8*(7).

Mulmi, P., Masters, W. A., Ghosh, S., Namirembe, G., Rajbhandary, R., Manohar, S., . . . Webb, P. (2017). Household food production is positively associated with dietary diversity and intake of nutrient-dense foods for older preschool children in poorer families: Results from a nationally-representative survey in Nepal. *PloS one*, *12*(11), e0186765.

Nguyen, P. H., Frongillo, E. A., Sanghvi, T., Wable, G., Mahmud, Z., Tran, L. M., . . . Menon, P. (2018). Engagement of Husbands in a Maternal Nutrition Program Substantially Contributed to Greater Intake of Micronutrient Supplements and Dietary Diversity during Pregnancy: Results of a Cluster-Randomized Program Evaluation in Bangladesh. *The Journal of Nutrition*, *148*(8), 1352-1363. doi: 10.1093/jn/nxy090

Poda, G. G., Hsu, C.-Y., & Chao, J. C.-J. (2017). Factors associated with malnutrition among children < 5 years old in Burkina Faso: evidence from the Demographic and Health Surveys IV 2010. *International Journal for Quality in Health Care*, *29*(7), 901-908.

Rakotonirainy, N. H., Razafindratovo, V. r., Remonja, C. R., Rasoloarijaona, R., Piola, P., Raharintsoa, C., & Randremanana, R. V. (2018). Dietary diversity of 6-to 59-month-old children in rural areas of Moramanga and Morondava districts, Madagascar. *PloS one*, *13*(7), e0200235.

- Rodriguez, L. A., Mundo-Rosas, V., Mendez-Gomez-Humaran, I., Perez-Escamilla, R., & Shamah-Levy, T. (2017). Dietary quality and household food insecurity among Mexican children and adolescents. *Matern Child Nutr*, 13(4). doi: 10.1111/mcn.12372
- Ruel, M. T., & Alderman, H. (2013). Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *The Lancet*, 382(9891), 536-551. doi: 10.1016/s0140-6736(13)60843-0
- Saaka, M., Osman, S. M., & Hoeschle-Zeledon, I. (2017). Relationship between agricultural biodiversity and dietary diversity of children aged 6-36 months in rural areas of Northern Ghana. *Food & nutrition research*, 61(1), 1391668.
- Sauer, B., & J Vanderweele, T. (2013). Supplement 2. Use of Directed Acyclic Graphs. In P. Velentgas, N. Dreyer, P. Nourjah, S. Smith & M. Torchia (Eds.), *Developing a Protocol for Observational Comparative Effectiveness Research: A User's Guide*: Agency for Healthcare Research and Quality.
- Shrier, I., & Platt, R. W. (2008). Reducing bias through directed acyclic graphs. *BMC medical research methodology*, 8, 70-70. doi: 10.1186/1471-2288-8-70
- Shrimpton, R., Victora, C. G., de Onis, M., Lima, R. C., Blossner, M., & Clugston, G. (2001). Worldwide timing of growth faltering: implications for nutritional interventions. *Pediatrics*, 107(5), E75.
- Sibhatu, K. T., Krishna, V. V., & Qaim, M. (2015). Production diversity and dietary diversity in smallholder farm households. *Proceedings of the National Academy of Sciences of the United States of America*, 112(34), 10657-10662. doi: 10.1073/pnas.1510982112
- Solomon, D., Aderaw, Z., & Tegegne, T. K. (2017). Minimum dietary diversity and associated factors among children aged 6â€“23 months in Addis Ababa, Ethiopia. *International journal for equity in health*, 16(1), 181.

- Textor, J., van der Zander, B., Gilthorpe, M. S., Liškiewicz, M., & Ellison, G. T. H. (2016). Robust causal inference using directed acyclic graphs: the R package ‘dagitty’. *International Journal of Epidemiology*, 45(6), 1887-1894. doi: 10.1093/ije/dyw341
- UNICEF. (2016). *From the First Hour of Life: Making the Case for Improved Infant and Young Child Feeding Everywhere*. New York, NY: UNICEF.
- UNICEF. (2017). *State of the World's Children*. In UNICEF (Ed.). New York City, NY: UNICEF.
- UNICEF. (2018). *Expanded Global Databases on Infant and Young Child Feeding*. Retrieved from: <https://data.unicef.org/resources/dataset/infant-young-child-feeding/>
- Wachs, T. D., Creed-Kanashiro, H., Cueto, S., & Jacoby, E. (2005). Maternal education and intelligence predict offspring diet and nutritional status. *J Nutr*, 135(9), 2179-2186.
- Wang, A., Scherpbier, R. W., Huang, X., Guo, S., Yang, Y., Josephs-Spaulding, J., . . . Wang, Y. (2017). The dietary diversity and stunting prevalence in minority children under 3 years old: a cross-sectional study in forty-two counties of Western China. *British Journal of Nutrition*, 118(10), 840-848.
- Westreich, D., & Greenland, S. (2013). The table 2 fallacy: presenting and interpreting confounder and modifier coefficients. *Am J Epidemiol*, 177(4), 292-298. doi: 10.1093/aje/kws412
- WHO, UNICEF, IFPRI, Davis, U., USAID, FANTA, & International, M. (2010). *Indicators for assessing infant and young child feeding practices: Part II Measurement*. In World Health Organisation (Ed.). Geneva.
- WHO Multicentre Growth Reference Study Group, & de Onis, M. (2006). WHO Child Growth Standards based on length/height, weight and age. *Acta Pædiatrica*, 95, 76-85. doi: 10.1111/j.1651-2227.2006.tb02378.x

World Health Organisation. (2006). WHO Anthro 2005: Software for assessing growth and development of the world's children. In WHO (Ed.). Geneva.

Accepted Article




Figure 1: Directed Acyclic Graph (DAG) mapping causal relationships. DAGitty web-based software was used to develop the diagram and to determine the minimal adjustment variable set to estimate total effect when regressing each explanatory risk factor on the outcome of interest – in this study, dietary diversity or LAZ (Textor et al., 2016).

Accepted

Table 1. Baseline socioeconomic and demographic characteristics of children 6-23 months

(n=1,848)

	Child dietary diversity, food groups					LAZ (n=1,775)
Explanatory variable n (%)	0 n=106 (6.0)	1 n=296 (16.6)	2 n=555 (31.1)	3 n=479 (26.9)	4-7 n=347 (19.5)	Mean (std dev)
Children dietary diversity score, food groups (n=1783)						
0						-1.11 (1.5)
1						-1.18 (1.6)
2						-1.07 (1.6)
3						-1.23 (1.5)
4-7						-1.11 (1.5)
Age in months (mean, SD) (n=1848)	11.1 (4.5)	12.4 (4.7)	14.5 (4.9)	15.5 (4.8)	15.6 (4.7)	-1.14 (1.5)
Sex (n=1847)						
Male	47 (44.3)	153 (51.7)	276 (49.7)	251 (52.4)	175 (50.4)	-1.23 (1.6)
Female	59 (55.7)	143 (48.3)	279 (50.3)	228 (47.6)	171 (49.3)	-1.05 (1.5)
SES (n=1848)						
Poorest	42 (39.6)	123 (41.6)	194 (35.0)	139 (29.0)	92 (26.5)	-1.29 (1.6)
Middle	36 (34.0)	96 (32.4)	197 (35.5)	176 (36.7)	91 (26.2)	-1.14 (1.5)
Richest	28 (26.4)	77 (26.0)	164 (29.6)	164 (34.2)	164 (47.3)	-0.99 (1.5)
Maternal education, yrs (median, IQR) (n=1848)	0 [0, 2]	0 [0, 4]	0 [0, 5]	0 [0, 4]	3 [0, 7]	-1.14 (1.5)
Decision making on major household purchases (n=1848)						
Father or others only	40 (37.7)	103 (34.8)	170 (30.6)	132 (27.6)	106 (30.6)	-1.27 (1.5)
Jointly or mother only	66 (62.3)	193 (65.2)	385 (69.4)	347 (69.4)	241 (69.5)	-1.08 (1.5)
Paternal childcare support (n=1752)						
Never	10 (9.8)	26 (9.4)	56 (10.6)	38 (8.3)	12 (3.7)	-1.16 (1.7)
Rarely	35 (34.3)	94 (33.9)	189 (35.6)	156 (34.2)	105 (32.2)	-1.17 (1.6)
Weekly or everyday	57 (55.9)	157 (56.7)	286 (53.9)	262 (57.5)	209 (64.1)	-1.12 (1.5)
Paternal domestic chores (n=1752)						
Never	43 (42.2)	109 (39.4)	227 (42.8)	190 (41.7)	123 (37.7)	-1.21 (1.6)
Rarely	25 (24.5)	82 (29.6)	133 (25.1)	107 (23.5)	89 (27.3)	-1.11 (1.5)
Weekly or everyday	34 (33.3)	86 (31.1)	171 (32.2)	159 (34.9)	114 (35.0)	-1.09 (1.5)
Food insecurity score (median, IQR) (n=1848)	0 [0, 7]	1 [0, 7]	0 [0, 6]	0 [0, 5]	0 [0, 3]	-1.14 (1.5)
Women's DD score (median, IQR) (n=1848)	3 [2, 4]	3 [2, 4]	3 [3, 4]	3 [3, 4]	4 [3, 4]	-1.14 (1.5)
Animal source food types produced, household (median, IQR) (n=1848)	1 [0, 3]	1 [0, 2]	1 [0, 2]	1 [0, 3]	1 [0, 3]	-1.14 (1.5)
Fruit and vegetable types produced, household (median, IQR) (n=1848)	1 [0, 2]	0 [0, 2]	1 [0, 2]	1 [0, 2]	1 [0, 2]	-1.14 (1.5)
Hectares (median, IQR) (n=1729)	0.5 [0, 1]	0.5 [0.25, 1]	0.5 [0.25, 1]	0.5 [0.25, 1]	0.5 [0.25, 1.5]	-1.14 (1.5)
Irrigation (n=1847)						
No	85 (80.2)	263 (88.9)	489 (88.1)	417 (87.2)	292 (84.2)	-1.14 (1.5)
Yes	21 (19.8)	33 (11.2)	66 (11.9)	61 (12.8)	55 (15.9)	-1.13 (1.6)
Fertiliser (n=1848)						

No	19 (17.9)	40 (13.5)	75 (13.5)	60 (12.5)	44 (12.7)	-1.09 (1.5)
Yes	87 (82.1)	256 (86.5)	480 (86.5)	419 (87.5)	303 (87.3)	-1.15 (1.5)
Diarrhoea, past 2 weeks ( <i>n</i> =1783)						
No/Don't know	68 (64.2)	192 (64.9)	422 (76.0)	339 (70.8)	246 (70.9)	-1.15 (1.5)
Yes	38 (35.9)	104 (35.1)	133 (24.0)	140 (29.2)	101 (29.1)	-1.11 (1.6)
Fever, past two weeks ( <i>n</i> =1783)						
No/Don't know	73 (68.9)	207 (69.9)	405 (73.0)	344 (71.8)	247 (71.2)	-1.12 (1.5)
Yes	33 (31.1)	89 (30.1)	150 (27.0)	135 (28.2)	100 (28.8)	-1.18 (1.5)

Accepted Article

Table 2. Protective and risk factors for dietary diversity among children 6-23 months (n=1,848)

Variable (n, multivariate models)	Unadjusted (OR)	CIs	P-value	Adjusted (OR)	CIs	p-value
Age (months) (n=1,783)	1.11	(1.09, 1.13)	<0.001	1.11*	(1.09, 1.13)	<0.001
Sex (n=1,782)						
Male	1		0.07	1**		0.05
Female	0.96	(0.81, 1.14)		0.94	(0.79, 1.12)	
Socioeconomic status (SES) (n=1,667)						
Poorest	1		<0.001	1***		0.004
Middle	1.21	(0.98, 1.50)		1.07	(0.85, 1.35)	
Richest	1.86	(1.49, 2.33)		1.50	(1.16, 1.93)	
Maternal education (years) (n=1,782)	1.07	(1.04, 1.09)	<0.001	1.07†	(1.05, 1.10)	<0.001
Women's decision-making power – major household expenses (n=1,667)						
Husband/other	1		0.05	1‡		0.02
Jointly/self	1.22	(1.00, 1.48)		1.27	(1.03, 1.55)	
Paternal childcare support (n=1,691)						
Never	1		0.02	1†		0.02
Rarely	1.52	(1.08, 2.13)		1.52	(1.08, 2.14)	
Weekly/every day	1.61	(1.16, 2.22)		1.62	(1.17, 2.24)	
Paternal domestic chores (n=1,691)						
Never	1		0.59	1†		0.64
Rarely	1.02	(0.82, 1.29)		1.03	(0.82, 1.30)	
Weekly/every day	1.11	(0.90, 1.38)		1.11	(0.89, 1.37)	
Food insecurity score (HFIAS) (n=1,667)	0.96	(0.94, 0.98)	<0.001	0.98§	(0.96, 1.00)	0.04
Minimum women's dietary diversity (MDD-W) (n=1,782)	1.68	(1.53, 1.85)	<0.001	1.63‡	(1.48, 1.80)	<0.001
Animal source food types produced, household (n=1,667)	1.12	(1.05, 1.18)	<0.001	1.06‡	(1.00, 1.13)	0.06
Fruit and vegetable types produced, household (n=1,667)	1.16	(1.09, 1.23)	<0.001	1.16‡	(1.09, 1.24)	<0.001
Land owned, hectares (n=1,668)	1.11	(1.01, 1.21)	<0.02	1.11†	(1.02, 1.21)	0.02
Irrigation (n=1,781)						
No	1		0.44	1†		0.50
Yes	1.12	(0.84, 1.48)		1.10	(0.83, 1.46)	
Fertiliser (n=1,782)						
No	1		0.22	1†		0.17
Yes	1.18	(0.90, 1.55)		1.21	(0.92, 1.59)	
Diarrhoea, past 2 weeks (n=1,782)						
No	1		0.32			0.33
Yes	0.91	(0.75, 1.10)		0.91†	(0.75, 1.10)	
Fever, past 2 weeks (n=1,782)						
No	1		0.79			0.68
Yes	0.97	(0.80, 1.18)		0.96†	(0.79, 1.17)	

Table 2 legend

Variables adjusted for in multivariate ordinal logistic regression modelling:

\* Sex

\*\* Age

\*\*\* Age, sex, maternal education, land ownership, women's empowerment, irrigation, fertiliser, fruit and vegetables produced (FV), animal source foods produced (ASF)

† Age, sex

‡ Age, sex, SES, maternal education, land ownership, food security, irrigation, fertiliser, FV, ASF

§ Age, sex, SES, maternal education, land ownership, women's empowerment, irrigation, fertiliser, FV, ASF

‖ Age, sex, women's empowerment, food security, FV, ASF

Φ Age, sex, SES, maternal education, land ownership, women's empowerment, irrigation, fertiliser, FV (n=1,667)

□ Age, sex, SES, maternal education, land ownership, women's empowerment, irrigation, fertiliser, ASF

Accepted Article

Table 3. Protective and risk factors for length-for-age z score among children 6-23 months (n=1,848)

Variable (n, multivariate models)	Unadjusted Coefficient	CI	P-value	Adjusted Coefficient	CI	p-value
Child dietary diversity (food groups) (n=1,579)						
0	1			1*		
1	-0.11	(-0.45, 0.23)		0.08	(-0.26, 0.41)	
2	-0.02	(-0.33, 0.30)		0.42	(0.10, 0.75)	
3	-0.15	(-0.47, 0.17)		0.35	(0.02, 0.68)	
4-7	-0.05	(-0.38, 0.28)	0.64	0.42	(0.08, 0.77)	0.006
Age (in months) (n=1,775)	-0.25	(-0.34, -0.17)	<0.001	-0.25**	(-0.34, -0.16)	<0.001
Age squared (n=1,775)	0.01	(0.003, 0.01)	<0.001	0.005**	(0.002, 0.01)	<0.001
Sex (n=1,775)						
Male	1			1***		
Female	0.18	(0.04, 0.32)	0.01	0.19	(0.06, 0.32)	0.005
SES (n=1,663)						
Poorest	1			1****		
Middle	0.15	(-0.27, 0.32)		0.11	(-0.06, 0.29)	
Richest	0.30	(0.12, 0.48)	0.005	0.30	(0.10, 0.49)	0.009
Maternal education, years (n=1,775)	0.03	(0.01, 0.05)	0.01	0.02†	(0.004, 0.04)	0.02
Women's decision-making power – major household expenses (n=1,783)						
Husband/other	1			1		
Jointly/self	0.17	(0.01, 0.32)	0.03	0.11‡	(-0.03, 0.26)	0.13
Paternal domestic chores (n=1,684)						
Never	1			1†		
Rarely	0.08	(-0.10, 0.26)		0.08	(-0.09, 0.26)	
Weekly/every day	0.10	(-0.07, 0.27)	0.49	0.09	(-0.07, 0.26)	0.47
Food insecurity score (HFIAS) (n=1,663)	-0.02	(-0.03, -0.002)	0.03	-0.01§	(-0.02, 0.01)	0.38
Minimum dietary diversity-women (MDD-W) (n=1,775)	0.04	(-0.03, 0.11)	0.25	0.01 <sup>l</sup>	(-0.06, 0.08)	0.79
Animal source food types produced (n=1,663)	-0.02	(-0.06, 0.03)	0.50	-0.04 <sup>o</sup>	(-0.09, 0.004)	0.08
Fruit and vegetable types produced (n=1,775)	0.06	(0.01, 0.11)	0.01	0.05 <sup>o</sup>	(0.005, 0.10)	0.03
Land owned, hectares (n=1,664)	0.07	(0.007, 0.14)	0.03	0.08 <sup>†</sup>	(0.02, 0.15)	0.01

Table 3 legend

Variables adjusted for in multivariate linear regression modelling:

\* Age, age squared, sex, SES, maternal education, women's empowerment, paternal childcare support, food security, land ownership, irrigation, fertiliser, fruit and vegetables produced (FV), animal source foods produced (ASF)

\*\* Sex

\*\*\* Age, age squared

\*\*\*\* Age, age squared, maternal education, land ownership, irrigation, fertiliser, FV, ASF

† Age, age squared, sex

‡ Age, age squared, sex, maternal education

§ Age, age squared, sex, SES, maternal education, land ownership, irrigation, fertiliser, FV, ASF (n=1663)

<sup>l</sup> Age, age squared, sex, women's empowerment, food security, FV, ASF

Φ Age, age squared, sex, SES, maternal education, land ownership, women's empowerment, irrigation, fertiliser, FV

□ Age, age squared, sex, SES, maternal education, land ownership, women's empowerment, irrigation, fertiliser, ASF

Accepted Article