

CORRECTION

Open Access



Correction to: Navigating environmental constraints to injection preparation: the use of saliva and other alternatives to sterile water among unstably housed PWID in London

Magdalena Harris^{1*}, Jenny Scott², Vivian Hope³, Talen Wright¹, Catherine McGowan¹ and Daniel Ciccarone⁴

Correction to: Harm Reduction Journal (2020) 17:24
<https://doi.org/10.1186/s12954-020-00369-0>

Following publication of the original article [1], the authors would like to correct two sentences in the article:

'Drug-related deaths in the UK are higher than any European Union (EU) country; in the year before Britians exit they accounted for one third of all reported in the EU country and, in the year prior to Britain's exit, accounted for a third of all reported in the EU'

should read: 'Drug-related deaths in the UK are higher than any European Union (EU) country; in the year prior to Britain's exit, they accounted for a third of all reported in the EU'.

and

'Current harm reduction advice advocates the use of WFI, with the "next best" alternative boiled and cooled potable (drinkable cold tap) water, and after this, cold tap water'.

should read: 'Current harm reduction advice advocates the use of WFI, with the "next best" alternative boiled and cooled potable (drinkable) tap water, and after this, cold tap water'.

Author details

¹Department of Public Health, Environments and Society, London School of Hygiene & Tropical Medicine, 15-17 Tavistock Place, London WC1H 9SH, UK. ²Department of Pharmacy & Pharmacology, University of Bath, Claverton Down, Bath BA2 7AY, UK. ³Public Health Institute, Liverpool John Moores University, Tithebarn Street, Liverpool L2 2QP, UK. ⁴School of Medicine, University of California, San Francisco, 513 Parnassus Ave, San Francisco, CA 94143-0410, USA.

Published online: 09 June 2020

Reference

1. Harris M, et al. Navigating environmental constraints to injection preparation: the use of saliva and other alternatives to sterile water among unstably housed PWID in London. *Harm Reduct J*. 2020;17:24. <https://doi.org/10.1186/s12954-020-00369-0>.

The original article can be found online at <https://doi.org/10.1186/s12954-020-00369-0>.

* Correspondence: magdalena.harris@lshtm.ac.uk

¹Department of Public Health, Environments and Society, London School of Hygiene & Tropical Medicine, 15-17 Tavistock Place, London WC1H 9SH, UK
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.