



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



COVID-19 and maternal and perinatal outcomes

Authors' reply

We thank all correspondents for their interest in our report.¹ A common point is the need for better quality studies in this area, which we wholeheartedly echo. Quality and consistency in reporting, particularly of key outcomes such as maternal death, stillbirth, and preterm birth, are essential if we are to tackle these scourges of maternity care.

As always, systematic reviews are restricted to covering what has been published. Susannah Leisher and the International Stillbirth Alliance call for the inclusion of a relevant preprint paper. We recognised that the COVID-19 literature is a rapidly evolving field with high rates of preprint publication. In consultation with editors and reviewers, however, we agreed to exclude such papers to avoid the pseudo-inflation of the sample size and undermine confidence in the findings of the analysis, but acknowledge that this exclusion might have a greater effect on data from LMICs, which are recognised to be less likely to secure publication.

We agree with Clara Calvert and colleagues, and did highlight in our Article, that the finding of an increase in maternal mortality relied on just two data sources, only one of which presented national-level data. The finding is consistent with another report from India not included in the quantitative analysis,² and with national-level evidence from Kenya reported in a preprint publication.³ But we agree that this early identification of increased maternal mortality cannot necessarily be extrapolated throughout the global population and that further investigation is required.

We share Lizbeth Ochoa and colleagues' confidence in the finding that preterm birth seems to have decreased, in high-income countries at least, during the COVID-19 pandemic. We support their call for

further robust investigations using population-level data of a high quality and considering stillbirth and preterm birth together. They suggest that we should have included another study in the quantitative synthesis.⁴ However, we were unable to include this otherwise excellent study because the report included four different time frames around three different implementation dates without extractable raw data.

We read with interest Zeus Aranda and colleagues' experiences in delivering maternity care at their Partners in Health project sites and congratulate them on their successes. Key lessons include the importance of responses to local need based on acceptable and available models of care and resources, highlighted in their community health worker programmes in Lesotho, Malawi, and Mexico. The response to the COVID-19 pandemic has driven an explosion in remote access to health care, community reporting, and the use of technology in a diverse range of settings. We should use this opportunity to improve the coverage of pregnancy and neonatal surveillance data.

EM reports grants from Chugai Pharma and Gedeon Richter; personal fees from Gedeon Richter and Pfizer; and other from Chugai Pharma and Kebomed, outside the submitted work; and is the president and a trustee of Royal College of Obstetricians & Gynaecologists, is a trustee of the British Menopause Society; and is the chair of trustees for Group B Strep Support. All other authors declare no competing interests.

Copyright © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Barbara Chmielewska, Imogen Barratt, Rosemary Townsend, Erkan Kalafat, Jan van der Meulen, Ipek Gurol-Urganci, Pat O'Brien, Edward Morris, Tim Draycott, Shakila Thangaratnam, Kirsty Le Doare, Shamez Ladhani, Peter von Dadelszen, Laura A Magee, *Asma Khalil akhalil@sgul.ac.uk

Fetal Medicine Unit, Department of Obstetrics and Gynaecology, St George's University Hospitals NHS

Foundation Trust (BC, IB, RT, AK), Paediatric Infectious Diseases Research Group and Vaccine Institute, Institute of Infection and Immunity (KLD, SL), and Vascular Biology Research Centre, Molecular and Clinical Sciences Research Institute (AK), St George's University of London, London SW17 0QT, UK; Middle East Technical University, Faculty of Arts and Sciences, Department of Statistics, Ankara, Turkey (EK); Koc University, School of Medicine, Department of Obstetrics and Gynaecology, Istanbul, Turkey (EK); Department of Health Service Research and Policy, London School of Hygiene and Tropical Medicine (JvdM, IG-U); The Royal College of Obstetricians and Gynaecologists, London, UK (PO'B, TD); University College London Hospitals NHS Foundation Trust, London, UK (PO'B); Norfolk and Norwich University Hospitals NHS Foundation Trust, Norwich, Norfolk, UK (EM); North Bristol NHS Trust Department of Women's Health, Westbury on Trym, UK (TD); Institute of Metabolism and Systems Research, WHO Collaborating Centre for Women's Health, University of Birmingham, Birmingham, UK (ST); Immunisation and Countermeasures Division, Public Health England (SL); British Paediatric Surveillance Unit, Royal College of Paediatrics and Child Health, London, UK (SL, LAM); Department of Women and Children's Health, School of Life Course Sciences, King's College London, London, UK (PvD)

- 1 Chmielewska B, Barratt I, Townsend R, et al. Effects of the COVID-19 pandemic on maternal and perinatal outcomes: a systematic review and meta-analysis. *Lancet Glob Health* 2021; **9**: e759-72.
- 2 Goyal M, Singh P, Singh K, Shekhar S, Agrawal N, Misra S. The effect of the COVID-19 pandemic on maternal health due to delay in seeking health care: experience from a tertiary center. *Int J Gynaecol Obstet* 2021; **152**: 231-35.
- 3 Shikuku D, Nyaoko I, Gichuru S, et al. Early indirect impact of COVID-19 pandemic on utilization and outcomes of reproductive, maternal, newborn, child and adolescent health services in Kenya. *medRxiv* 2020; published online Sept 9. <https://doi.org/10.1101/2020.09.09.20191247> (preprint).
- 4 Been JV, Burgos Ochoa L, Bertens LCM, Schoenmakers S, Steegers EAP, Reiss IKM. Impact of COVID-19 mitigation measures on the incidence of preterm birth: a national quasi-experimental study. *Lancet Public Health* 2020; **5**: e604-11.