



Original Research

Trastuzumab-lapatinib as neoadjuvant therapy for HER2-positive early breast cancer: Survival analyses of the CHER-Lob trial



Valentina Guarneri ^{a,b,*}, Maria V. Dieci ^{a,b}, Gaia Griguolo ^{a,b},
Federica Miglietta ^{a,b}, Fabio Girardi ^b, Giancarlo Bisagni ^c,
Daniele G. Generali ^{d,e}, Katia Cagossi ^f, Samanta Sarti ^g,
Antonio Frassoldati ^h, Lorenzo Gianni ⁱ, Luigi Cavanna ^j,
Graziella Pinotti ^k, Antonino Musolino ^l, Federico Piacentini ^m,
Saverio Cinieri ⁿ, Aleix Prat ^{o,p,q}, PierFranco Conte ^{a,b} on behalf of the
CHER-Lob study team

^a Department of Surgery, Oncology and Gastroenterology (DISCOG), University of Padova, 35128 Padova, Italy

^b Medical Oncology 2, Istituto Oncologico Veneto IOV-IRCCS, 35128 Padova, Italy

^c Pathology Unit, Azienda Unità Sanitaria Locale–IRCCS di Reggio Emilia, Reggio Emilia, Emilia-Romagna, Italy

^d Breast Cancer Unit, Azienda Socio Sanitaria Territoriale di Cremona, Cremona, Italy

^e Department of Medical, Surgery and Health Sciences, University of Trieste, Trieste, Italy

^f Ospedale Bernardino Ramazzini, Carpi, Italy

^g IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) "Dino Amadori", 47014 Meldola, Italy

^h Clinical Oncology, Department of Morphology, Surgery and Experimental Medicine, S Anna University Hospital, Ferrara, Italy

ⁱ Oncology Unit Rimini, Azienda USL Romagna, Rimini, Italy

^j Oncologia Medica, ASL di Piacenza, Piacenza, Italy

^k Department of Oncology, ASST-Settelaghi, Varese, Italy

^l Department of Medicine and Surgery, University of Parma, Medical Oncology and Breast Unit, University Hospital of Parma, Parma, Italy

^m Department of Medical and Surgical Sciences, University of Modena and Reggio Emilia, Modena, Italy

ⁿ Medical Oncology & Breast Unit, Antonio Perrino Hospital, Brindisi, Italy

^o Department of Medical Oncology, Hospital Clinic of Barcelona, Barcelona, Spain

^p Translational Genomics and Targeted Therapeutics in Solid Tumors, August Pi i Sunyer Biomedical Research Institute (IDIBAPS), Barcelona, Spain

^q Department of Medicine, University of Barcelona, Barcelona, Spain

Received 22 March 2021; received in revised form 30 April 2021; accepted 8 May 2021

Available online 19 June 2021

* Corresponding author: Department of Surgery, Oncology and Gastroenterology, University of Padova, Division of Medical Oncology 2, Istituto Oncologico Veneto IRCCS. Via Gattamelata 64, 35128 Padova, Italy. Fax: +39 049 821 5932.

E-mail address: valentina.guarneri@unipd.it (V. Guarneri).

KEYWORDS

HER2-positive breast cancer;
Neoadjuvant treatment;
Lapatinib;
Trastuzumab;
Escalation

Abstract Aim: The Cher-LOB randomised phase II study showed that the combination of lapatinib-trastuzumab plus chemotherapy increases pathologic complete response (pCR) rate compared with chemotherapy plus either trastuzumab or lapatinib. Here, we report the post hoc survival analysis as per treatment arm, pCR and biomarkers.

Methods: The Cher-LOB study randomised 121 patients with human epidermal growth factor receptor 2-positive, stage II–III breast cancer. A specific protocol to collect recurrence-free survival (RFS) and overall survival (OS) data was designed. Tumour-infiltrating lymphocytes (TILs) and PAM50-intrinsic subtyping were evaluated at baseline.

Results: At 9-year median follow-up, a trend towards RFS improvement with lapatinib-trastuzumab over trastuzumab was observed (hazard ratio [HR] 0.44, 95% confidence interval [CI] 0.18–1.05). Combining treatment arms, pCR was significantly associated with both RFS (HR 0.12, 95% CI 0.03–0.49) and OS (HR 0.12, 95% CI 0.03–0.49). TILs were significantly associated with RFS (HR = 0.978 for each 1% increment). Luminal-A subtype was a significant and independent predictor of improved RFS as compared with other PAM50-based intrinsic subtypes at the multivariate analysis including the most relevant clinicopathologic variables (HR 0.29, 95% CI 0.09–0.94, $p = 0.040$).

Conclusions: Cher-LOB trial survival analysis confirmed the prognostic role of pCR and TILs and showed a signal for a better outcome with lapatinib-trastuzumab over trastuzumab.

Trial registration: NCT00429299.

© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Human epidermal growth factor receptor 2 (HER2) is overexpressed/amplified in around 20% of early breast cancers (BCs) and, in the absence of targeted treatment, represents a poor prognostic factor.

Over the past three decades, the implementation of HER2-targeted treatments has progressively improved the prognosis of HER2-positive BC [1–3]. The neoadjuvant model represents an ideal platform to evaluate new therapeutic strategies, and the adoption of pathologic complete response (pCR) as a surrogate end-point in neoadjuvant trials with regulatory intent is currently endorsed by US Food and Drug Administration (FDA) [4]. Indeed, irrespective of tumour biology and disease stage at diagnosis, patients achieving pCR have a better outcome than patients with residual disease [5]. In the neoadjuvant setting, the addition of trastuzumab to chemotherapy has been associated with a significant increase in pCR rates over chemotherapy alone and also with an improved outcome [6–8]. Therefore, several studies testing different strategies have been set up, with the primary aim of increasing pCR rate. In particular, dual HER2-blockade with trastuzumab and lapatinib provided meaningful clinical results. As compared with chemotherapy and trastuzumab, the combination of trastuzumab-lapatinib and chemotherapy was associated with an increased pCR rate in several randomised trials [9–13], thus providing evidence supporting the incorporation of this escalated approach in the neoadjuvant management of HER2-positive disease.

However, the surrogacy of pCR for drug efficacy is still object of extensive debate. Indeed, in the above-mentioned FDA-promoted CTNeoBC meta-analysis [5], authors failed to formally establish a trial-level relationship between pCR differences between treatment arms and survival. Emblematically, although a trend towards increased survival rates has been consistently observed with dual versus single HER2-blockade across escalated neoadjuvant trials, only the CALGB 40601 phase III trial so far reported a significant improvement in survival outcomes [9,10].

We previously reported findings from the phase II non-comparative randomised Cher-LOB trial, showing a relative 80% increase in pCR rates (ypT0is/N0) in patients treated with lapatinib plus trastuzumab plus chemotherapy versus chemotherapy plus either lapatinib or trastuzumab [13].

Here, we report the post hoc analysis of long-term survival outcomes (recurrence-free survival, RFS; overall survival, OS) from the Cher-LOB trial, as per treatment arms, pCR status and biomarkers.

2. Methods

2.1. Cher-LOB study design, patients and study procedures

The Cher-LOB study design, procedures and primary efficacy results have been previously published and are detailed in [Supplementary Methods](#) [13–15]. Briefly, 121 patients with stage II–III HER2-positive BC were

centrally randomised (1:1:1) to receive paclitaxel–anthracycline–based neoadjuvant chemotherapy plus trastuzumab (arm A), lapatinib (arm B) or concomitant trastuzumab plus lapatinib (arm C). Ethical Committees of participating sites approved the study, and informed consent was obtained from patients before study entry.

pCR was defined as the absence of residual invasive tumour in breast and axillary nodes. Methodology used to evaluate tumour-infiltrating lymphocytes (TILs) and PAM50 intrinsic subtyping [14,15] is detailed in [Supplementary Methods](#).

2.2. Outcomes

A specific protocol to continue survival data collection of patients enrolled in the Cher-LOB trial was approved by participating centers' institutional review boards.

RFS was calculated from randomisation to BC recurrence (locoregional or distant; contralateral BC excluded) or death from any cause, whichever first. OS was calculated from randomisation to death from any cause or last follow-up. Patients without an event were censored at time of the last clinical assessment.

2.3. Statistical analysis

The Kaplan-Meier method was adopted to estimate survival curves and 5-year rates, whereas the Cox proportional hazard model was used to compare RFS and OS among subgroups. We examined the relationship between survival end-points and treatment arm, pathologic response at surgery (pCR vs no-pCR), oestrogen receptor (ER) status and biomarkers (TILs, PAM50-based intrinsic subtyping). The association between RFS and biomarkers was evaluated at univariate and multivariate analyses. Hazard ratios (HRs) and 95% confidence intervals (CIs) were also reported. All tests were two-sided and significant for $P \leq 0.05$. Statistical analyses were performed using SPSS v.26.

Because the Cher-LOB trial was not designed or powered to detect treatment differences with respect to survival end-points, RFS and OS analyses are to be considered for descriptive purposes only.

3. Results

In the Cher-LOB trial, 36, 39 and 46 patients were randomly assigned to trastuzumab (A), lapatinib (B) and concomitant trastuzumab plus lapatinib (C) arms, respectively. Patients' clinicopathologic features have been previously reported [13].

Survival data were available for 114 of the 121 patients originally included in the Cher-LOB trial (CONSORT chart in [Supplementary Fig. S1](#)). Patient characteristics were in line with the overall Cher-LOB study cohort, as shown in [Table 1](#). Distribution per arm was as follows: 36 in arm A, 35 in arm B and 43 in

arm C. Patient clinicopathologic features as per treatment arm are summarised in [Table 2](#). pCR was observed in 37 patients (32%).

At a median follow-up of 9 years (95% CI 8.4–9.7), 32 RFS events and 15 deaths were recorded. Patients experiencing an RFS event were 14 (39%), 10 (29%) and 8 (19%) in the trastuzumab, lapatinib or concomitant trastuzumab plus lapatinib arm, respectively. Patterns of first recurrence and causes of death as per treatment arm are reported in [Supplementary Table I](#). The 5-year RFS was 77.8% in the trastuzumab arm, 77.1% in the lapatinib arm and 85.8% in the combination arm ([Fig. 1A](#)). RFS was not statistically different among the three treatment arms (log rank $P = 0.162$); however, a trend towards a benefit for the combination as compared with the trastuzumab arm (HR 0.44, 95% CI 0.18–1.05) was observed. Globally, the 5-year RFS was 77.5% for patients treated with chemotherapy plus single HER2-blockade (A + B) and 85.8% for patients treated with chemotherapy plus dual HER2-blockade (C) (HR 0.52, 95% CI 0.23–1.15; log-rank $P = 0.102$; [Fig. 1B](#)). There were no significant differences in RFS as per ER expression (5-year RFS 79.8% vs 81.1% for ER-negative and ER-positive tumours, respectively; HR 1.032, 95% CI 0.50–2.11; log-rank $P = 0.932$).

Patients experiencing an OS event were 5 (14%), 4 (11%) and 6 (14%) in the trastuzumab, lapatinib or concomitant trastuzumab plus lapatinib arm, respectively. No significant difference was observed in OS between the combination and the trastuzumab arm (HR 1.00, 95% CI 0.31–3.27).

Combining treatment groups, a significant association between pCR and RFS was observed. Patients achieving pCR had a significantly higher 5-year RFS (97.3%) than those without pCR (72.7%) (HR 0.12, 95% CI 0.03–0.49; log-rank $P < 0.001$; [Fig. 2A](#)). Results were consistent in the ER-negative (5-year RFS in pCR vs no-pCR: 94.7% vs 69.2%, HR 0.11, 95% CI 0.01–0.85; log-rank $P = 0.010$) and ER-positive (5-year RFS in pCR vs no-pCR: 100% vs 74.5%, HR 0.12, 95% CI 0.02–0.90; log-rank $P = 0.014$) cohorts separately.

Patients achieving pCR also presented a significantly higher 5-year OS (97.2% vs 89.5%) than those without pCR (HR 0.12, 95% CI 0.03–0.49; log-rank $P = 0.028$; [Fig. 2B](#)).

Finally, the impact of tumour biology and microenvironment on RFS was assessed.

Baseline stromal TIL (strTIL) evaluation was available for 102 of 114 patients assessable for the long-term outcome and significantly correlated with RFS at univariate analysis: an HR of 0.978 for each 1% TIL increment was observed (95% CI 0.957–0.999, $P = 0.041$). At multivariate analysis including strTILs, pCR, clinical stage, treatment arm (single versus dual HER2-blockade), ER status and age, only pCR maintained a significant association with RFS, as shown in [Supplementary Table 2](#).

Table 1
Clinicopathological characteristics of patients included in the survival analysis compared with the overall CherLOB population.

Variable	Overall CherLOB population			Survival analysis population			p
	N (%)	Mean (range)	IQ (Q1; Q3)	N (%)	Mean (range)	IQ (Q1; Q3)	
Age							
Available	121 (100)	49 (26–68)	12 (44–56)	114 (100)	49 (26–68)	13 (44–57)	ns
Missing	0 (0)						
Clinical stage							
IIA	38 (31.4)			37 (32.5)			ns
IIB	61 (50.4)			57 (50.0)			
IIIA	22 (18.2)			20 (17.5)			
Missing	0 (0)			0 (0)			
Histotype							
Ductal	115 (95)			108 (94.7)			ns
Lobular	6 (5)			6 (5.3)			
NA	0 (0)			0 (0)			
Grade							
1–2	23 (19)			19 (16.7)			ns
3	77 (63.9)			74 (64.9)			
Missing	21 (17.4)			21 (18.4)			
ER							
ER-	50 (41.3)			45 (39.5)			ns
ER+	71 (58.7)			69 (60.5)			
Missing	0 (0)			0 (0)			
Ki67							
Available	113 (93.4)	29.6 (4–90)	15 (20–35)	107 (93.8)	29.5 (4–90)	15 (20–35)	ns
Missing	8 (6.6)			7 (6.2)			
Arm							
A (CT + T)	36 (29.8)			36 (31.5)			ns
B (CT + L)	39 (32.2)			35 (30.7)			
C (CT + T + L)	46 (38)			43 (37.8)			
Missing	0 (0)			0 (0)			
strTILs							
Available	105 (86.8)	26.8 (0–100)	31 (9–40)	102 (89.5)	26.8 (0–100)	30.5 (9.5–40)	ns
Missing	16 (13.2)			12 (10.5)			
PAM50							
Luminal A	23 (19.0)			21 (18.5)			ns
Luminal B	14 (11.6)			14 (12.3)			
HER2-enriched	22 (18.2)			22 (19.3)			
Basal-like	12 (9.9)			12 (10.5)			
Normal-like	15 (12.4)			15 (13.1)			
Missing	35 (28.9)			30 (26.3)			

N, number; IQ, interquartile; Q, quartile; P, P-value; ER, oestrogen receptor; CT, chemotherapy; T, trastuzumab; L, lapatinib; strTILs, stromal tumour infiltrating lymphocytes; ns, non-significant.

PAM50 intrinsic subtyping was assessable for 84 of 114 patients included in the long-term outcome analysis. Subtype distribution was as follows: HER2-enriched = 26.2% (N = 22), luminal A = 25.0% (N = 21), luminal B = 16.7% (N = 14), basal-like = 14.3% (N = 12) and normal-like = 17.9% (N = 15). Excluding normal-like cases, no significant difference in RFS as per PAM50 intrinsic subtyping was observed (evaluatable N = 69; [Supplementary Fig. S2](#)); however, at multivariate analysis including PAM50 subtypes (luminal A versus other subtypes), pCR, clinical stage, treatment arm (single versus dual HER2-blockade), ER status and age, both pCR and luminal A subtype (as compared with other PAM50 subtypes) were confirmed to retain an independent prognostic value, as shown in [Table 3](#).

4. Discussion

Findings from the post hoc survival analysis of the phase II randomised Cher-LOB trial regarding the prognostic role of pCR in patients with HER2-positive BC receiving neoadjuvant chemotherapy plus anti-HER2 treatment are consistent with previous reports. By combining treatment arms, we showed that achieving pCR was highly and significantly associated with long-term survival. Patients with pCR experienced a 78% relative decrease in the risk of an RFS or OS event at 5 years as compared with those with residual invasive disease. This supports the strong prognostic role that the achievement of pCR retains in HER2-positive BC at a single-patient level, as already highlighted in the context of the CtNeoBC meta-analysis [5] and in studies testing

Table 2
Clinicopathological characteristics of patients included in the survival analysis as per the treatment arm.

Variable	Arm A (CT + T)			Arm B (CT + L)			Arm C (CT + T + L)		
	N (%)	Mean (range)	IQ (Q1; Q3)	N (%)	Mean (range)	IQ (Q1; Q3)	N (%)	Mean (range)	IQ (Q1; Q3)
Number of patients	36			35			43		
Age									
Available	36 (100)	50 (34–65)	13 (44–57)	35 (100)	49 (34–68)	13 (43–56)	43 (100)	48 (26–65)	13 (43–56)
Missing	0 (0)			0 (0)			0 (0)		
Clinical stage									
IIA	11 (30.6)			13 (37.1)			13 (30.2)		
IIB	19 (52.8)			17 (48.6)			21 (48.8)		
IIIA	6 (16.7)			5 (14.3)			9 (20.9)		
Missing	0 (0)			0 (0)			0 (0)		
Histotype									
Ductal	32 (88.9)			33 (94.3)			43 (100)		
Lobular	4 (11.1)			2 (5.7)			0 (0)		
NA	0 (0)			0 (0)			0 (0)		
Grade									
1–2	8 (22.3)			4 (11.4)			7 (16.3)		
3	24 (66.7)			23 (65.7)			27 (62.8)		
Missing	4 (11.1)			8 (22.9)			9 (20.9)		
ER									
ER-	16 (44.4)			12 (34.3)			17 (39.5)		
ER+	20 (55.6)			23 (65.7)			26 (60.5)		
Missing	0 (0)			0 (0)			0 (0)		
Ki67									
Available	34 (94.4)	26.8 (10–60)	13.5 (17.7–31.2)	33 (94.3)	29.3 (12–65)	16.5 (20–36.5)	40 (93)	32.1 (4–90)	20 (20–40)
Missing	2 (5.6)			2 (5.7)			3 (7)		
pCR									
Yes	9 (25)			9 (25.7)			19 (44.2)		
No	27 (75.029)			26 (74.3)			24 (55.8)		

N, number; IQ, interquartile; Q, quartile; ER, oestrogen receptor; CT, chemotherapy; T, trastuzumab; L, lapatinib; pCR, pathological complete response.

dual HER2-blockade as the escalated neoadjuvant approach [7,9,16–18]. Importantly, the positive prognostic impact of pCR observed in our analysis was consistent in both ER-positive and ER-negative subgroups, in line with previous evidence [5,16–19].

Although the Cher-LOB trial was not powered to formally detect survival differences across treatment arms, we confirmed the trend towards improved survival with the combination of lapatinib plus trastuzumab over single HER2-blockade. In the CALGB 40601 phase III

trial, a notable 68% and 66% relative improvement in the 7-year RFS and OS rates, respectively, was reported with lapatinib plus trastuzumab as compared with the trastuzumab single agent [9]. In addition, consistent with our findings, in both the NSABP B41 and NeoALTTO trials, a numerical increase in long-term survival rates with dual HER2-targeting was observed [16,17].

Correlative studies regarding the association between survival and tumour microenvironment and biology in the context of the Cher-LOB study were also performed.




Fig. 1. Kaplan-Meier RFS curves as per A. treatment arm (chemotherapy + trastuzumab vs chemotherapy + lapatinib vs chemotherapy + trastuzumab + lapatinib) and B. dual HER2 blockade vs single HER2 blockade (CT + trastuzumab and CT + lapatinib arms combined). RFS, recurrence-free survival; CI, confidence interval; HR, hazard ratio; CT, chemotherapy; HER2, human epidermal growth factor receptor 2.




Fig. 2. Kaplan-Meier curves by pCR. A. RFS curve; B. OS curve. RFS, recurrence-free survival; CI, confidence interval; HR, hazard ratio; pCR, pathological complete response; OS, overall survival.

We previously reported a positive association between higher baseline TIL levels and pCR in the Cher-LOB trial. Here, we found baseline TILs to be prognostic in the overall Cher-LOB study population. In particular, similar to the findings of the NeoALTT0 translational analysis [20], each 1% TIL increase corresponded to a 2.2% decrease in the risk of an RFS event across treatment groups. These results are further supported by a pooled analysis assessing the prognostic role of TILs as per BC subtype [21]. Indeed, among 1379 patients with HER2-positive BC undergoing neoadjuvant chemotherapy (the majority also receiving anti-HER2 treatment), a positive association between TILs and disease-free survival was detected. Our findings therefore add further evidence in support to the clinical validity of TILs as a prognostic biomarker in HER2-positive early BC [20–25]. Importantly, TIL evaluation is currently endorsed by both European Society for Medical Oncology guidelines [26] and World Health Organisation Classification of Tumours, 5th edition, (<http://publications.iarc.fr>) in this setting.

Besides TILs, PAM50-based intrinsic subtyping is gaining increasing clinical relevance in HER2-positive disease. We previously reported that HER2-enriched and luminal A subtypes account for most of the Cher-LOB study population (HER2-enriched = 26.2%, luminal A = 25%), with the first being associated with the highest (50%) and the second with the lowest (21%) pCR rate [14]. In the present work, no significant association was identified between RFS and intrinsic subtypes. However, the luminal A phenotype was significantly and independently associated with longer 5-year RFS than other PAM50 subtypes, and this is consistent with the CALGB 40601 translational analysis, where, similarly, patients with luminal A BC showed the lowest pCR rate (14.3%) while experiencing the most favourable long-term outcome as compared with other subtypes (no RFS events at 7 years) [9]. The notion that the luminal A phenotype exhibits a lower degree of sensitivity to chemotherapy plus HER2-blockade while retaining a positive impact on survival apparently challenges the prognostic value of

pCR. However, this is an example of so-called Simpson's paradox [27,28], describing the confounding impact of unmeasured, but causal, variables on associations, which translates into an inverse relationship between chemosensitiveness and prognosis in case of residual disease after neoadjuvant treatment. Consistently, although pCR is prognostic in all traditionally defined BC subtypes (hormone receptor-positive/HER2-negative, HER2-positive and triple-negative), this association is stronger in most chemosensitive subgroups, namely triple-negative and HER2-positive. These observations [7,9,16,29,30] suggest that even among patients with HER2-positive BC, PAM50 analysis may add a further layer of stratification, beyond hormone receptor status [31]. Indeed, PAM50 analysis may allow controlling for those causal variables potentially affecting the strength of the association between pCR and prognosis and which would otherwise dilute the observed prognostic impact of pCR itself. Our findings therefore emphasise that this source of

Table 3

Multivariate analysis for RFS including PAM50 subtypes (luminal-A versus other subtypes), pCR, clinical stage (IIA vs IIB vs IIIA), treatment arm (single versus dual HER2-blockade), ER status (positive versus negative) and age (continuous variable).

Variable	HR (95% CI)	p
PAM50		
Luminal-A vs other	0.29 (0.09–0.94)	0.040
Pathologic response	0.09 (0.01–0.77)	0.027
pCR vs no pCR		
Clinical stage		
Stage IIA	0.65 (0.16–2.70)	0.554
Stage IIB	0.84 (0.22–3.27)	0.806
Stage IIIA	ref	0.810
Treatment arm		
Dual vs single HER2-blockade	0.83 (0.32–2.18)	0.711
ER status		
Positive vs negative	1.69 (0.61–4.73)	0.314
Age (cont.)	0.99 (0.95–1.04)	0.863

p values < 0.05 are highlighted in bold. RFS, recurrence-free survival; CI, confidence interval; HR, hazard ratio; HER2, human epidermal growth factor receptor 2; cont, continuous; vs, versus; pCR, pathologic complete response; ER, oestrogen receptor.

heterogeneity within HER2-positive BC should be taken into consideration when designing future neoadjuvant trials.

The neoadjuvant platform, albeit endorsed by the FDA [32] for the evaluation of investigational interventions given the more favourable cost-effective balance as compared with the adjuvant setting, can be underpowered to demonstrate significant survival differences across treatment arms. Indeed, the sample size required to capture meaningful pCR differences is significantly smaller than that required to detect significant survival differences. Indeed, although nearly all the trials comparing the combination of trastuzumab + lapatinib versus trastuzumab alone (in association with chemotherapy) for the neoadjuvant management of HER2-positive breast cancer reported a substantial improvement in pCR rates with the dual HER2 targeting strategy as compared with single HER2-blockade [9–13], many of them, with the exception of the CALGB 40601 trial [9], failed to demonstrate a statistically significant improvement in survival end-points. However, they were all consistent in confirming the positive prognostic impact of pCR at a single-patient level [5,7,9,16–18]. Overall, these observations suggest that such lack of survival improvement with dual over single HER2-blockade may reflect a statistical limitation, rather than a real absence of long-term advantage.

For this reason, although possibly limited by the relatively small sample size, long-term survival results from the Cher-LOB study are worthy of attention because they add further evidence in the same direction as that provided by other neoadjuvant studies of dual HER2-blockade.

Author contributions

Concept and design: Guarneri, Conte. Acquisition, analysis, or interpretation of data: All authors. Statistical analysis: Dieci. Drafting of the manuscript: Guarneri, Griguolo, Miglietta. Critical revision of the manuscript: All authors.

Funding

The authors acknowledge the following funding: DOR funding from the University of Padova—Department of Surgery, Oncology and Gastroenterology BIRD 2019 (to V.G., M.V.D. and P.F.C.) and BIRD 2020 (to V.G., M.V.D. and G.G.). A.P. would like to acknowledge funds from the Instituto de Salud Carlos III (PI13/01718) and a Career Catalyst Grant from the Susan Komen Foundation and Banco Bilbao Vizcaya Argentaria (BBVA) Foundation.

Role of the funder/sponsor

The funders had no role in the design and conduct of the study; collection, management, analysis and

interpretation of the data; preparation, review or approval of the manuscript and decision to submit the manuscript for publication.

Data sharing statement

The ChER-LOB study data sets are not publicly available to protect patient privacy. Gene expression data from the ChER-LOB study have been previously deposited (GEO database, GSE66399).

Conflict of interest statement

V.G. reports personal fees from Roche, Novartis, Eli Lilly and MSD outside the submitted work. D.G.G. reports personal fees from Roche, Novartis, Eli Lilly and Pierre Fabre, all outside the submitted work. A.F. reports personal fees from Roche, Novartis, Pfizer, Lilly, Daiichi and Seagen, all outside the submitted work. L.G. reports honoraria for lectures and ad boards from Lilly, Pfizer, Novartis and AstraZeneca, all outside the submitted work, reports travel accommodation and expenses from Novartis and Daiichi Sankyo and is an uncompensated member of the Olympia steering committee. A.M. reports grants and personal fees from Roche and Eisai; personal fees from Lilly, MacroGenics and Novartis and grants from Pfizer, all outside the submitted work. S.C. reports personal fees from Lilly Oncology outside the submitted work. A.P. reports grants and personal fees from Roche, AstraZeneca, Daiichi Sankyo, Merck Sharp & Dohme (MSD), PUMA Biotechnology, Novartis and Nanostring Technologies, personal fees from Seattle Genetics, Lilly, Pfizer, Guardant Health, Oncolytics Biotech and Abbvie, all outside the submitted work, and a patent (WO2018/103834A1) licensed to Nanostring Technologies, a patent (WO/2018/096191) issued. M.V.D. reports personal fees from Lilly, Genomic Health, Novartis and Celgene, all outside the submitted work. P.F.C. reports personal fees from Novartis, Eli Lilly, AstraZeneca, Tesaro, BMS and Roche, all outside the submitted work.

All remaining authors have declared no conflicts of interest.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejca.2021.05.018>.

References

- [1] Moja L, Tagliabue L, Balduzzi S, Parmelli E, Pistotti V, Guarneri V, et al. Trastuzumab containing regimens for early breast cancer. *Cochrane Database Syst Rev* 2012 Apr 18;(4): CD006243. doi(4):CD006243.

- [2] Balduzzi S, Mantarro S, Guarneri V, Tagliabue L, Pistotti V, Moja L, et al. Trastuzumab-containing regimens for metastatic breast cancer. *Cochrane Database Syst Rev* 2014 Jun 12;(6): CD006242. doi(6):CD006242.
- [3] Guarneri V, Frassoldati A, Giovannelli S, Borghi F, Conte P. Primary systemic therapy for operable breast cancer: a review of clinical trials and perspectives. *Canc Lett* 2007 Apr 18;248(2):175–85.
- [4] U.S. food and drug administration. guidance for industry. Pathological complete response in neoadjuvant treatment of high-risk early-stage breast cancer: use as an endpoint to support accelerated approval. October 2014. <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM305501.pdf>. accessed 30 october 2014.
- [5] Cortazar P, Zhang L, Untch M, Mehta K, Costantino JP, Wolmark N, et al. Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. *Lancet* 2014 Jul 12;384(9938):164–72.
- [6] Buzdar AU, Ibrahim NK, Francis D, Booser DJ, Thomas ES, Theriault RL, et al. Significantly higher pathologic complete remission rate after neoadjuvant therapy with trastuzumab, paclitaxel, and epirubicin chemotherapy: results of a randomized trial in human epidermal growth factor receptor 2-positive operable breast cancer. *J Clin Oncol* 2005 Jun 1;23(16):3676–85.
- [7] Gianni L, Eiermann W, Semiglazov V, Manikhas A, Lluch A, Tjulandin S, et al. Neoadjuvant chemotherapy with trastuzumab followed by adjuvant trastuzumab versus neoadjuvant chemotherapy alone, in patients with HER2-positive locally advanced breast cancer (the NOAH trial): a randomised controlled superiority trial with a parallel HER2-negative cohort. *Lancet* 2010 Jan 30;375(9712):377–84.
- [8] Guarneri V, Barbieri E, Dieci MV, Piacentini F, Conte P. Anti-HER2 neoadjuvant and adjuvant therapies in HER2 positive breast cancer. *Canc Treat Rev* 2010 Nov;36(Suppl 3):S62–6.
- [9] Fernandez-Martinez A, Krop IE, Hillman DW, Polley MY, Parker JS, Huebner L, et al. Survival, pathologic response, and genomics in CALGB 40601 (alliance), a neoadjuvant phase III trial of paclitaxel-trastuzumab with or without lapatinib in HER2-positive breast cancer. *J Clin Oncol* 2020 Dec 10;38(35):4184–93.
- [10] Carey LA, Berry DA, Cirincione CT, Barry WT, Pitcher BN, Harris LN, et al. Molecular heterogeneity and response to neoadjuvant human epidermal growth factor receptor 2 targeting in CALGB 40601, a randomized phase III trial of paclitaxel plus trastuzumab with or without lapatinib. *J Clin Oncol* 2016 Feb 20;34(6):542–9.
- [11] Robidoux A, Tang G, Rastogi P, Geyer Jr CE, Azar CA, Atkins JN, et al. Lapatinib as a component of neoadjuvant therapy for HER2-positive operable breast cancer (NSABP protocol B-41): an open-label, randomised phase 3 trial. *Lancet Oncol* 2013 Nov;14(12):1183–92.
- [12] Baselga J, Bradbury I, Eidtmann H, Di Cosimo S, de Azambuja E, Aura C, et al. Lapatinib with trastuzumab for HER2-positive early breast cancer (NeoALTT0): a randomised, open-label, multicentre, phase 3 trial. *Lancet* 2012 Feb 18;379(9816):633–40.
- [13] Guarneri V, Frassoldati A, Bottini A, Cagossi K, Bisagni G, Sarti S, et al. Preoperative chemotherapy plus trastuzumab, lapatinib, or both in human epidermal growth factor receptor 2-positive operable breast cancer: results of the randomized phase II CHER-LOB study. *J Clin Oncol* 2012 Jun 1;30(16):1989–95.
- [14] Dieci MV, Prat A, Tagliafico E, Pare L, Ficarra G, Bisagni G, et al. Integrated evaluation of PAM50 subtypes and immune modulation of pCR in HER2-positive breast cancer patients treated with chemotherapy and HER2-targeted agents in the CherLOB trial. *Ann Oncol* 2016 Oct;27(10):1867–73.
- [15] Guarneri V, Dieci MV, Frassoldati A, Maiorana A, Ficarra G, Bettelli S, et al. Prospective biomarker analysis of the randomized CHER-LOB study evaluating the dual anti-HER2 treatment with trastuzumab and lapatinib plus chemotherapy as neoadjuvant therapy for HER2-positive breast cancer. *Oncologist* 2015 Sep;20(9):1001–10.
- [16] Robidoux A, Tang G, Rastogi P, Geyer C, Azar C, Atkins J, et al. Evaluation of lapatinib as a component of neoadjuvant therapy for HER2+ operable breast cancer: 5-year outcomes of NSABP protocol B-41. *J Clin Oncol* 2016;34(15_suppl). https://doi.org/10.1200/JCO.2016.34.15_suppl.501 (May 20, 2016) 501-501.
- [17] de Azambuja E, Holmes AP, Piccart-Gebhart M, Holmes E, Di Cosimo S, Swaby RF, et al. Lapatinib with trastuzumab for HER2-positive early breast cancer (NeoALTT0): survival outcomes of a randomised, open-label, multicentre, phase 3 trial and their association with pathological complete response. *Lancet Oncol* 2014 Sep;15(10):1137–46.
- [18] Gianni L, Pienkowski T, Im YH, Tseng LM, Liu MC, Lluch A, et al. 5-year analysis of neoadjuvant pertuzumab and trastuzumab in patients with locally advanced, inflammatory, or early-stage HER2-positive breast cancer (NeoSphere): a multicentre, open-label, phase 2 randomised trial. *Lancet Oncol* 2016 Jun;17(6):791–800.
- [19] Jackisch C, Hegg R, Stroyakovskiy D, Ahn JS, Melichar B, Chen SC, et al. HannaH phase III randomised study: association of total pathological complete response with event-free survival in HER2-positive early breast cancer treated with neoadjuvant-adjuvant trastuzumab after 2 years of treatment-free follow-up. *Eur J Canc* 2016 Jul;62:62–75.
- [20] Salgado R, Denkert C, Campbell C, Savas P, Nuciforo P, Aura C, et al. Tumor-infiltrating lymphocytes and associations with pathological complete response and event-free survival in HER2-positive early-stage breast cancer treated with lapatinib and trastuzumab: a secondary analysis of the NeoALTT0 trial. *JAMA Oncol* 2015 Jul;1(4):448–54.
- [21] Denkert C, von Minckwitz G, Darb-Esfahani S, Lederer B, Heppner BI, Weber KE, et al. Tumor-infiltrating lymphocytes and prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with neoadjuvant therapy. *Lancet Oncol* 2018 Jan;19(1):40–50.
- [22] Dieci MV, Miglietta F, Guarneri V. Immune infiltrates in breast cancer: recent updates and clinical implications. *Cells* 2021 Jan 23;10(2). <https://doi.org/10.3390/cells10020223>.
- [23] Dieci MV, Conte P, Bisagni G, Brandes AA, Frassoldati A, Cavanna L, et al. Association of tumor-infiltrating lymphocytes with distant disease-free survival in the ShortHER randomized adjuvant trial for patients with early HER2+ breast cancer. *Ann Oncol* 2019 Mar 1;30(3):418–23.
- [24] Kim RS, Song N, Gavin PG, Salgado R, Bandos H, Kos Z, et al. Stromal tumor-infiltrating lymphocytes in NRG Oncology/NSABP B-31 adjuvant trial for early-stage HER2-positive breast cancer. *J Natl Cancer Inst* 2019 Aug 1;111(8):867–71.
- [25] Krop I, Paulson J, Campbell C, Kiermaier A, André F, Fumagalli D, et al. Genomic correlates of response to adjuvant trastuzumab and pertuzumab in HER2+ breast cancer: biomarker analysis of the APHINITY trial. Presented at ASCO 2017, May 31-June 4, Chicago, IL, USA, abstr 1012.
- [26] Cardoso F, Ohno S, Penault-Llorca F, Poortmans P, Rubio IT, Zackrisson S, et al. Early breast cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2019 Aug 1;30(8):1194–220.
- [27] Carey LA, Dees EC, Sawyer L, Gatti L, Moore DT, Collichio F, et al. The triple negative paradox: primary tumor chemosensitivity of breast cancer subtypes. *Clin Canc Res* 2007 Apr 15;13(8):2329–34.

- [28] Carey LA. CCR 20th anniversary commentary: Simpson's paradox and neoadjuvant trials. *Clin Canc Res* 2015 Sep 15; 21(18):4027–9.
- [29] Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, Mejia JA, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol* 2008 Mar 10;26(8):1275–81.
- [30] Rouzier R, Perou CM, Symmans WF, Ibrahim N, Cristofanilli M, Anderson K, et al. Breast cancer molecular subtypes respond differently to preoperative chemotherapy. *Clin Canc Res* 2005 Aug 15;11(16):5678–85.
- [31] Dieci MV, Miglietta F, Griguolo G, Guarneri V. Biomarkers for HER2-positive metastatic breast cancer: beyond hormone receptors. *Canc Treat Rev* 2020 Aug;88:102064.
- [32] Pathological complete response in neoadjuvant treatment of high-risk early-stage breast cancer: use as an endpoint to support accelerated approval guidance for industry. U.S. Department of Health and Human Services Food and Drug Administration Oncology Center of Excellence Center for Drug Evaluation and Research (CDER) Center for Biologics Evaluation and Research (CBER); July 2020.