





























COVID-19 hospitalization	TAF/FTC	156	2.0	(1.6, 2.3)	0 (ref.)	..	1 (ref.)	..
	TDF/FTC	5	0.9	(0.1, 1.7)	-1.1	(-1.9, -0.2)	0.43	(0.07, 0.87)
	ABC/3TC	63	2.2	(1.6, 2.8)	0.2	(-0.4, 0.9)	1.09	(0.79, 1.48)
	Other	60	2.4	(1.8, 3.1)	0.4	(-0.3, 1.1)	1.21	(0.88, 1.62)
COVID-19 ICU admission	TAF/FTC	43	0.5	(0.4, 0.7)	0 (ref.)	..	1 (ref.)	..
	TDF/FTC	2	0.3	(0.0, 1.4)	-0.2	(-0.6, 0.8)	0.57	(0.00, 2.63)
	ABC/3TC	18	0.5	(0.3, 0.8)	0.0	(-0.3, 0.3)	0.96	(0.51, 1.81)
	Other	24	0.7	(0.4, 1.1)	0.2	(-0.1, 0.6)	1.38	(0.75, 2.42)

\* Standardized by age, race/ethnicity, smoking status, CD4 cell count, HIV viral load, maximum HIV viral load over the past 12 months, BMI, systolic blood pressure, diastolic blood pressure, average eGFR over the past 12 months, months since February 2020, hospitalization in previous month, comorbidities including: cardiovascular disease, hypertension, liver disease, chronic pulmonary disease, cancer, chronic kidney disease, alcohol/substance use, diabetes, and dementia.