



Near-Complete Genome Sequences of Measles Virus Strains from 10 Years of Uganda Country-wide Surveillance

Prossy Namuwulya,^a Henry Bukenyua,^a Phionah Tushabe,^a Robert Tweyongyere,^b  Josephine Bwogi,^a  Matthew Cotten,^{c,d}  My V. T. Phan^c

^aUganda Virus Research Institute, Entebbe, Uganda

^bDepartment of Veterinary Pharmacy Clinical and Comparative Medicine, Makerere University, Kampala, Uganda

^cU.K. Medical Research Council-Uganda Virus Research Institute and London School of Hygiene and Tropical Medicine Uganda Research Unit, Entebbe, Uganda

^dU.K. Medical Research Council-University of Glasgow Centre for Virus Research, Glasgow, Scotland, United Kingdom

ABSTRACT Measles remains a global health challenge despite the availability of a safe and effective vaccine. Sporadic outbreaks of measles virus infections continue in Uganda. We report eight near-complete genome sequences of measles virus strains from Uganda cases from 2011 to 2020, providing useful data for assessing vaccine escape and local/regional transmission.

Measles virus (MV) is classified in the *Measles morbillivirus* species (genus *Morbillivirus*, family *Paramyxoviridae*) and is encoded by a negative-sense single-stranded RNA linear genome of about 16,000 nucleotides (nt). Measles is a very transmissible viral infection and, despite an effective vaccine, MV continues to infect and sicken individuals throughout the world. Disruptions of routine vaccination by the coronavirus disease 2019 (COVID-19) epidemic are resulting in further increases in global measles cases (1). The currently used MV vaccines are based on a genotype A MV isolate from 1954 (2). Continued MV evolution may erode the efficacy of the vaccines currently in use, requiring adjustments in vaccines or vaccination schedules for protection against currently circulating MV strains. Accordingly, full genome sequences of MV strains isolated from contemporary measles cases from all parts of the world are essential for keeping ahead of this pathogen. We have developed a simple amplicon-based sequencing method suitable for the MinION sequencing platform to generate genome sequences of MV strains, which is especially functional in resource-poor settings. Here, we report eight near-complete genome sequences from Uganda cases from 2011 to 2020, addressing the lack of MV full genomes from Uganda and from East Africa.

MV IgM enzyme-linked immunosorbent assay (ELISA)-positive cases (3) were identified by Expanded Programme on Immunization (EPI) MV surveillance activity in Uganda (4), and throat swab samples from IgM-positive cases were cultured on Vero/hSLAM cells (5) for 1 or 2 passages of 5 days. The MV-positive cultures were sequenced using MV-specific genotype B3 amplicon primers that amplify the full-length MV genome (primer sequences are available at https://github.com/mlcotten13/Measles_primers). Briefly, total viral RNA was extracted from cell culture material using QIAamp viral RNA minikits following the manufacturer's instructions. MV RNA was reverse transcribed using amplicon-specific primers and SuperScript III reverse transcriptase (Invitrogen), followed by PCR amplification using Phusion high-fidelity DNA polymerase (Thermo Fisher Scientific). PCR amplicon products were subjected to MinION sequencing library preparation using the SQK-LSK109 barcoding kit (Oxford Nanopore Technologies) according to the manufacturer's instructions, followed by sequencing on MinION SpotON R9.4.1 flow cells. The resulting fast5 files were base called and demultiplexed using Oxford Nanopore Technologies Guppy software v.5.0.11+2b6dbff (6). Demultiplexed reads for each sample were used for reference-based assembly using minimap2.1 (7) with MVs/California.USA/34.19/[B3] (GenBank accession

Editor Jelle Matthijnssens, KU Leuven

Copyright © 2022 Namuwulya et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to My V. T. Phan, my.phan@lshtm.ac.uk.

The authors declare no conflict of interest.

Received 14 June 2022

Accepted 8 July 2022

Published 25 July 2022

TABLE 1 Sampling details, assembly statistics, and accession numbers for the strains in this study

Sample identifier	Strain	BioProject accession no.	SRA accession no.	GenBank accession no.	Location	Collection date (yr-mo-day)	No. of raw reads ^a	No. of MV-specific reads	Genome length (nt)	GC content (%)	No. of nucleotide differences from reference ^b	Identity to reference (%) ^c	No. of nucleotide differences from earliest Uganda strain ^d	Identity to earliest Uganda strain (%) ^e
GEMV_11_2	MV/Kyenjojo.UGA/24.13 [B3]	PRJNA843031	SRX15483626	ON642794	Kyenjojo	2013-6-11	183,718	170,266	15,657	47.1	349	97.76	20	99.88
GEMV_11_3	MV/Otuke.UGA/13.19 [B3]	PRJNA843031	SRX15483627	ON642795	Otuke	2019-3-28	283,310	256,334	15,700	47.1	174	98.88	354	97.72
GEMV_11_4	MV/Kole.UGA/05.20 [B3]	PRJNA843031	SRX15483628	ON642796	Kole	2020-1-30	426,125	393,265	15,709	46.5	193	98.76	373	97.60
GEMV_12_9	MV/Jinja.UGA/50.11 [B3]	PRJNA843031	SRX15483629	ON642797	Jinja	2011-12-7	120,874	106,074	15,624	47.6	338	98.82	0	100
GEMV_12_12	MV/Kabarole.UGA/13.13 [B3]	PRJNA843031	SRX15483630	ON642798	Kabarole	2013-3-26	298,611	273,575	15,645	47.6	344	97.79	14	99.92
GEMV_12_18	MV/Kampala.UGA/32.17 [B3]	PRJNA843031	SRX15483631	ON642799	Kampala	2017-8-10	167,430	165,049	15,710	47.4	164	98.95	341	97.81
GEMV_12_20	MV/Bushenyi.UGA/01.12 [B3]	PRJNA843031	SRX15483632	ON642800	Bushenyi	2012-1-2	144,609	129,043	15,543	47.1	348	97.76	30	99.81
GEMV_12_23	MV/Hoima.UGA/01.13 [B3]	PRJNA843031	SRX15483633	ON642801	Hoima	2013-1-4	1,588,359	1,387,323	15,678	47.4	350	97.75	20	99.87

^a Read lengths of 2,000 to 2,500 nt.

^b Number of nucleotide differences from the genome of MVs/California.USA/34.19/[B3] (GenBank accession number [MT789820](#)).

^c Percentage of nucleotide differences from the genome of MVs/California.USA/34.19/[B3] (GenBank accession number [MT789820](#)).

^d Number of nucleotide differences from the earliest Uganda MV strain (sample identifier GEMV_12_9) near-complete genome (GenBank accession number [ON642797](#)).

^e Percentage of nucleotide differences from the earliest Uganda MV strain (sample identifier GEMV_12_9) near-complete genome (GenBank accession number [ON642797](#)).

number [MT789820](#)) as the reference sequence. Consensus genome sequences and final sequence checks were determined using Geneious Prime v.2022.1.1 (8). All tools were run with default parameters unless otherwise specified. Differences or ambiguities in the genome sequences were resolved manually by counting coverage with quality-controlled read data. Details for the samples are found in Table 1, including case dates and locations, sequencing metrics, and GenBank and SRA accession numbers for the sequence data.

The eight near-complete MV genome sequences reported here were obtained from Uganda measles cases identified in 2011 to 2020 from locations across the country. Despite small numbers, these eight sequences help address the lack of MV genome sequences from Uganda and from East Africa, where outbreaks of measles occur regularly. The sequences were all classified as genotype B3, the genotype most frequently reported from Africa (9, 10). The MV sequencing primers and method for the MinION platform reported here provide a useful tool for monitoring MV evolution and transmission.

The study was approved by the Uganda Ministry of Health (reference number 105/197/01), the Uganda Virus Research Institute Research and Ethics Committee (reference number GC/127/19/12/740), and the Uganda National Council for Science and Technology (reference number HS2741).

Data availability. The genome sequences described here have been deposited in GenBank with the accession numbers [ON642794](#) to [ON642801](#). The unassembled read data are available in the SRA with the accession numbers [SRX15483626](#) to [SRX15483633](#) under BioProject accession number [PRJNA843031](#) (Table 1). Primer sequences are available at https://github.com/mlcotten13/Measles_primers.

ACKNOWLEDGMENTS

This work was supported by the National Institute for Health and Care Research (NIHR) (project reference number 17/63/82) using aid from the U.K. Government to support global health research. The study was additionally funded by a Wellcome-Department for International Development-Wellcome Epidemic Preparedness-Coronavirus grant (grant 220977/Z/20/Z) awarded to M.C.

The views expressed in this publication are those of the authors and not necessarily those of the NIHR or the U.K. Department of Health and Social Care.

REFERENCES

1. Our World in Data. 2022. Reported cases of measles, 1980 to 2020. https://ourworldindata.org/grapher/reported-cases-of-measles?country=~OWID_WRL. Accessed 15 May 2022.
2. Bankamp B, Takeda M, Zhang Y, Xu W, Rota PA. 2011. Genetic characterization of measles vaccine strains. *J Infect Dis* 204(Suppl 1):S533–S548. <https://doi.org/10.1093/infdis/jir097>.
3. Hübschen JM, Bork SM, Brown KE, Mankertz A, Santibanez S, Ben Mamou M, Mulders MN, Muller CP. 2017. Challenges of measles and rubella laboratory diagnostic in the era of elimination. *Clin Microbiol Infect* 23: 511–515. <https://doi.org/10.1016/j.cmi.2017.04.009>.
4. Mulders M. 2018. Manual for the laboratory-based surveillance of measles, rubella, and congenital rubella syndrome. <https://www.technet-21.org/en/topics/laboratory-measles-rubella-manual>.
5. Ono N, Tatsuo H, Hidaka Y, Aoki T, Minagawa H, Yanagi Y. 2001. Measles viruses on throat swabs from measles patients use signaling lymphocytic activation molecule (CDw150) but not CD46 as a cellular receptor. *J Virol* 75:4399–4401. <https://doi.org/10.1128/JVI.75.9.4399-4401.2001>.
6. Wick RR, Judd LM, Holt KE. 2019. Performance of neural network basecalling tools for Oxford Nanopore sequencing. *Genome Biol* 20:129. <https://doi.org/10.1186/s13059-019-1727-y>.
7. Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. <https://doi.org/10.1093/bioinformatics/bty191>.
8. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
9. Rota PA, Brown K, Mankertz A, Santibanez S, Shulga S, Muller CP, Hübschen JM, Siqueira M, Beirnes J, Ahmed H, Triki H, Al-Busaidy S, Dosseh A, Byabamazima C, Smit S, Akoua-Koffi C, Bwogi J, Bukenya H, Wairagkar N, Ramamurty N, Incomserb P, Pattamadilok S, Jee Y, Lim W, Xu W, Komase K, Takeda M, Tran T, Castillo-Solorzano C, Chenoweth P, Brown D, Mulders MN, Bellini WJ, Featherstone D. 2011. Global distribution of measles genotypes and measles molecular epidemiology. *J Infect Dis* 204(Suppl 1):S514–S523. <https://doi.org/10.1093/infdis/jir118>.
10. World Health Organization. 2015. Genetic diversity of wild-type measles viruses and the global measles nucleotide surveillance database (MeaNS). *Wkly Epidemiol Rec* 90:373–380.