

Associations Between Late Pregnancy Dietary Inflammatory Index (DII) and Offspring Bone Mass: A Meta-Analysis of the Southampton Women's Survey (SWS) and the Avon Longitudinal Study of Parents and Children (ALSPAC)

Stephen J Woolford,¹ Stefania D'Angelo,¹ Giulia Mancano,² Elizabeth M Curtis,¹ Shanze Ashai,¹ Nitin Shivappa,³ James R Hébert,^{3,4} Sarah R Crozier,^{1,5} Catherine M Phillips,⁶ Matthew Suderman,² Caroline L Relton,² Cyrus Cooper,^{1,7,8} and Nicholas C Harvey,^{1,8}  and the ALPHABET Consortium Investigators

¹MRC Lifecourse Epidemiology Centre, University of Southampton, Southampton, UK

²MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

³Cancer Prevention and Control Program and Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC, USA

⁴Department of Nutrition, Connecting Health Innovations LLC, Columbia, SC, USA

⁵NIHR Applied Research Collaboration Wessex, Southampton Science Park, Southampton, UK

⁶School of Public Health, Physiotherapy, and Sports Science, University College Dublin, Belfield, Ireland

⁷NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK

⁸NIHR Biomedical Research Centre, University of Oxford, Oxford, UK

ABSTRACT

Systemic inflammation is associated with reduced bone mineral density and may be influenced by pro-inflammatory diets. We undertook an observational analysis of associations between late pregnancy energy-adjusted dietary inflammatory index (E-DII) scores and offspring bone outcomes in childhood. E-DII scores (higher scores indicating pro-inflammatory diets) were derived from food frequency questionnaires in late pregnancy in two prospective mother-offspring cohorts: the Southampton Women's Survey (SWS) and the Avon Longitudinal Study of Parents and Children (ALSPAC). The mean (SD) offspring age at dual-energy X-ray absorptiometry (DXA) scanning was 9.2 (0.2) years. Linear regression was used to assess associations between E-DII and bone outcomes, adjusting for offspring sex and age at DXA and maternal age at childbirth, educational level, pre-pregnancy body mass index (BMI), parity, physical activity level, and smoking in pregnancy. Associations were synthesized using fixed-effect meta-analysis. Beta coefficients represent the association per unit E-DII increment. In fully adjusted models (total $n = 5910$) late pregnancy E-DII was negatively associated with offspring whole body minus head bone area (BA: $\beta = -3.68$ [95% confidence interval $-6.09, -1.27$] cm^2/unit), bone mineral content (BMC: $\beta = -4.16$ [95% CI $-6.70, -1.62$] g/unit), and areal bone mineral density (aBMD: $\beta = -0.0012$ [95% CI $-0.0020, -0.0004$] $\text{g}\cdot\text{cm}^{-2}/\text{unit}$), but there was only a weak association with BMC adjusted for BA ($\beta = -0.48$ [95% CI $-1.11, 0.15$] g/unit) at 9 years. Adjustment for child height partly or, for weight, fully attenuated the associations. Higher late pregnancy E-DII scores (representing a more pro-inflammatory diet) are negatively associated with offspring bone measures, supporting the importance of maternal and childhood diet on longitudinal offspring bone health. © 2022 The Authors. *Journal of Bone and Mineral Research* published by Wiley Periodicals LLC on behalf of American Society for Bone and Mineral Research (ASBMR).

KEY WORDS: OSTEOPOROSIS; EPIDEMIOLOGY; CHILDHOOD; BONE; DXA; E-DII; DIET; INFLAMMATION; SWS; ALSPAC

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Received in original form August 16, 2021; revised form April 20, 2022; accepted June 4, 2022.

Address correspondence to: Nicholas Harvey, PhD, MRC Lifecourse Epidemiology Centre (University of Southampton), Southampton General Hospital, Southampton, SO16 6YD, UK. E-mail: nch@mrc.soton.ac.uk

Additional Supporting Information may be found in the online version of this article.

*The ALPHABET Consortium Investigators are Adrien M Aubert, Jonathan Y Bernard, Ling-Wei Chen, Liesbeth Duijts, Aisling A Geraghty, Wojciech Hanke, Barbara Heude, Cecily C Kelleher, Fionnuala M McAuliffe, John Mehegan, Sara M Mensink-Bout, Kinga Polanska, and Elzbieta Trafalska.

Journal of Bone and Mineral Research, Vol. 37, No. 8, August 2022, pp 1511–1519.

DOI: 10.1002/jbmr.4623

© 2022 The Authors. *Journal of Bone and Mineral Research* published by Wiley Periodicals LLC on behalf of American Society for Bone and Mineral Research (ASBMR).

Introduction

Chronic systemic inflammation has been associated with several musculoskeletal outcomes including osteoarthritis, osteoporosis, and fragility fractures.^(1–4) Dietary components, such as certain carbohydrates and fatty acids, have been found to contribute to a chronic inflammatory state,⁽⁵⁾ which may detrimentally affect long-term health. For example, Western diets rich in red meat, high fat, sugar, dairy products, and refined grains have been associated with greater inflammatory load.^(6–8)

The dietary inflammatory index (DII) was created to quantify the inflammatory potential of diet⁽⁹⁾ and, after the first validation,⁽¹⁰⁾ has now been validated with pro-inflammatory cytokines and other biomarkers in more than 40 studies from around the world.⁽¹¹⁾ Both higher DII⁽¹²⁾ and higher energy-adjusted DII (E-DII) scores (both indicating a more pro-inflammatory diet) have been associated with lower bone mineral density and higher fracture risk in adults.^(13,14) Consistent with these musculoskeletal outcomes, higher DII/E-DII score has been associated with a wide range of noncommunicable diseases, including cardiovascular diseases,^(15,16) colorectal cancer,⁽¹⁷⁾ and depression.⁽¹⁸⁾

Additionally, higher maternal E-DII during pregnancy has been associated with lower birth weight and shorter birth length,⁽¹⁹⁾ which may have detrimental implications for early-life bone development. These observations, together with well-established links between the early environment and later musculoskeletal development, give rise to the hypothesis that greater dietary inflammatory load in early life, for example, in utero, may impair bone accrual,⁽²⁰⁾ which might lead to a reduction in peak bone mass (PBM) achieved in early adulthood,^(21,22) with implications for osteoporosis and fracture risk in older age.^(23–25)

We therefore aimed to investigate the associations between late pregnancy and early childhood E-DII and offspring bone outcomes in later childhood in two longitudinal mother–child cohorts within the EU ALPHABET consortium: the Southampton Women’s Survey (SWS) and the Avon Longitudinal Study of Parents and Children (ALSPAC).

Materials and Methods

The ALPHABET consortium is a European Union and national partner organization–funded collaboration bringing together seven mother–child cohorts across five European countries. Its aim is to investigate the associations between maternal diet quality, dietary inflammatory potential, epigenetic markers, and offspring health, ultimately informing future public health policy. The Southampton Women’s Survey (SWS) and the Avon Longitudinal Study of Parents and Children (ALSPAC) are two UK-based prospective mother–offspring cohort studies contributing data to the ALPHABET consortium, and both investigate the effects of anthropometric, environmental, and genetic characteristics on maternal and offspring health. These two cohorts form the basis of the current analysis because of the availability of childhood dual-energy X-ray absorptiometry (DXA) data in addition to the E-DII data. The structure and methods of the SWS and ALSPAC cohorts have been described in detail elsewhere.^(26–28)

In brief, 12,583 non-pregnant women aged 20 to 34 years were recruited to the SWS between 1998 and 2002 from the general population of Southampton, UK, with no other exclusion criteria. For ALSPAC, pregnant women with an expected delivery date between April 1, 1991, and December 31, 1992, and living

in the area of Avon, England, were recruited, with 14,541 women entering the study. For both cohorts, only the first child born during the study time frame was included in further analyses. There was a total of 3158 and 14,062 singleton live births being followed for the SWS and ALSPAC, respectively.

In both cohorts, anthropometry and lifestyle assessments were undertaken at study recruitment, with comparable methodology. Maternal height was measured using a stadiometer and weight with a calibrated digital scale, with body mass index (BMI) being calculated. Research nurses underwent training and regular reassessment to ensure measurement consistency. Information on parity, educational level, smoking habits, and regular physical activity was also gathered via questionnaire at study recruitment. Within the SWS, subsets of mothers from specific general practitioner (GP) practices were approached to participate in a bone assessment substudy and for their child to undergo DXA assessment at several time points, including 9 years of age. Specific GP practices were chosen to avoid participants being recruited into multiple substudies. In ALSPAC, at 9 years all children with known addresses who were still participating in the study were invited to a “Focus @ 9” clinic, with DXA scanning at clinic attendance being offered. In neither study was there a specific inclusion criteria other than the ability to undergo DXA scanning.

The SWS was approved by the Southampton and South West Hampshire Local Research Ethics Committee. Ethics approval for ALSPAC was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. Informed consent for the use of data collected via questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time. Written consent was obtained from the parent or caregiver of all participants at each study stage.

Dietary inflammatory index measurement

The design and development of the original DII was published in 2009.⁽²⁹⁾ Methods were updated about 4 years later, which produced a revised tool addressing shortfalls in the original model, for example, standardizing units of measurement to global referent values and using a markedly expanded literature.⁽⁹⁾ The E-DII was later created to account for differences in energy intake that could influence inflammatory capacity of the diet and could be related to sex, physical activity, and body size.⁽¹¹⁾ At this juncture, the E-DII fits the data better (produces greater model explanatory ability and better goodness of fit) than the DII in about two-thirds of instances. The current version of both the DII and E-DII reflect evidence-based scoring systems derived from a comprehensive review yielding around 6500 articles relating dietary parameters to six inflammatory markers (IL-1b, IL-4, IL-6, IL-10, TNF-a, and C-reactive protein). A total of 1943 studies were found to report on primary associations between these six inflammatory markers and 45 food parameters, including 10 whole-food items and 35 nutrient measures.⁽⁹⁾ Individual food parameters were assigned a positive score (+1) if associated with a pro-inflammatory response, a negative score (–1) if associated with an anti-inflammatory response, or a score of zero if not associated with inflammatory response.⁽⁹⁾ In the present study, dietary information from participants was converted to an amount consumed per 1000 kcal and then linked to a regionally representative database with similarly energy-adjusted values. This database provided an overall estimate of mean and standard deviation of energy-standardized intakes for each of the dietary parameters collected

for the participant (ie, nutrients, foods, and other food components), and these data also were adjusted for energy using the density method. By subtracting the mean of the energy-adjusted regionally representative database from the individual's reported amount and dividing this value by the parameter's representative standard deviation, Z-scores for each dietary parameter were derived. These Z-scores were converted to cumulative proportions (ie, with values ranging from 0 to 1) and then centered by doubling each and subtracting 1. The resulting value was then multiplied by the corresponding nutrient parameter effect score. These food parameter-specific E-DII scores were then summed to yield an overall E-DII score. In this way, an overall E-DII score can be calculated for an individual, with more positive E-DII scores indicating a more pro-inflammatory dietary pattern and lower E-DII scores indicating a more anti-inflammatory diet. Construct validity has since been assessed in relation to inflammatory biomarkers in more than 39 studies.^(10,29,30)

Full methodology regarding how E-DII scores were calculated for the SWS and ALSPAC has been published previously.⁽³¹⁾ Briefly, in both the SWS and ALSPAC, maternal E-DII scores were calculated from food-frequency questionnaires (FFQ) at late pregnancy follow-up clinics (34 weeks' gestation in the SWS and 32 weeks' gestation in ALSPAC). Offspring diet was assessed at 3 years of age, via postal FFQs⁽⁶⁾ from which children's-DII (C-DII) scores were generated.⁽⁶⁾ A total of 100 food items were measured by FFQ in the SWS (previously validated against 4-day food diary⁽³²⁾), and a total of 43 food items were measured by FFQ in ALSPAC,⁽³³⁾ with additional questions about bread, milk, fats, and drinks. Of these food items, 24 and 28 dietary parameters were derived to calculate DII from the SWS and ALSPAC, respectively.⁽³¹⁾ The specific food/nutrient items considered in each cohort are documented in Supplemental Table S2.⁽³¹⁾

Childhood DXA assessment

In the SWS, the child's height (without shoes), using a stadiometer (Seca, Birmingham, UK) and weight (without shoes and wearing light clothing), using calibrated digital scales (Seca), were measured. Similar measurements were taken in ALSPAC, using a stadiometer and a Tanita body fat analyzer (Tanita Europe BV, Amsterdam, Netherlands). The child's age at the time of DXA assessment was also recorded. A Hologic Discovery DXA scanner was used in the SWS children (Hologic Inc., Bedford, MA, USA), whereas a Lunar Prodigy DXA scanner was used in the ALSPAC (GE Healthcare, Chalfont St Giles, UK). Whole-body scans were obtained, generating data on bone indices. Coefficients of variation for whole-body bone mineral density (BMD) were 0.75% and 0.84% in the SWS and ALSPAC, respectively. DXA scans were reviewed, and those with excessive movement or clothing artifacts were omitted from the analysis. In the SWS, 1024 children underwent DXA assessment at 9 years, with 990 having useable images for analysis. In ALSPAC, 7722 children underwent 9-year DXA assessment, with 7333 having useable images for analysis. Bone outcomes of interest obtained directly from the DXA assessment were whole-body bone area (BA), whole-body bone mineral content (BMC), and whole-body areal bone mineral density (aBMD).

Statistical analysis

All baseline characteristics were checked for normality of distribution and described using mean (SD), median (IQR), or number (%), as appropriate. Whole-body minus head DXA data were

analyzed.⁽³⁴⁾ As a further measure to correct for body size, BMC adjusted for BA was used. Outcomes of interest were initially related to late pregnancy E-DII using univariable linear regression analyses. Based on prior literature, biological plausibility, and availability across both cohorts, the following covariates were considered in fully adjusted models: offspring sex and age at DXA scanning and maternal age at child's birth, educational level (A level-equivalent qualification or above in both cohorts), pre-pregnancy BMI, parity, physical activity level (hours of strenuous activity per week in the SWS and regular physical activity at least once a week in ALSPAC), and smoking status in pregnancy. In further exploratory analyses, the potential mediating effect of offspring height or weight was investigated. Additional exploratory analyses used the childhood 3-year C-DII measure as the exposure. Finally, because obesity has relevance for both systemic inflammation and bone health and is clearly linked to diet, we examined whether there were any interactions between maternal E-DII and BMI on offspring bone outcomes. Beta coefficients generated represent increase in the bone outcome of interest per unit greater E-DII score.

Analyses were carried out in each study separately and a meta-analysis of 8- to 9-year SWS and ALSPAC results was then performed. Heterogeneity of effect estimates from the two studies was assessed by the Cochran's Q-statistic and quantified by the I^2 statistic.⁽³⁵⁾ Because there was no statistical evidence of heterogeneity ($p > 0.1$), the effects were combined in a fixed-effects meta-analysis model⁽³⁵⁾ to estimate the pooled effect of late pregnancy and 3-year offspring DII on bone outcomes at 9 years. Forest plots were used as a graphical display of the results of the meta-analysis. Stata V15.1 (StataCorp LP, College Station, TX, USA) was used for all analyses.

Results

Characteristics of mothers and offspring

In the SWS, 931 mothers had E-DII calculated at 34 weeks' gestation and offspring DXA data at 9 years. Their baseline characteristics are summarized in Table 1. The mean (SD) age at delivery was 30.7 (3.8) years, with 48.6% of mothers being multiparous; 61.5% had a senior school high school (A level-equivalent) qualification or above, and 15.1% smoked during pregnancy; 13.7% were obese pre-pregnancy; and 65.4% reported any strenuous physical activity each week. The mean (SD) age of offspring at DXA assessment was 9.2 (0.2) years. Approximately half of all offspring were male. Mean (SD) E-DII score was 0.60 (1.46) for mothers and C-DII -0.04 (1.06) for offspring at age 3 years. Compared with individuals who did not take part in the DXA follow-up, those who did were more likely to have a higher level of education and less likely to smoke during pregnancy. All other characteristics were similar across the groups (Supplemental Table S3a).

In ALSPAC, 6334 mothers had E-DII calculated at 32 weeks' gestation and offspring who underwent DXA scanning at 9 years. Their baseline characteristics are summarized in Table 2. The mean (SD) age at delivery was 29.2 (4.5) years. Most (53.8%) mothers were multiparous, 33.1% had at least an A level-equivalent qualification, and 14.4% smoked during pregnancy. Mean (SD) pre-pregnancy BMI was 22.9 (3.7) kg/m², and 68.0% of participants reported regular physical activity at least once a week. Of the children, 48.9% were male and the mean (SD) age at DXA scanning was 9.9 years (0.3). Mean (SD) E-DII was 0.37 (1.80) for mothers and C-DII 0.53 (1.29) for offspring. Compared with mothers whose child did not take part in the DXA follow-

Table 1. SWS Maternal and Offspring Characteristics

Dietary inflammatory index	<i>n</i>	
Maternal, 34 weeks' gestation (E-DII)	931	Mean 0.48 (1.47); median 0.54 (−0.52, 1.53)
Offspring, 3 years (C-DII)	800	Mean 0.23 (1.47); median −0.03 (−0.76, 0.67)
Maternal characteristics	<i>n^a</i>	Mean (SD) or <i>n</i> (%)
Age at delivery (years)	931	30.8 (3.7)
Parity (≥primiparous)	931	442 (47.5)
Educational level (≥A level)	929	587 (63.2)
Smoked during pregnancy	930	115 (12.4)
Height (cm)	927	163.6 (6.4)
Pre-pregnancy weight (kg)	924	67.3 (13.2)
Pre-pregnancy BMI (kg/m ²)	923	
<18.5 (underweight)		12 (1.3)
18.5–25 (normal)		525 (56.9)
25–30 (overweight)		264 (28.6)
>30 (obese)		122 (13.2)
Hours/week of strenuous physical activity (>0)	928	608 (65.5)
9-year offspring characteristics	<i>n^b</i>	Mean(SD) or <i>n</i> (%)
Sex (male)	969	487 (50.3)
Age at scan (years)	969	9.2 (0.2)
Height (cm)	969	135.6 (6.0)
Weight (kg)	966	31.2 (6.4)
BA (cm ²)	969	1124.3 (157.8)
BMC (g)	969	724.8 (120.4)
aBMD (g/cm ²)	969	0.64 (0.06)
BMC adjusted for BA (g)	969	725.2 (65.3)

SWS = Southampton Women's Study; E-DII = energy-adjusted dietary inflammatory index; C-DII = children's dietary inflammatory index; BMI = body mass index; BA = bone area; BMC = bone mineral content; aBMD = areal bone mineral density.

Data are mean (SD), median (IQR), or number (%).

^aNumbers are mothers with 34 weeks' gestation DII calculated and offspring who underwent DXA assessment at 9 years.

^bNumbers are offspring with 3-year DII calculated and BMC available at 9 years.

Note that numbers of individuals missing values for descriptive variables are summarized in Supplemental Table S1.

up, the mothers included in this study had lower E-DII score, were more likely to have at least an A level, and to be slightly taller (Supplemental Table S3b).

The participant flow through the two cohorts is summarized in Supplemental Fig. S1.

Maternal late pregnancy E-DII, offspring 3-year C-DII, and offspring bone outcomes in SWS and ALSPAC

Associations between either maternal late pregnancy E-DII and offspring bone outcomes are presented for both the SWS and ALSPAC in Table 3, in unadjusted and fully adjusted models. The general pattern is one of greater E-DII (indicating a more pro-inflammatory dietary pattern) associated with poorer bone outcomes (BA, BMC, and aBMD), supported by greater statistical

Table 2. ALSPAC Maternal and Offspring Characteristics

Dietary inflammatory index	<i>n</i>	
Maternal, 32 weeks' gestation (E-DII)	6334	0.37 (1.80)
Offspring, 3 years (C-DII)	5710	0.53 (1.29)
Maternal characteristics	<i>n^a</i>	Mean(SD) or <i>n</i> (%)
Age at delivery (years)	6103	29.2 (4.5)
Parity (≥primiparous)	6194	3331 (53.8)
Educational level (≥A level)	6331	2096 (33.1)
Smoked during pregnancy	5924	852 (14.4)
Height (cm)	5768	164.2 (6.6)
Pre-pregnancy weight (kg)	5768	61.8 (10.7)
Pre-pregnancy BMI (kg/m ²)	5826	22.9 (3.7)
Regular physical activity at least once a week	6010	4090 (68.0)
9-year offspring characteristics	<i>n^b</i>	Mean(SD) or <i>n</i> (%)
Sex (male)	5710	2808 (49.2)
Age at scan (years)	5710	9.9 (0.3)
Height (cm)	5311	139.5 (6.2)
Weight (kg)	5311	34.6 (7.2)
BA (cm ²)	5710	1136.5 (162.7)
BMC (g)	5710	890.6 (182.5)
aBMD (g/cm ²)	5710	0.78 (0.05)
BMC adjusted for BA (g)	5710	894.4 (39.7)

^aNumbers are mothers with 34 weeks' gestation E-DII calculated and offspring who underwent DXA assessment at 9 years of age.

^bNumbers are offspring with 3-year C-DII calculated and BMC available at that time point.

ALSPAC = Avon Longitudinal Study of Parents and Children; E-DII = energy-adjusted dietary inflammatory index; C-DII = children's dietary inflammatory index; BMI = body mass index; BA = bone area; BMC = bone mineral content; aBMD = areal bone mineral density.

Data are mean (SD), median (IQR), or number (%).

evidence in the ALSPAC cohort, consistent with the much larger sample size. In general, associations were robust to adjustment for confounding factors and in exploratory analyses associations with bone outcomes appeared similar for offspring 3-year C-DII (Supplemental Table S3). There was no evidence of an interaction between maternal E-DII and BMI on offspring bone outcomes. Further adjustment for either childhood height (Supplemental Table S4) or weight (Supplemental Table S5) led to partial or full attenuation of the associations, respectively. There was no evidence of associations between either maternal pregnancy E-DII or childhood C-DII and childhood BMC adjusted for BA. As an illustration of the effect size, comparing offspring of mothers in the top compared with the bottom quarter of E-DII distribution in ALSPAC, BMC was on average 15 g greater, representing a mean difference of 0.08 SD.

Meta-analysis of 9-year bone outcomes

Associations between maternal late pregnancy E-DII and 9-year offspring bone outcomes in the SWS and ALSPAC cohorts are presented using forest plots in Fig. 1. In fully adjusted models, late pregnancy E-DII was negatively associated with offspring BA, BMC, and aBMD at 9 years, with weaker negative associations apparent for BMC adjusted for BA. In further exploratory analyses, associations between childhood 3-year C-DII score and

Table 3. Associations Between Maternal Late Pregnancy (34 Weeks) E-DII and Offspring Bone Outcomes at 8 to 9 Years in the SWS or ALSPAC

	Late pregnancy E-DII (units)							
	Unadjusted				Adjusted ^a			
	<i>n</i>	β	SE	<i>p</i>	<i>n</i>	β	SE	<i>p</i>
SWS								
BA (cm ²)	931	-2.49	3.53	0.48	917	-3.25	3.80	0.39
BMC (g)	931	-1.45	2.69	0.59	917	-2.58	2.86	0.37
aBMD (g/cm ²)	931	0.0003	0.0012	0.80	917	-0.0003	0.0013	0.84
BMC for BA (g)	931	0.14	1.44	0.92	917	-0.49	1.49	0.74
ALSPAC								
BA (cm ²)	6334	-2.22	1.14	0.05	4993	-3.73	1.29	<0.01
BMC (g)	6334	-2.95	1.28	0.02	4993	-4.57	1.45	<0.01
aBMD (g/cm ²)	6334	-0.0011	0.0004	<0.01	4993	-0.0013	0.0004	<0.01
BMC for BA (g)	6334	-0.52	0.28	0.06	4993	-0.478	0.33	0.15

SWS = Southampton Women's Study; ALSPAC = Avon Longitudinal Study of Parents and Children; E-DII = energy-adjusted dietary inflammatory index; BA = bone area; BMC = bone mineral content; aBMD = areal bone mineral density.

Table shows regression coefficient and standard error from univariable and multivariable linear regression analyses. Outcomes are whole-body measurements, without heads. Results with $p \leq 0.05$ shown in bold.

^aAdjusted for offspring sex and age at DXA and maternal age at childbirth, educational level, pre-pregnancy body mass index, parity, physical activity level, and smoking in pregnancy status.

9-year bone outcomes in the SWS and ALSPAC cohorts are presented using forest plots in Supplemental Fig. S2. In fully adjusted models, there was evidence of marginally greater associations with bone outcomes compared with maternal late pregnancy E-DII score as the exposure for BA, BMC, and aBMD. Again, associations with BMC adjusted for BA were less robust than with the other three bone measures.

Discussion

To our knowledge, this is the first study to demonstrate associations between consumption of a pro-inflammatory diet during pregnancy or early childhood and offspring bone measures. Higher late pregnancy levels of pro-inflammatory dietary components were associated with lower offspring bone measures




Fig. 1. Meta-analysis of associations between late pregnancy energy-adjusted dietary inflammatory index and 9-year offspring bone outcomes in the SWS and ALSPAC cohorts. SWS = Southampton Women's Study; ALSPAC = Avon Longitudinal Study of Parents and Children; BA = bone area; BMC = bone mineral content; aBMD = areal bone mineral density.

at 9 years. The associations were much weaker for BMC adjusted for BA than for BA, BMC, or aBMD, with associations at least partly explained by child height and weight. Additionally, a more pro-inflammatory diet at 3 years of age was negatively associated with offspring BA and BMC at 9 years. Thus, overall, our findings support the notion that greater exposure to a more pro-inflammatory diet in early life is associated with poorer bone health in childhood.

We have not been able to identify any prior evidence linking childhood bone accrual with either maternal diet-associated inflammation during pregnancy or via the child's own diet earlier in postnatal life. However, other work has previously associated DII score in pregnancy and early childhood with a range of other detrimental health outcomes, including reduced offspring birthweight and greater childhood adiposity,⁽³⁶⁻³⁹⁾ including within the larger ALPHABET consortium.^(19,31) Our findings augment the existing evidence base by demonstrating persistent negative associations between pro-inflammatory diet in pregnancy and childhood with offspring bone mass. Furthermore, although data regarding the musculoskeletal sequelae of elevated DII in pregnancy and early life are somewhat limited, our findings are consistent with those observed at older ages, when DII or other markers of systemic inflammation are associated negatively with measures of bone health.^(1-4,12-14) However, several of the studies mentioned above have examined DII, as opposed to E-DII, during pregnancy and early childhood; although DII and E-DII are scored similarly and scaled identically,⁽¹¹⁾ direct comparison to our results should last be undertaken with this caveat in mind.

Although causality cannot be inferred from our results because of the observational nature of this study, several mechanisms have been proposed by which systemic inflammation during pregnancy may detrimentally affect fetal growth and, by extension, offspring bone health. Raised levels of inflammatory cytokines⁽⁴⁰⁾ may directly transfer across the placenta to the fetus and subsequently impair fetal development.⁽⁴¹⁾ Maternal inflammation may also lead to local inflammation of the placenta, impairing its ability to facilitate maternal-fetal nutrient transfer.⁽⁴²⁻⁴⁴⁾ Although the mother-offspring associations were partly attenuated through adjustment for offspring height, consistent with the much-attenuated associations with BMC for BA, it was apparent that the relationship was not entirely explained through linear growth. Adjustment for child's body weight largely removed the associations, and given that both lean and fat mass tend to be positively related to bone measures, this raises the possibility that the bone relationships might be secondary to differences in lean or fat mass. A further possibility that arises from the inverse association between pro-inflammatory diets and other measures of a healthy diet (discussed in more detail below) relates to the greater calcium intake associated with a healthy diet.⁽⁴⁵⁾ Greater calcium intake did not appear to be pro-inflammatory in the studies underpinning the genesis of the E-DII,⁽⁹⁾ but a less pro-inflammatory diet might be associated with better bone health via greater calcium intake through the inverse association between E-DII and healthy dietary patterns.⁽⁴⁵⁾ However, the evidence linking maternal calcium intake during pregnancy and offspring bone development is limited, and indeed even calcium intake during childhood is inconsistently associated with bone mineral density.^(46,47) Given that vitamin D intake is part of the E-DII, it is difficult to disaggregate its effect as part of the score from any other potential contribution to the associations.

In this collaborative work, cognizant of the complexity and limitations of nutrition research, as documented in a recent position paper,⁽⁴⁸⁾ we studied two large and well-characterized

longitudinal cohorts, with "gold standard" measurements of offspring bone indices. However, despite these strengths, there are several limitations that should be considered in the interpretation of results from this study. First, the data produced by the DXA assessment of children can be affected by their lower BMD when compared with adults and the greater likelihood of children moving during scanning. To allow for this, specific pediatric software was used in both cohorts at all time points, which minimizes the loss of edge detection, and images with excessive movement artifacts were excluded from the analysis.⁽⁴⁹⁾ Second, although the SWS and ALSPAC are largely comparable cohorts in terms of methodology and setting, the characteristics of their participants differed in several ways. Specifically, mothers in ALSPAC were less likely to have an A-level equivalent qualification and had a lower mean DII; offspring in ALSPAC had greater mean E-DII, when compared with the SWS. Undertaking analysis within each cohort and synthesizing by meta-analysis should mitigate any potential impact of these differences in the associations observed. Third, although E-DII has been extensively validated against a wide range of inflammatory biomarkers in a variety of settings,^(10,29,30,50) no internal validation against biomarkers taken from participants within the SWS or ALSPAC was able to be performed. Fourth, as in virtually all observational studies, diet was self-assessed, which introduces the potential for information bias. Fifth, there is limited validation of the C-DII measure in younger children, with the previous validation study using C-DII in rather older children aged 6 to 14 years. The tracking of DII over age is not well characterized and so the relevance of the 3-year measure for bone health at 9 years is uncertain. Hence, we view the associations between the childhood measures as intriguing but clearly requiring further replication. Sixth, we were not able to broaden our study to the remaining ALPHABET cohorts because DXA measures at a comparable age were not available for those children. Seventh, it was not appropriate in this setting to discern an E-DII threshold above which a diet might be regarded as "unhealthy" for bone outcomes. Although defining a threshold might be a helpful concept for clinical and public health practice, it would clearly require replication in other cohorts across diverse populations. Finally, we did not have longitudinal repeated measures of diet in mothers or children, and common to observational designs, the potential for residual confounding will always remain. Thus, future studies in independent cohorts including consistent longitudinal assessments of diet are warranted to confirm these findings.

If our observed associations persist into the later lifecourse, there are implications for musculoskeletal health in adulthood. Early bone accrual during childhood is an important contributor to peak bone mass, achieved by approximately the third decade of age.^(21,22) Therefore, it is conceivable that E-DII in early life might impact the magnitude of an individual's peak bone mass with implications for the onset of osteoporosis⁽²³⁾ and increased hip fracture risk⁽²⁴⁾ in later life. The effect size observed in our analysis is relatively modest, with a mean difference of 0.08 SD in childhood BMC between the bottom and top quarters of the distribution of maternal E-DII. However, the very strong relationship between bone mass and fracture risk suggests that, if maintained, it is still likely to be relevant at the population level.^(51,52) For example, a recent modeling analysis has demonstrated that even a modest 0.1 SD increase in the BMD distribution across a population may reduce the overall burden of hip fracture by around 7%.⁽⁵³⁾ Given that there are more than 320,000 hip fractures annually in the USA,⁽⁵⁴⁾ although such extrapolation is

speculative, it demonstrates that a relatively small difference has the potential for substantial change to the burden of disease in a large population. Thus, although direct clinical implementation of these findings would be currently inappropriate based on this initial analysis, the results in a general sense point toward potential benefits for foods associated with low dietary inflammatory potential. This notion is consistent with the wider observation of inverse relationships between dietary inflammatory load and markers of a healthy diet with correlations between scores around -0.5 to -0.7 .⁽⁵⁵⁻⁶²⁾ Indeed, Western diets rich in red meat, high fat, sugar, dairy products, and refined grains have been associated with greater inflammatory load.⁽⁶⁻⁸⁾ Such findings support the importance of using other approaches to dietary evaluation, such as the “prudent” dietary pattern derived in the SWS mothers, characterizing a healthy diet as one based on high consumption of fruit and vegetables, high fiber, and low intake of saturated fats.^(9,45-47,63)

In summary, we have shown that diets with a greater inflammatory component, measured in both late pregnancy and in offspring childhood, are associated with lower bone mineral density in the offspring at 9 years old, with associations at least partly mediated through height and/or body weight. These observations further support the importance of maternal and early childhood diet on the longitudinal bone health of offspring, which may contribute to later musculoskeletal disease and fracture risk in older age.

Disclosures

JRH owns controlling interest in Connecting Health Innovations LLC (CHI), a company that has licensed the right to his invention of the dietary inflammatory index (DII) from the University of South Carolina in order to develop computer and smart phone applications for patient counseling and dietary intervention in clinical settings. NS is an employee of CHI. The subject matter of this article will not have any direct bearing on that work, nor has that activity exerted any influence on this project. All other authors declare no conflicts of interest in relation to this work.

Acknowledgments

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole SWS and ALSPAC teams, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. Please note that the ALSPAC study website (<http://www.bristol.ac.uk/alspac/researchers/our-data/>) contains details of all the data that are available through a fully searchable data dictionary and variable search tool.

The UK Medical Research Council and the Wellcome Trust (grant no. 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. Wellcome Trust grants 079960 and 084632 funded the pQCT scans. The SWS work was supported by grants from Medical Research Council (MRC), Bupa Foundation, British Heart Foundation, National Institute for Health Research (NIHR) Southampton Biomedical Research Centre, University of Southampton, and University Hospital Southampton NHS Foundation Trust, NIHR Oxford Biomedical Research Centre, University of Oxford, and the UK Royal Osteoporosis Society Osteoporosis and Bone Research Academy. NCH is supported by the UK Medical Research Council (MC_PC_21003;

MC_PC_21001). EMC has been supported by the Wellcome Trust (201268/Z/16/Z). The work leading to these results was supported by the European Union’s Seventh Framework Programme (FP7/2007–2013), projects EarlyNutrition, ODIN, and LifeCycle under grant agreements numbers 289346, 613977, and 733206, and by the BBSRC (HDHL-Biomarkers, BB/P028179/1 and BB/P028187/1), as part of the ALPHABET project, supported by an award made through the ERA-Net on Biomarkers for Nutrition and Health (ERA HDHL), Horizon 2020 grant agreement number 696295. CMP was supported by an award from Science Foundation Ireland, Ireland (grant no. SFI/16/ERA-HDHL/3360).

Authors’ roles: SW: conceptualization, data curation, investigation, methodology, project administration, writing—original draft, and writing—review & editing. SD: formal analysis, methodology, writing—original draft, and writing—review & editing. GM: data curation, formal analysis, investigation, methodology, writing—original draft, and writing—review & editing. EMC: conceptualization, methodology, project administration, writing—original draft, and writing—review & editing. SA: investigation, project administration, and writing—review & editing. NS: conceptualization, data curation, methodology, resources, and writing—review & editing. JH: formal analysis, funding acquisition, methodology, resources, and writing—review & editing. SRC: formal analysis, methodology, supervision, and writing—review & editing. CP: conceptualization, data curation, funding acquisition, investigation, resources, and writing—review & editing. MS: conceptualization, data curation, formal analysis, investigation, methodology, resources, and writing—review & editing. CR: conceptualization, data curation, funding acquisition, methodology, project administration, and writing—review & editing. CC: conceptualization, data curation, funding acquisition, project administration, resources, supervision, and writing—review & editing. NCH: conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, supervision, writing—original draft, and writing—review & editing.

AUTHOR CONTRIBUTIONS

Stephen J Woolford: Conceptualization; data curation; investigation; methodology; project administration; writing – original draft; writing – review and editing. **Stefania D’Angelo:** Formal analysis; methodology; writing – original draft; writing – review and editing. **Giulia Mancano:** Data curation; formal analysis; investigation; methodology; writing – original draft; writing – review and editing. **Elizabeth M Curtis:** Conceptualization; methodology; project administration; writing – original draft; writing – review and editing. **Shanze Ashai:** Investigation; project administration; writing – review and editing. **Nitin Shivappa:** Conceptualization; data curation; methodology; resources; writing – review and editing. **James R Hébert:** Formal analysis; funding acquisition; methodology; resources; writing – review and editing. **Sarah R Crozier:** Formal analysis; methodology; supervision; writing – review and editing. **Catherine M Phillips:** Conceptualization; data curation; funding acquisition; investigation; resources; writing – review and editing. **Matthew Suderman:** Conceptualization; data curation; formal analysis; investigation; methodology; resources; writing – review and editing. **Caroline L Relton:** Conceptualization; data curation; funding acquisition; methodology; project administration; writing – review and editing. **Cyrus Cooper:** Conceptualization; data curation; funding acquisition; project administration; resources; supervision; writing – review and editing. **Nicholas C Harvey:**

Conceptualization; formal analysis; funding acquisition; investigation; methodology; project administration; supervision; writing – original draft; writing – review and editing.

DATA AVAILABILITY STATEMENT

All data access enquiries should be made to Professor Cyrus Cooper, Director, MRC Lifecourse Epidemiology Centre, University of Southampton: cc@mrc.soton.ac.uk

References

- Orchard T, Yildiz V, Steck SE, et al. Dietary inflammatory index, bone mineral density, and risk of fracture in postmenopausal women: results from the Women's Health Initiative. *J Bone Miner Res.* 2017; 32(5):1136-1146.
- Cauley JA, Danielson ME, Boudreau RM, et al. Inflammatory markers and incident fracture risk in older men and women: the Health Aging and Body Composition Study. *J Bone Miner Res.* 2007;22(7):1088-1095.
- Barbour KE, Boudreau R, Danielson ME, et al. Inflammatory markers and the risk of hip fracture: the Women's Health Initiative. *J Bone Miner Res.* 2012;27(5):1167-1176.
- Sokolove J, Lepus CM. Role of inflammation in the pathogenesis of osteoarthritis: latest findings and interpretations. *Therapeu Adv Muscul Skel Dis.* 2013;5(2):77-94.
- Ricker MA, Haas WC. Anti-inflammatory diet in clinical practice: a review. *Nutr Clin Pract.* 2017;32(3):318-325.
- Khan S, Wirth MD, Ortaglia A, et al. Design, development and construct validation of the children's dietary inflammatory index. *Nutrients.* 2018;10(8):993.
- Hickling S, Hung J, Knuiman M, Divitini M, Beilby J. Are the associations between diet and C-reactive protein independent of obesity? *Prev Med.* 2008;47(1):71-76.
- Lopez-Garcia E, Schulze MB, Manson JE, et al. Consumption of (n-3) fatty acids is related to plasma biomarkers of inflammation and endothelial activation in women. *J Nutr.* 2004;134(7):1806-1811.
- Shivappa N, Steck SE, Hurley TG, Hussey JR, Hébert JR. Designing and developing a literature-derived, population-based dietary inflammatory index. *Public Health Nutr.* 2014;17(8):1689-1696.
- Shivappa N, Steck SE, Hurley TG, et al. A population-based dietary inflammatory index predicts levels of C-reactive protein in the seasonal variation of blood cholesterol study (SEASONS). *Public Health Nutr.* 2014;17(8):1825-1833.
- Hébert JR, Shivappa N, Wirth MD, Hussey JR, Hurley TG. Perspective: the dietary inflammatory index (DII)—lessons learned, improvements made, and future directions. *Adv Nutr.* 2019;10(2):185-195.
- Na W, Park S, Shivappa N, Hébert JR, Kim MK, Sohn C. Association between inflammatory potential of diet and bone-mineral density in Korean postmenopausal women: data from fourth and fifth Korea National Health and Nutrition Examination Surveys. *Nutrients.* 2019; 11(4):885.
- Mazidi M, Shivappa N, Wirth M, Hebert J, Vatanparast H, Kengne A. The association between dietary inflammatory properties and bone mineral density and risk of fracture in US adults. *Eur J Clin Nutr.* 2017;71(11):1273-1277.
- Cervo MM, Shivappa N, Hebert JR, et al. Longitudinal associations between dietary inflammatory index and musculoskeletal health in community-dwelling older adults. *Clin Nutr.* 2020;39(2):516-523.
- García-Arellano A, Ramallal R, Ruiz-Canela M, et al. Dietary inflammatory index and incidence of cardiovascular disease in the PREDIMED study. *Nutrients.* 2015;7(6):4124-4138.
- Ramallal R, Toledo E, Martinez-Gonzalez MA, et al. Dietary inflammatory index and incidence of cardiovascular disease in the SUN cohort. *PLoS One.* 2015;10(9):e0135221.
- Shivappa N, Godos J, Hébert JR, et al. Dietary inflammatory index and colorectal cancer risk—a meta-analysis. *Nutrients.* 2017;9(9):1043.
- Shivappa N, Hebert JR, Neshatbini Tehrani A, Bayzai B, Naja F, Rashidkhani B. A pro-inflammatory diet is associated with an increased odds of depression symptoms among Iranian female adolescents: a cross-sectional study. *Front Psych.* 2018;9:400.
- Chen L-W, Aubert AM, Shivappa N, et al. Associations of maternal dietary inflammatory potential and quality with offspring birth outcomes: an individual participant data pooled analysis of 7 European cohorts in the ALPHABET consortium. *PLoS Med.* 2021;18(1): e1003491.
- Vansant G, Wallace S, Godderis L. Effect of maternal and paternal nutrition on DNA methylation in the offspring: a systematic review of human and animal studies. *Adv Obes Weight Manag Control.* 2016;4(4):81-97.
- Harvey N, Dennison E, Cooper C. Osteoporosis: a lifecourse approach. *J Bone Miner Res.* 2014;29(9):1917-1925.
- Dennison EM, Syddall HE, Sayer AA, Gilbody HJ, Cooper C. Birth weight and weight at 1 year are independent determinants of bone mass in the seventh decade: the Hertfordshire cohort study. *Pediatr Res.* 2005;57(4):582-586.
- Hernandez C, Beaupre G, Carter D. A theoretical analysis of the relative influences of peak BMD, age-related bone loss and menopause on the development of osteoporosis. *Osteoporos Int.* 2003;14(10): 843-847.
- Cooper C, Eriksson JG, Forsen T, Osmond C, Tuomilehto J, Barker DJ. Maternal height, childhood growth and risk of hip fracture in later life: a longitudinal study. *Osteoporos Int.* 2001;12(8):623-629.
- Curtis E, Suderman M, Phillips C, Relton C, Harvey N. Early-life dietary and epigenetic influences on childhood musculoskeletal health: update on the UK component of the ALPHABET project. *Nutr Bull.* 2018;43(2):158-173.
- Inskip HM, Godfrey KM, Robinson SM, et al. Cohort profile: the Southampton Women's Survey. *Int J Epidemiol.* 2006;35(1):42-48.
- Fraser A, Macdonald-Wallis C, Tilling K, et al. Cohort profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol.* 2013;42(1):97-110.
- Boyd A, Golding J, Macleod J, et al. Cohort profile: the 'children of the 90s'—the index offspring of the Avon longitudinal study of parents and children. *Int J Epidemiol.* 2013;42(1):111-127.
- Cavicchia PP, Steck SE, Hurley TG, et al. A new dietary inflammatory index predicts interval changes in serum high-sensitivity C-reactive protein. *J Nutr.* 2009;139(12):2365-2372.
- Marx W, Veronese N, Kelly JT, et al. The dietary inflammatory index and human health: an umbrella review of meta-analyses of observational studies. *Adv Nutr.* 2021;12(5):1681-1690.
- Chen L-W, Aubert AM, Shivappa N, et al. Maternal dietary quality, inflammatory potential and childhood adiposity: an individual participant data pooled analysis of seven European cohorts in the ALPHABET consortium. *BMC Med.* 2021;19(1):1-14.
- Robinson S, Godfrey K, Osmond C, Cox V, Barker D. Evaluation of a food frequency questionnaire used to assess nutrient intakes in pregnant women. *Eur J Clin Nutr.* 1996;50(5):302-308.
- Rogers I, Emmett P. Diet during pregnancy in a population of pregnant women in south West England. ALSPAC study team. Avon Longitudinal Study of Pregnancy and Childhood. *Eur J Clin Nutr.* 1998; 52(4):246-250.
- Crabtree NJ, Arabi A, Bachrach LK, et al. Dual-energy X-ray absorptiometry interpretation and reporting in children and adolescents: the revised 2013 ISCD Pediatric Official Positions. *J Clin Densitom.* 2014;17(2):225-242.
- Normand SL. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med.* 1999;18(3):321-359.
- Sen S, Rifas-Shiman SL, Shivappa N, et al. Associations of prenatal and early life dietary inflammatory potential with childhood adiposity and cardiometabolic risk in project viva. *Pediatr Obes.* 2018;13(5): 292-300.
- Moore BF, Sauder KA, Starling AP, et al. Proinflammatory diets during pregnancy and neonatal adiposity in the healthy start study. *J Pediatr.* 2018;195(121-7):e2.
- Navarro P, Shivappa N, Hébert JR, et al. Predictors of the dietary inflammatory index in children and associations with childhood

- weight status: a longitudinal analysis in the lifeways cross-generation cohort study. *Clin Nutr*. 2020;39(7):2169-2179.
39. Navarro P, Shivappa N, Hébert JR, et al. Intergenerational associations of dietary inflammatory index with birth outcomes and weight status at age 5 and 9: results from the Lifeways Cross-generation Cohort Study. *Pediatr Obes*. 2020;15(3):e12588.
 40. Ingvorsen C, Brix S, Ozanne S, Hellgren L. The effect of maternal inflammation on foetal programming of metabolic disease. *Acta Physiol*. 2015;214(4):440-449.
 41. Sureshchandra S, Marshall NE, Wilson RM, et al. Inflammatory determinants of pregravid obesity in placenta and peripheral blood. *Front Physiol*. 2018;9:1089.
 42. Girard S, Tremblay L, Lepage M, Sébire G. IL-1 receptor antagonist protects against placental and neurodevelopmental defects induced by maternal inflammation. *J Immunol*. 2010;184(7):3997-4005.
 43. Challier J, Basu S, Bintein T, et al. Obesity in pregnancy stimulates macrophage accumulation and inflammation in the placenta. *Placenta*. 2008;29(3):274-281.
 44. Martin R, Harvey N, Crozier S, et al. Placental calcium transporter (PMCA3) gene expression predicts intrauterine bone mineral accrual. *Bone*. 2007;40(5):1203-1208.
 45. Crozier SR, Robinson SM, Borland SE, Inskip HM. Dietary patterns in the Southampton Women's Survey. *Eur J Clin Nutr*. 2006;60(12):1391-1399.
 46. Ganpule A, Yajnik CS, Fall CH, et al. Bone mass in Indian children—relationships to maternal nutritional status and diet during pregnancy: the Pune Maternal Nutrition Study. *J Clin Endocrinol Metab*. 2006;91(8):2994-3001.
 47. Winzenberg T, Shaw K, Fryer J, Jones G. Effects of calcium supplementation on bone density in healthy children: meta-analysis of randomised controlled trials. *BMJ*. 2006;333(7572):775.
 48. Lewis JR, Voortman T, Ioannidis JP. Evaluating and strengthening the evidence for nutritional bone research: ready to break new ground? *J Bone Miner Res*. 2021;36(2):219-226.
 49. Bianchi ML, Baim S, Bishop NJ, et al. Official positions of the International Society for Clinical Densitometry (ISCD) on DXA evaluation in children and adolescents. *Pediatr Nephrol*. 2010;25(1):37-47.
 50. Phillips CM, Chen L-W, Heude B, et al. Dietary inflammatory index and non-communicable disease risk: a narrative review. *Nutrients*. 2019;11(8):1873.
 51. Curtis EM, Harvey NC, D'Angelo S, et al. Bone mineral content and areal density, but not bone area, predict an incident fracture risk: a comparative study in a UK prospective cohort. *Arch Osteopor*. 2016;11(1):39.
 52. Johnell O, Kanis JA, Oden A, et al. Predictive value of BMD for hip and other fractures. *J Bone Miner Res*. 2005;20(7):1185-1194.
 53. Harvey NC, Kanis JA, Liu E, et al. Impact of population-based or targeted BMD interventions on fracture incidence. *Osteoporos Int*. 2021;32(10):1973-1979.
 54. Healthcare Cost and Utilization Project (HCUPnet). Rockville, MD, USA: Agency for Healthcare Research and Quality; 2017. <https://www.ahrq.gov/data/hcup/index.html>
 55. Wirth MD, Hébert JR, Shivappa N, et al. Anti-inflammatory dietary inflammatory index scores are associated with healthier scores on other dietary indices. *Nutr Res*. 2016;36(3):214-219.
 56. Shivappa N, Hebert JR, Kivimaki M, Akbaraly T. Alternative healthy eating index 2010, dietary inflammatory index and risk of mortality: results from the Whitehall II Cohort Study and meta-analysis of previous dietary inflammatory index and mortality studies. *Br J Nutr*. 2017;118(3):210-221.
 57. Kim Y, Chen J, Wirth MD, Shivappa N, Hebert JR. Lower dietary inflammatory index scores are associated with lower glycemic index scores among college students. *Nutrients*. 2018;10(2):182.
 58. Asadi Z, Ghaffarian Zirak R, Yaghooti Khorasani M, et al. Dietary inflammatory index is associated with healthy eating index, alternative healthy eating index, and dietary patterns among Iranian adults. *J Clin Lab Anal*. 2020;34(12):e23523.
 59. Turner-McGrievy G, Wirth MD, Hill KL, Dear ER, Hébert JR. Examining commonalities and differences in food groups, nutrients, and diet quality among popular diets. *Clin Nutr ESPEN*. 2021;41:377-385.
 60. Jandari S, Mosalmanzadeh N, Shadmand Foumani Moghadam MR, et al. Dietary inflammatory index and healthy eating index-2015 are associated with rheumatoid arthritis. *Public Health Nutr*. 2021;24(18):6007-6014.
 61. Hodge AM, Karim MN, Hébert JR, Shivappa N, de Courten B. Association between diet quality indices and incidence of type 2 diabetes in the Melbourne Collaborative Cohort Study. *Nutrients*. 2021;13(11):4162.
 62. Kranz S, Hasan F, Kennedy E, et al. Diet quality and dietary inflammatory index score among women's cancer survivors. *Int J Environ Res Public Health*. 2022;19(4):1916.
 63. Cole ZA, Gale CR, Javaid MK, et al. Maternal dietary patterns during pregnancy and childhood bone mass: a longitudinal study. *J Bone Miner Res*. 2009;24(4):663-668.