




BMJ Open Cost-effectiveness of emergency preparedness measures in response to infectious respiratory disease outbreaks: a systematic review and econometric analysis

Constantine Vardavas,¹ Katerina Nikitara ,¹ Konstantinos Zisis,¹ Konstantinos Athanasakis,² Revati Phalkey,³ Jo Leonardi-Bee,³ Helen Johnson ,⁴ Svetla Tsoлова,⁵ Massimo Ciotti,⁵ Jonathan E Suk ⁵

To cite: Vardavas C, Nikitara K, Zisis K, *et al.* Cost-effectiveness of emergency preparedness measures in response to infectious respiratory disease outbreaks: a systematic review and econometric analysis. *BMJ Open* 2021;**11**:e045113. doi:10.1136/bmjopen-2020-045113

► Prepublication history and supplemental material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-045113>).

Received 30 September 2020
Revised 25 January 2021
Accepted 23 February 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Jonathan E Suk;
jonathan.suk@ecdc.europa.eu

ABSTRACT

Objectives Respiratory infectious disease outbreaks pose a threat for loss of life, economic instability and social disruption. We conducted a systematic review of published econometric analyses to assess the direct and indirect costs of infectious respiratory disease outbreaks that occurred between 2003 and 2019.

Setting Respiratory infectious disease outbreaks or public health preparedness measures or interventions responding to respiratory outbreaks in OECD countries (excluding South Korea and Japan) so as to assess studies relevant to the European context. The cost-effectiveness of interventions was assessed through a dominance ranking matrix approach. All cost data were adjusted to the 2017 Euro, with interventions compared with the null. We included data from 17 econometric studies.

Primary and secondary outcome measures Direct and indirect costs for disease and preparedness and/or response or cost-benefit and cost-utility were measured. **Results** Overall, the economic burden of infectious respiratory disease outbreaks was found to be significant to healthcare systems and society. Indirect costs were greater than direct costs mainly due to losses of productivity. With regard to non-pharmaceutical strategies, prehospitalisation screening and the use of protective masks were identified as both an effective strategy and cost-saving. Community contact reduction was effective but had ambiguous results for cost saving. School closure was an effective measure, but not cost-saving in the long term. Targeted antiviral prophylaxis was the most cost-saving and effective pharmaceutical intervention.

Conclusions Our cost analysis results provide evidence to policymakers on the cost-effectiveness of pharmaceutical and non-pharmaceutical intervention strategies which may be applied to mitigate or respond to infectious respiratory disease outbreaks.

INTRODUCTION

Emerging, re-emerging and endemic respiratory and influenza-like infectious diseases represent a threat for loss of life, economic

Strengths and limitations of this study

- A systematic approach was followed, and the assessment of data quality indicated that the majority of studies included were of high quality.
- The synthesis of the results was performed using the dominance ranking matrix approach, which allowed for a direct comparison of the cost-effectiveness of each intervention to the null.
- Costs and resources varied between different countries, different regional settings and over time, making the cost component comparison of cost-effectiveness measures complex to interpret.
- We only focused on EU and OECD analogous countries excluding Japan and South Korea, and hence our cost-effectiveness analyses are not applicable to other countries or settings.
- Discrepancies in context and populations likely affect the implementation and efficacy of interventions.
- This study was conducted prior to the COVID-19 pandemic.

instability and social disruption as they can rapidly spread within communities and across countries, affecting the whole globe. Annually, it is estimated that 5%–15% of the population will suffer from influenza-related respiratory tract infections, while 3–5 million people face severe illness due to influenza.¹ In 2018, a total number of 109.5 million influenza virus episodes were identified among children under 5 years globally, with approximately 34 800 overall deaths. In Europe, seasonal influenza is estimated to lead to 4–50 million symptomatic cases and 15 000–70 000 deaths annually; however, this may differ between years, as the severe 2017/2018 influenza season led to an estimated 152 000 deaths in Europe alone.^{2 3}

In order for robust national preparedness systems and response strategies to outbreaks to be established in the Europe, it is crucial for public health officers to receive recent data of the health impact and the economic burden of respiratory infectious disease outbreaks in contrast to emergency response and preparedness actions. This evidence will ensure well-informed decisions regarding, among others, the proper allocation of resources.^{4,5} To this extent, although there is substantial literature from previously published systematic reviews on the value of public health emergency preparedness, they either refer to an older timeframe⁶ or use mathematical models to predict the effectiveness and cost-effectiveness of measures.⁷ Hence, there is limited recent information on the economic evaluations of infectious respiratory disease outbreaks that provide an overview of the cost effectiveness of response measures.⁸

Within the above context, the aim of this systematic review of econometric analyses was to assess the economic impact of response and preparedness measures when contrasted with the cost of infectious respiratory disease outbreaks. We further synthesise the cost-effectiveness for each intervention using a dominance ranking matrix (DRM) approach.

METHODS

Search strategy and selection criteria

A comprehensive systematic literature review of published econometric analyses was conducted between July and August 2019 using the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines⁹ and the Consolidated Health Economic Evaluation Reporting Standards (CHEERS)¹⁰ to identify peer-reviewed articles using two biomedical literature databases (PUBMED and EMBASE) and two economic literature databases (ECONLIT, IDEAS REPEC). The search strategy was designed for a broader study aiming to identify econometric studies on all types of infectious diseases, but due to the outbreak of COVID-19, and for the purposes of this specific article, we retained only those referring to respiratory infectious diseases. The complete search strategy and search terms are available in online supplemental appendix 1.

The inclusion criteria were as follows:

- ▶ Exposure: respiratory infectious disease outbreaks or public health preparedness measures or interventions responding to respiratory outbreaks in OECD countries (excluding Asian countries South Korea and Japan due to the wide cultural differences with the EU context as this study was performed under contract for the European Center for Disease Prevention and Control (ECDC)).
- ▶ Comparator: (i) no intervention (cost of inaction) or current practice, (ii) cost of preparedness versus cost of response (for studies reporting cost and benefit of public health preparedness).

- ▶ Outcome measures: direct and indirect costs for disease and preparedness and/or response or cost-benefit and cost-utility. Typical outcome measures of economic evaluations included: life years gained or cost per life-year gained with the intervention under investigation when incremental costs are combined, cost per quality-adjusted life year (QALY) gained, cases averted, monetary outcomes.
- ▶ Perspective: all direct and indirect costs pertaining to all relevant perspectives (eg, individual, hospital, insurance and societal—including national and regional) and all direct and indirect costs pertaining to all relevant perspectives according to York Health Economics Consortium¹¹ (health system perspective, including hospital, public health units; societal perspective; governmental perspective).
- ▶ Study designs: all relevant analytical epidemiological designs which estimate cost either as full economic evaluation studies, including cost-minimisation, cost-effectiveness, cost-utility and cost-benefit studies; cost-outcome and economic modelling studies or partial economic evaluations.
- ▶ Timeframe: from 2003 until August 2019, to reflect the timepoint from the 2003 SARS outbreak and onward¹²—this review refers to the pre-COVID-19 published evidence.

Studies that met the above inclusion criteria but did not report or perform any econometric analysis were excluded.

Data analysis and extraction

Studies identified from the searches were uploaded into a bibliographic database in which duplicate entries were removed. Initially, a pilot training screening process was used, where a random sample of 100 titles and abstracts were screened independently for eligibility by four reviewers (KN, KZ, RP, JLB) to enable consistency in screening and identify areas for amendments in the inclusion criteria. Following this, a random sample of 50% of titles and abstracts was screened independently by two reviewers. Since a high measure of inter-rater agreement was achieved (percentage agreement >88.7% and/or Cohen's Kappa >0.646), the remaining titles and abstracts were screened for eligibility by one reviewer. Where insufficient information was available in the title and abstract to make a decision, the full-text article of the document was retrieved for further inspection. Full-text documents of potentially eligible studies were retrieved for the records marked for inclusion. All full-text documents were independently double-screened by two reviewers, and inter-rater agreement measures were calculated at 88.3%. Disagreements in every step of the process were subsequently discussed and agreed on. Documents that passed the inclusion criteria on the basis of the full-text screening were included in the current review.

Appraisal of methodological quality

For evaluating the methodological quality of the included studies, the Consensus on Health Economic Criteria (CHEC) checklist¹³ was used. This specific tool has been designed for the assessment of full economic evaluations and includes 19 items (questions) with answers of 'Yes' or 'No'. For each positive answer on full economic evaluation studies, a single point was being assigned for the methodological quality, with a maximum score of 19. For the quality appraisal of partial economic evaluations, we used items from the CHEC checklist that were applicable—hence, the maximum score was 16. The quality appraisal process was completed by two reviewers, with a percentage of agreement in the three pilot studies, initially assessed by both, of 83.7%.

Comparative economic analysis approach

All cost data were adjusted to a common currency (Euro in 2017 (€²⁰¹⁷)) and price year using the Campbell and Cochrane Economics Methods Group—Evidence for Policy and Practice Information and Coordinating Centre cost converter.¹⁴ We adjusted the original estimate of cost from the original price year to a target price year of the €²⁰¹⁷, using a gross domestic product deflator index (GDPD), obtained from the International Monetary Fund World Economic Outlook Database GDPD index data set.¹⁵ Subsequently, we converted the price-year adjusted cost estimate from the original currency to €²⁰¹⁷, using conversion rates based on purchasing power parities (PPP) for GDP (the 2017 implied conversion factor was US\$1=€1.13, the €²⁰¹⁷ conversion factor was €1=US\$1.2, while with regard to British pounds, the conversion factor was £1=€0.88). PPP values adjust appropriately for differences in current price levels between countries, thus allowing comparisons based on a common set of average international prices; this is an advantage over pure exchange-rate conversions and GDP per capita approaches as PPPs eliminate differences in price levels between countries in the process of conversion. For studies that did not state the year of cost calculation, the costs were calculated 1 year before the publication year of each respective study.

Synthesis of cost-effectiveness

In order to synthesise the cost-effectiveness results, the DRM approach was used, which is a classification system developed for summarising and interpreting the results of economic evaluations in systematic reviews.¹⁶ The DRM is a three-by-three matrix with the following classification options:

1. Strong dominance for the intervention when the incremental cost-effectiveness measure shows the intervention compared with no intervention as: (i) more effective and less costly or (ii) as effective and less costly or (iii) more effective and equal cost.
2. Weak dominance for the intervention when the measure shows the intervention compared with no intervention as: (iv) effective and equally costly or (v) more

effective and more costly or (vi) less effective and less costly.

3. Non-dominance for the intervention when the measure shows the intervention compared with no intervention as: (vii) less effective and more costly or (viii) less effective and equally as costly or (ix) as effective and more costly.

Within our DRM, only studies that compared interventions to no intervention were included in the matrix.

Patient and public involvement

This study was performed under contract for the European Center for Disease Prevention and Control (ECDC). Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

RESULTS

The initial study search yielded 20 513 studies after removal of the duplicates and according to the specified selection criteria, only 66 were further assessed for eligibility via full text. Through the assessment of the full-texts, 52 studies were excluded for the following reasons: inadequate data on costs and/or cost-effectiveness (n=2), they were reviews (n=15), not referring to respiratory outbreaks (n=29), not referring to outbreaks of infectious diseases (n=2) and conference abstracts with no full text available (n=4). Additionally, three full-text papers were identified through the screening of the reference lists of the selected manuscripts, and hence, a total number of 17 econometric studies were considered in our analysis. The flowchart of the study selection process is presented in figure 1.

Overall, 11 out of the 17 studies were of high methodological quality (>80%), 5 were categorised as of good quality (60%–80%) and only 1 was of medium quality (40%–60%) due to missing quality criteria not mentioned by the authors including the comparative intervention, sensitivity analysis, incremental costs and outcomes. Online supplemental appendix 2 presents the overall quality appraisal score, for studies related to cost of infectious disease outbreaks and for sources related to preparedness, preventive and response measures concerning infectious disease outbreaks. The quality appraisal of partial and full economic evaluation studies is in online supplemental appendices 3 and 4, respectively. It is important to note that for the studies where a partial economic evaluation was performed, we only performed calculations for the items of the quality appraisal tool that were applicable.

Comparative cost analysis of infectious respiratory disease outbreaks

Regarding infectious respiratory disease outbreaks, six studies were included.^{17–22} All studies referred to influenza as the disease, either relating to pandemic H1N1 or seasonal Influenza B. Geographically, the studies




Figure 1 Flowchart.

were performed in the USA,¹⁷ Spain,^{18 22} France,¹⁹ New Zealand and Australia.^{20 21} Five out of the six studies were observational in design (cross-sectional or retrospective) and used collected data;^{18–22} one study was based on a simulation model.¹⁷ Similarly, five out of the six studies assessed costs from a healthcare system perspective;^{17 18 20–22} however, societal (n=3),^{17–19} governmental (n=1)¹⁷ and payer (n=1)¹⁹ perspectives were also assessed. Discounting in costs was not necessary for any of the included studies as the implementation timeframe had a duration of less than 1 year, and sensitivity analyses were performed only in three studies.^{17 19 21} A detailed description of the characteristics of the included illness studies is presented in online supplemental appendix 5.

Table 1 presents an analytical overview of the direct and indirect costs associated with influenza outbreaks. Direct costs mainly refer to medical and healthcare costs related to the outbreaks, along with the costs of response measures. Indirect costs included the loss of income, the loss of business and the loss of productivity. The overall direct costs reported in the studies were calculated at the patient level where possible.

The most recent study was a simulation study by Prager *et al*,¹⁷ in which multiple scenarios were assessed through simulation models for the US population so as to estimate the total economic burden of pandemic influenza outbreaks in the USA, taking into account both the scenario of an adequately vaccinated population and the opposite. The results indicated that medical expenditures for a pandemic influenza outbreak could reach 83.2 billion €²⁰¹⁷ in the no vaccination scenario and 67.3 billion €²⁰¹⁷ in the vaccination scenario. Notably, for indirect cost estimations, vaccination in a pandemic scenario would reduce workday losses by 22.2 million days, when compared with no vaccination.

Silva *et al*¹⁹ focused on an influenza outbreak in France between 2010 and 2011 and extrapolated the results to the entire country with a hypothetical approximate number of 2 million influenza cases (3.2% of the French population), for which they calculated an overall cost of 151 million €²⁰¹⁷ for the French Health Insurance System. Direct costs per patient ranged between 35.26 €²⁰¹⁷ and 73.91 €²⁰¹⁷, with higher indirect costs of 97.88 €²⁰¹⁷ per

Table 1 Characteristics of cost of illness studies of influenza outbreaks*, expressed in Euros (base year 2017)

Study (Publication Year)	Setting, year	Perspective	Direct costs (€, 2017)	Indirect costs (€, 2017)
Prager <i>et al</i> (2017) ¹⁷	USA, n/a	Healthcare system, governmental, societal	Seasonal (no vaccination): €5.92 billion Seasonal (vaccination): €9.96 billion Pandemic (no vaccination): €81.18 billion Pandemic (vaccination): €65.59 billion	<i>Illness-related workdays losses</i> (a) Vaccination and no vaccination in a seasonal scenario: vaccination contributes to more workday losses than no vaccination (b) Vaccination and no vaccination in a pandemic scenario: vaccination reduces workday losses by 22.2 million days compared with no vaccination
Morales-Suárez-Varela <i>et al</i> (2016) ¹⁸	Spain, 2009–2010	Healthcare system, societal	<i>Total direct cost/patient</i> Non-pregnant women: €3 908.70 Pregnant women: €2 227.10	<i>Total indirect cost/patient</i> Non-pregnant women: €107.18 Pregnant women: €63.83
Silva <i>et al</i> (2014) ¹⁹	France, 2010–2011	Payer, societal	<i>Mean direct cost/patient</i> All ages—€53.43 0–4 years old—€73.91 5–14 years old—€52.79 15–65 years old—€35.26 ≥65 years old—€44.13 <i>Total direct costs</i> All ages—€107 883 835 0–4 years old—€18 908 254 5–14 years old—€52 474 781 15–65 years old—€21 590 741 ≥65 years old—€6 940 836	<i>Mean daily allowance cost due to work leave/patient</i> All ages—€22.38 0–4 years old—€0 5–14 years old—€0 15–65 years old—€97.88 ≥65 years old—€0
Higgins <i>et al</i> (2011) ²⁰	Australia and New Zealand, 2009	Healthcare system	Total mean cost: €19296 136 Total ICU costs: €6 107 069 Total non-ICU costs: €12 961 942 Mean cost of ICU/patient: €61 368 Mean cost of non-ICU/patient: €10 755 Mean cost in ICU/per patient and per day: €4 767	Non-reported
Wilson <i>et al</i> (2009) ²¹	New Zealand, 2009	Healthcare system	Total ICU costs: €40 807 660 Median ICU cost/patient: €22 540 Mean ICU cost/patient: €32 168 <i>Total hospital costs/patient</i> Median hospital cost: €39 696 Mean hospital cost: 53 553 <i>Treatment costs in ICU per subgroup</i> (a) Cost/patient with and without pre-existing comorbidity €16 100 and €28 980, respectively (b) Cost/patient with viral pneumonitis and with other influenza syndromes €22 212 and €12 880, respectively	Non-reported
Rodríguez-Rieiro <i>et al</i> (2009) ²²	Spain, 2009	Healthcare system	Total cost: €36 700 000 Median cost per hospitalisation (concomitant chronic disease): €2 205 Median cost per hospitalisation (without a medical condition): €1 172	Non-reported

The adjustment was performed from Canadian \$, US\$, Australian \$, British pounds £ and converted to Euro (Germany has been selected as target currency in these cases). Currencies from European Union countries adjusted to their currency.

The cost data include all forms of cost derived from inclusion studies, such as overall/total cost, mean/average cost, income loss, labour cost, household cost, savings, cost per case, etc.

For studies without currency year indicated, the previous year of publication was selected for adjustment.

*Confirmed or extrapolated/hypothetical cases on which they base the economic evaluation.
ICU, intensive care unit.

day due to absence from work, for those within the 15–65 age group.

Two studies assessed the cost of an influenza outbreak from an intensive care unit (ICU) and hospital perspective.^{20 21} One focused on ICU and hospital costs derived from an influenza pandemic in 2009 in New Zealand (among 1224 cases, of which 122 were admitted to ICUs), which surpassed 40.8 million €²⁰¹⁷ at an average cost of

32 167 €²⁰¹⁷ per patient, with significantly increased costs for patients with underlining comorbidities.²¹ The mean total hospitalisation cost (normal and ICU) per case surpassed 53 553 €²⁰¹⁷. Similarly, in a study that included 762 H1N1 cases from both Australia and New Zealand, the mean cost per ICU patient was 61 368 €²⁰¹⁷, with a per-day cost of 4767 €²⁰¹⁷.²⁰ On the contrary, the non-ICU patient had a mean cost of 10 755 €²⁰¹⁷; however, overall

non-ICU patient costs surpassed those of ICU patients (12.96 million €²⁰¹⁷ vs 6.1 million €²⁰¹⁷), leading to a total hospitalisation cost of 19.3 million €²⁰¹⁷ for the 2009 influenza outbreak.

Similarly, Rodriquez-Rieiro *et al*²² studied the hospitalisation costs that occurred during the 2009 influenza pandemic in Spain, which reached 36.7 million €²⁰¹⁷ for 11 449 hospitalisations—during which the appearance of comorbidities led to higher average costs per patient (2205 €²⁰¹⁷ vs 1172 €²⁰¹⁷, respectively). Specific populations in Spain were assessed by Morales-Suárez-Varela *et al*¹⁸ who estimated direct costs for medical visits, medication and diagnostic tests at €3908 €²⁰¹⁷ for non-pregnant women and 2227 €²⁰¹⁷ for pregnant women of reproductive age, with indirect costs estimated at 107 €²⁰¹⁷ and 64 €²⁰¹⁷, respectively.

Cost-effectiveness studies of measures in averting and/or responding to infectious respiratory disease outbreaks

We identified 11 studies^{23–33} referring to *preparedness, preventative and response measures*, to influenza outbreaks, presented in detail in online supplemental appendix 6. Two studies were observational (based in the Netherlands and the UK),^{23 24} and the remaining nine were simulation models (four US models, with one study each modelled for Canada, France, Australia, Israel and one referring to developed countries in general). All included studies either used a cost-effectiveness or a cost-utility economic evaluation approach. The studies' timeframes ranged from 2004 to 2018. Regarding the perspective for direct and indirect costs, a healthcare system or society approach was consistently presented.

The preparedness, preventive and response measures described included three pharmaceutical interventions (vaccination as a response measure, general vaccination, antiviral drug therapy and stockpiling),^{31–33} four non-pharmaceutical interventions (screening at the point of contact, community contact reduction, volunteer isolation/quarantine, school closure and the use of personal protective measures)^{23–25 28} and four combined pharmaceutical and non-pharmaceutical interventions.^{26 27 29 30}

Table 2 presents the details of the cost-effectiveness studies on preparedness and response measures for infectious respiratory disease outbreaks. Further details on the comparative analysis of health indexes gained when averting or responding to respiratory outbreaks can be found in online supplemental appendix 7.

With regard to studies that compared multiple interventions, a simulation model of pandemic influenza in the USA studied the cost-effectiveness of stockpile strategy and concluded that expanded adjuvanted vaccination seemed to be the most cost-effective strategy, averting 68% of infections and deaths and gaining 404 303 QALYs at \$10 844 (€9600 €²⁰¹⁷) per QALY gained relative to the stockpiling strategy.³⁰ Saunders-Hastings *et al*,²⁶ using a simulated population of 1.2 million people (reflective of Ottawa, Canada), performed a cost-effectiveness analysis of six interventions including vaccination, school closure,

antiviral prophylaxis and other measures. The authors concluded that vaccination was the most cost-effective intervention when compared with other interventions while the least cost-effective intervention was school closure in conjunction with community-contact reduction, personal protective measures, voluntary isolation and quarantine. In particular, the cost per life-year saved was estimated to be \$2581 (1700 €²⁰¹⁷) for combined vaccination and antiviral treatment, while an estimated cost of \$260 472/life-year saved (€171 590 €²⁰¹⁷) was noted for school closure in conjunction with other interventions. Finally, Halder *et al*²⁷ aimed to evaluate the most cost-effective strategies suitable for a future pandemic with H1N1 2009 characteristics in Australia. The results showed that the strategy with the lowest cost was the dual strategy of individual school closure for 2 weeks along with antiviral drug strategies, with a total cost of approximately AU\$632 (376.31 €²⁰¹⁷) per case averted. The strategy with the highest cost was the dual strategy of school closure along with the continuous—50% workplace closure, with a cost of \$103 million (61.3 million €²⁰¹⁷), per 100 000 population.

Comparative cost-effectiveness analysis

A DRM approach is presented in figure 2. These interventions include both pharmaceutical measures and non-pharmaceutical measures. The interventions were compared with the 'no intervention' scenario, with the exception of one study²⁹ in which the comparators were vaccination versus self-isolation, which was subsequently excluded from the DRM.

Pharmaceutical measures

Vaccination as a response measure

With the application of our inclusion and exclusion criteria, four studies assessed vaccination as a response measure in the context of an outbreak and included a cost analysis. Overall, as highlighted in the majority of the studies, vaccination as a response measure was noted to have a more significant clinical effect than comparators and was more cost-saving in most cases. According to Sander *et al*,³⁰ the most clinically effective intervention was expanded adjuvant vaccination which contributed to 404 030 QALYs. Similarly, Khazeni calculated that with expanded adjuvanted vaccination, 45 941 deaths would be averted.³¹ Additionally, Saunders-Hastings *et al*²⁶ concluded that the most cost-effective approach for controlling a pandemic was vaccination in combination with antiviral therapy and prophylaxis. However, a review of the results showed that much of the cost-effectiveness of pharmaceutical interventions were driven by vigorous vaccination campaigns, while the contribution of antiviral drugs' was not of significance. Finally, Madema *et al*³³ through a simulation model of an influenza pandemic among developing countries calculated the costs and assessed the effectiveness of two types of vaccines, an egg-based and a cell culture-based, in

Table 2 Characteristics of full economic evaluation studies on preparedness and response measures of influenza outbreaks, expressed in Euros (base year 2017)

Study (publication year)	Setting, Year	Population (n)	Interventions	Comparator	Economic evaluation outcomes
<i>Non-pharmaceutical studies</i>					
Lankelma <i>et al</i> (2019) ²³	Netherlands, 2017–2018	Patients with acute RTI at the emergency department (1546 tests, 624 cases)	Point-of-care-testing for Influenza before hospital admission	2016–2017 influenza season	<p>Net savings €388 317 (after subtraction with costs)</p> <p>More than 80% of the total savings are due to the shorter length of stay and decreased hospital admissions.</p> <p>The overall cost of intervention: €98 968</p> <p>Laboratory costs at €72 202</p> <p>Clinical aspects costs at €26 767</p>
Orset (2018) ²⁵	France, 2014	200 participants, data extrapolated	7-day home confinement	No intervention	<p>Costs associated with home confinement</p> <p>(a) Direct costs For adults: €742/case For elderly: €1191/case</p> <p>(b) Indirect costs Productivity losses/case For adults: €550. For elderly: €125</p> <p>Costs of death/case The cost of death for children is estimated at €22–128, for adults at €63–361 and for elderly at €2667–15 389</p> <p>Loss of productivity due to influenza/case Productivity loss in case of adult sickness: €88.70 (incl. absent from work+reduced productivity)</p> <p>Productivity loss in case of a sick child for the adult (mainly mother): €97.62</p>
Sadique <i>et al</i> (2008) ²⁴	UK, 2005	Working parents with depending children	School closure	No intervention	<p>Cost of school closure: between €280 million—€2.8 billion/week</p> <p>Cost of absenteeism: €1.4 billion</p> <p>Adjusting for informal care, the cost reduced between €552–€635 million per week.</p> <p>Adjusting for the elasticity of production the cost reduced to €970 327 320—€1.1 billion per week</p>
Tracht <i>et al</i> (2012) ²⁸	USA, 2009–2010 influenza season	Simulation of the USA (302 million people:73 million children, 191 million adults and 38 million seniors)	Population use of face masks (N95) on the spread of a pandemic	No intervention	<p>Net savings if masks are worn by 10% of the adult population: €418.75 billion if masks are worn by 50% of the adult population: €501.9 billion</p> <p>Economic burden, if no intervention: €728.28 billion (incl. direct and indirect costs)</p>
<i>Combined pharmaceutical and non-pharmaceutical strategies</i>					
Saunders-Hastings <i>et al</i> (2017) ²⁶	Canada, n/a	A simulation of Ottawa, Canada (1.2 million)	<ol style="list-style-type: none"> 1. Vaccination+antiviral treatment 2. Vaccination+antiviral treatment+antiviral prophylaxis 3. Community contact reduction+personal protective measures+isolation 4. Community-contact reduction+personal protective measures+isolation+antiviral treatment 5. School closure+community contact reduction+personal protective measures+quarantine 6. All interventions 	No intervention	<p>Cost/LYG vs no intervention</p> <ol style="list-style-type: none"> 1. €1700/LYG 2. €1769/LYG 3. €4394/LYG 4. €4447/LYG 5. €171 590/LYG 6. €131 679/LYG <p>Total economic burden For all scenarios, the economic burden ranges between €75 758 to €1 416 351</p>

Continued

Table 2 Continued

Study (publication year)	Setting, Year	Population (n)	Interventions	Comparator	Economic evaluation outcomes
Halder <i>et al</i> (2011) ²⁷	Australia, 2009	A community in Western Australia (30 000)	Different combinations of durations of individual school closure, antiviral treatment, household antiviral prophylaxis, extended antiviral prophylaxis, 50% workplace closure, 50% community contact reduction	No intervention	Cost/case averted: Antiviral drug strategies+2 weeks school closure: €396 per case averted (cost-effective) Short-duration school closure: €820/case averted ISC, continuously+50% workplace, continuously: €6 204/case averted In case of 2 weeks for the above combination: €1891/case averted ISC, continuously: €2180/case averted Total cost, per 100 000 population The dual strategy of individual school closure for 2 weeks (ISC) along with the 50% community contact reduction (CCR): €3.39million The dual strategy of continuous ISC along with the continuous—50% WP: €61.3million. Productivity loss due to illness and interventions per 100 000 population ISC (cont.)+WP (cont.): €90.21 million Combined antiviral treatment, household antiviral prophylaxis and extended antiviral prophylaxis: €4.63
Yarmand <i>et al</i> (2010) ²⁹	USA, 2009–2010 influenza season	North Carolina State University undergraduate students (23 087)	Vaccination	Self-isolation	High levels of interventions Self-isolation is incrementally more cost-effective than vaccination. This has been presented for most of cost ratio values. Low levels of interventions Vaccination is incrementally more cost-effective than self-isolation. The results were robust, even in sensitivity analyses.
Sander <i>et al</i> (2009) ³⁰	USA, n/a	Residents of a 1.632-million-person city	<ol style="list-style-type: none"> HTAP25 with a stockpile for 25% of the population HTAP50 with a stockpile for 50% of the population HTAP with an unlimited stockpile School closure for 26 weeks Prevaccination 70% of the population with a low efficacy vaccine HTAP25+school closure HTAP50+school closure HTAP+school closure Prevaccination+school closure: prevaccinating 70% population with the low-efficacy vaccine, plus closing all schools for 26 weeks Treatment only: Treating all cases with antivirals FTAP25 for household contacts and 60% of work/school contacts, stockpile for 25% of the population FTAP50 for household contacts and 60% of work/school contacts, stockpile for 50% of population FTAP for household contacts and 60% of work/school contacts, stockpile unlimited FTAP25+school closure FTAP50+school closure FTAP+school closure 	No intervention	Cost/capita and cost-effectiveness outcomes 1. FTAP is cost-effective (54% reduction attack rate, €119 per capita) 2. Prevaccination (48% reduction attack rate, €131 per capita) 3. School closure in combination with each of the above is the least cost-effective (€2 524 per capita) ICUR of FTAP: €42 959 ICUR of prevaccination and school closure: €43 106 Cost-saving FTAP and prepandemic vaccination are cost-saving compared with no intervention
<i>Pharmaceutical only strategies</i>					

Continued

Table 2 Continued

Study (publication year)	Setting, Year	Population (n)	Interventions	Comparator	Economic evaluation outcomes
Khazeni <i>et al</i> (2009) ³¹	USA, n/a	A US metropolitan city (8.3 million)	<ol style="list-style-type: none"> 1. Stockpiled strategy 2. Expanded adjuvanted vaccination 3. Expanded antiviral prophylaxis 	No intervention	<p><i>Intervention and treatment costs</i></p> <ol style="list-style-type: none"> 1. Stockpiled strategy: Total cost of €30.1 million and contribution to €288 million treatment costs 2. Expanded adjuvanted vaccination: Total cost of €179 million and contribution to €166 million treatment costs 3. Expanded antiviral prophylaxis: Total cost of €58.4 million and contribution to €266 million treatment costs 4. No intervention: contribution to €462 million treatment costs <p><i>Cost/QALY gained</i></p> <ol style="list-style-type: none"> 1. Stockpiled strategy compared with no intervention: €7894/QALY 2. Expanded adjuvanted vaccination (at 80% effectiveness) relative to stockpiled strategy: €8600/QALY 3. Expanded antiviral prophylaxis has a less favourable cost-effectiveness ratio than adjuvanted vaccination <p>Expanded adjuvanted vaccination shown to be a cost-effective intervention because it contributes to 404 030 QALYs at \$10 844 per QALY gained relative to stockpiled strategy.</p>
Balicer <i>et al</i> (2005) ³²	Israel, n/a	Population of Israel (1 618 200 cases/patients)	<p>Stockpiling with antiviral drugs</p> <ol style="list-style-type: none"> 1. Therapeutic use (all patients) 2. Therapeutic use (high-risk patients) 3. Pre-exposure long-term prophylaxis (all population) 4. Pre-exposure long-term prophylaxis (high-risk population) 5. Short-term postexposure prophylaxis for all close contacts 	No intervention	<p><i>CBA</i></p> <p>Therapeutic use (incl. all and high-risk patients): 2.44–3.68 Pre-exposure (incl. entire and high-risk population): 0.37–0.38 Postexposure: 2.49</p> <p>Stockpiling with antiviral drugs for high-risk patients remain cost-saving strategy even if the annual probability of a pandemic remains >1 every 80 years.</p> <p><i>Overall cost</i></p> <p>The overall health-related costs: €56 234 057 The overall cost to the economy: €535 245 986 <i>Workdays lost due to illness</i> 6 536 240 or 4 days/patient</p>
Medema <i>et al</i> (2004) ³³	n/a,	Developed countries (1 billion people)	<ol style="list-style-type: none"> 1. Egg-based vaccines with 17% population coverage 2. Cell culture-based vaccines with 37% population coverage 	No intervention	<p><i>Cost per life-year gained</i></p> <p>In general, vaccination is cost-effective. Cell culture-based vaccines: €3376/LYG (cost-effective) <i>Cost per intervention</i> Egg-based: €2.6 billion Cell culture-based: €5.87 billion <i>Net savings</i> Egg-based: €8.5 billion Cell culture-based: €5.87 billion Savings: €1.84 billion</p>

CBA, cost-benefit ratio; CCR, community contact reduction; FTAP, full targeted antiviral prophylaxis; HTAP, household targeted antiviral prophylaxis; ICER, incremental cost-effectiveness ratio; ICUR, incremental cost-utility ratio; ISC, individual school closure; LYG, life-year gained; QALY, quality-adjusted life year; VSL, value of statistical life; WP, workplace closure.

Type of Intervention	Costs*	Health benefit**
Pharmaceutical strategies		
Vaccination as a response measure		
Sander et al. (2009)	–	+
Khazeni et al. (2009)	0	+
Madema et al. (2004)	–	+
Saunders-Hastings et al. (2017)	–	+
General vaccination		
Sander et al. (2009)	–	+
Antiviral drugs		
Sander et al. (2009)	–	+
Halder et al. (2011)	–	+
Balicer et al. (2005)	–	+
Saunders-Hastings et al. (2017)	–	+
Khazeni et al. (2009)	0	+
Stockpile strategy		
Sander et al. (2009)	–	+
Balicer et al. (2005)	–	+
Khazeni et al. (2009)	0	+
Non-Pharmaceutical strategies		
Volunteer isolation		
Orset (2018)	–	+
Saunders-Hastings et al. (2017)	–	+
Pre hospitalisation screening		
Lankelma et al., (2019)	–	+
Community contact reduction		
Saunders-Hastings et al. (2017)	–	+
Halder et al. (2011)	+	+
School closure		
Sadique et al. (2008)	+	+
Saunders-Hastings et al. (2017)	+	+
Halder et al. (2011)	+	+
Sander et al. (2009)	+	+
Personal protective measures		
Tracht et al. (2012)	–	+
Saunders-Hastings et al. (2017)	–	+

Figure 2 Dominance ranking matrix for pharmaceutical and non-pharmaceutical strategies. *+: the intervention is less cost saving than the comparator; 0: the intervention is equally cost saving with the comparator; –: the intervention is more cost saving than the comparator. **+: The intervention is more effective than the comparator; 0: the intervention is equally effective with the comparator; –: the intervention is less effective than the comparator.

comparison with no intervention. Overall, vaccination was more cost-effective than no intervention; however, vaccination with cell culture-based vaccines was the most cost-effective strategy with a cost of 3779 €²⁰¹⁷ per life-year gained. General vaccination was also assessed by Sander *et al.*,³⁰ who noted it to be both more cost-saving and effective than the unmitigated pandemic scenario, although when comparing prevaccination with low-efficacy vaccines with full targeted antiviral prophylaxis, it was less effective and more costly.

Antiviral drugs

Antiviral drug strategies were assessed in five studies, where it was noted that they were both more effective and cost-saving than the no intervention scenario, primarily when used as targeted prophylaxis. According to Halder *et al.*,²⁷ antiviral drug strategies

such as antiviral treatment and antiviral treatment in combination with household confinement and extended prophylaxis can result in reduced attack rates of 7.6% and 3.5% in comparison to the unmitigated attack rate of 13%. The costs of these strategies are also lower than the cost of no intervention.

Moreover, therapeutic treatment and postexposure prophylaxis for exposed individuals (targeted prophylaxis) were shown to be the most cost-saving.³² Consistent with the above, antiviral therapy in combination with a layered non-pharmaceutical approach seemed to reduce the overall economic costs the most and was identified as more effective when compared with no intervention.²⁶ Furthermore, it was noted that expanded antiviral prophylaxis could help delay a pandemic when additional strategies are implemented

and would also lead to averting 32 745 deaths in the USA.³¹ Finally, Sander *et al*³⁰ used a stochastic simulation model of pandemic influenza in the USA, aiming to evaluate the potential economic impact of 16 different mitigation interventions from a societal perspective. Conclusively, targeted antiviral prophylaxis was both the most cost-saving and effective intervention with a cost of \$127 per capita (€118.73 €²⁰¹⁷), with the scenario of implementation of expanded antiviral prophylaxis leading to a total of 282 329 QALYs gained.

Stockpile strategy

The stockpile strategy was assessed in three of the studies included in this systematic review. Based on the findings, stockpiling antiviral prophylaxis in the context of a pandemic was noted to be both cost-saving for the society and avert loss of life compared with no intervention.³⁰ Moreover, prepandemic stockpiling of antiviral drugs would be more effective and cost-saving than no intervention if antiviral drugs were administered either solely as a treatment or as short-term prophylaxis for exposed individuals.³² Finally, stockpiling was also found more effective than a no intervention scenario (averting 29 761 deaths in the USA), although when compared with other interventions, expanded vaccination and prophylaxis were found to be more effective.³¹

Non-pharmaceutical measures

Pre hospitalisation screening

Lankelma *et al*²³ assessed the cost-effectiveness of screening patients with acute respiratory tract infection for influenza before hospital admission. Overall costs of screening were estimated at 98 968 €²⁰¹⁷ for 1546 tests and 624 cases and reported net savings of 388 317 €²⁰¹⁷ for the healthcare system. Point-of-care testing for influenza before hospital admission was identified as a cost-effective intervention.²³

Community contact reduction

Community contact reduction was assessed in two studies, where it was either implemented solely or in combination with other pharmaceutical and non-pharmaceutical measures. Home confinement was noted as cost-effective as a preventive measure in the context of influenza epidemics, if the proportion of compliance is adequate and infected individuals ask for medical assistance, regardless of the severity level of the pandemic.²⁶ Isolation of infected individuals was found to be among the most effective interventions, whereas combined with community contact reduction, personal protective measures and antiviral treatment, self-isolation had the lowest cost.²⁷

School closure

The effectiveness and the economic burden of school closure were evaluated in four studies, highlighting that the duration of school closure and potentially

combined strategies significantly affect its impact. Sadique *et al*²⁴ estimated the economic burden of school closure in the UK from a societal perspective and showed that the estimated costs of school closure were high, at 0.28–1.68 billion €²⁰¹⁷ per week and the authors concluded that school closure was likely to significantly add an extra economic burden on the health system through staff absenteeism, even if school closure may delay infectious disease transmission. Similarly, Sander *et al*,³⁰ who studied school closure as an additional intervention to full targeted antiviral prophylaxis or prevaccination found that while school closure further improves health outcomes (gaining 51 QALYs), it was the least cost-effective measure as it increased the total cost to society by \$2700 per capita (€2524 €²⁰¹⁷). Additionally, school closure produced only a small reduction in attack rate, whether implemented in combination with other interventions or alone.²⁶ Finally, exclusive school closure for 2 weeks along with the continuous 50% workplace closure, antiviral treatment, household antiviral prophylaxis and extended antiviral prophylaxis, had the lowest illness attack rate (2.4%) and one of the lowest costs. On the contrary, school closure as a sole intervention to counterbalance infectious respiratory diseases was not a cost-effective measure.²⁷

Personal protective measures

Personal protective measures such as face masks and hand hygiene were assessed in two of the included studies, noting that they could contribute to the control of a pandemic, dependant though on the exposed and susceptible individuals' compliance rate, the setting and the overall burden of the respiratory pandemic.^{26 28} Tracht *et al* aimed to assess the cost-effectiveness of facemasks (N95 grade) in reducing the spread of pandemic (H1N1) 2009, using a simulation model of the US population and identified an economic burden of 728.28 billion €²⁰¹⁷ (incl. direct and indirect costs). Notably, if masks are worn by 10% and 50% of the adult population of the US net savings were calculated at 418.75 billion €²⁰¹⁷ and 501.9 billion €²⁰¹⁷, respectively. Hence, the use of face masks were identified as a cost-effective preventive measure depending on the population's level of compliance.

DISCUSSION

The aim of this systematic literature review of economic analysis studies was to assess the economics of preparedness when contrasted with the cost of infectious respiratory disease outbreaks primarily within the context of European and OECD countries (excluding Japan and South Korea). Overall, the economic burden of infectious disease outbreaks is costly to healthcare systems, or to governments and society reflecting the medical costs for response activities including both the treatment of the confirmed cases and the surveillance

and elimination of the disease's transmission, as well as indirect costs which were also substantial.

In general, the majority of direct costs seemed to primarily reflect cost of additional personnel hours, which are mandatory for the management of the infected cases, for the organisation of response planning and contact tracing, for providing educational training and materials as well as laboratory costs. With regard to indirect costs, these could in many cases be greater than the direct costs, especially when school closures and/or workplace closures are enacted across a population, which in turn impact productivity and increase the economic burden.

While all the identified pharmaceutical and non-pharmaceutical interventions lead to a health benefit for the individual or the society, the cost benefit of such interventions differs. With regard to the potential non-pharmaceutical strategies, we identified that the use of personal protective measures, such as a facemask, is both cost-saving and effective, as also is prehospitalisation screening among suspect cases. On the other hand, all studies that assessed the impact of school closure noted that although it is an effective measure in reducing transmission, it is not cost-saving as it leads to increased economic burden. Moreover, when school closure was used as a sole intervention, then the use of limited duration school closure was significantly more cost-effective compared with continuous school closure.²⁴ Community contact reduction was identified to have a positive health impact but had ambiguous results with regard to its potential cost saving as one study²⁶ noted that it is a cost-saving intervention, while the other²⁷ noted that social distancing strategies, such as reduced workplace attendance, were not a cost-saving measure primarily due to productivity losses, especially during longer periods of closure. Productivity losses primarily were noted to arise from pandemic related deaths and illness coupled with those losses due to interventions such as workplace closure and child-care of an ill child. It is important to note that non-pharmaceutical strategies were mostly applied complementary with a pharmaceutical measure or in combination with other non-pharmaceutical strategies in order to enhance their effectiveness. However, their cost-effectiveness highly depended on the duration, the level of compliance from the population and the type and burden of the infectious disease. It should moreover be noted that cost-effectiveness of measures will vary depending on the epidemiology of the disease in question.

With regard to pharmaceutical interventions, vaccination as a rapid response measure for infected and suspected individuals was noted to have a more significant clinical effect than comparators and was more cost-saving in most cases. As for antiviral treatment, the majority of the findings noted that it is a cost-effective strategy, especially when combined with other pharmaceutical and non-pharmaceutical interventions or

when used as targeted prophylaxis for exposed individuals. Targeted antiviral prophylaxis was the most cost-saving and effective intervention, while stockpiling was cost saving in most cases and averted loss of life when compared with no intervention.

The current number of economic evaluation or cost-effectiveness studies of influenza outbreak preparedness measures is small, with an increase shown since the 2009 influenza pandemic; however, it is important to note that these studies refer to the evidence published before the COVID-19 pandemic. There are only a limited number of related reviews, however of different scope focusing primarily on policy recommendations³⁴ or used dynamic transmission models in the included economic assessments of pandemic influenza preparedness measures based on significantly older studies.⁶ Additionally, most of the existing review studies either evaluate the overall economic burden of the disease or the cost-effectiveness of different pharmaceutical and non-pharmaceutical interventions without necessarily them reflecting the economics of outbreaks of infectious respiratory diseases.

Placing the above into context and following the assessment of the methodological approaches used across studies, it is essential to note the minimum contents that economic outbreaks of respiratory studies should include to help inform future and upcoming research, especially in light of the COVID-19 pandemic. These include clearly noting of the study year, the population at risk and, the population infected, the type of economic perspective (ie, healthcare, societal, etc), the study timeframe and discounting, as well as detailed reporting of the direct and indirect costs of the respiratory outcome and the interventions applied.

Strengths and limitations

A significant strength of this review is the comprehensive approach that was followed and the assessment of data quality—which indicated that the majority of the studies included were of high quality. Second, the synthesis of the results was performed using the DRM approach, which allowed for a direct comparison of the cost-effectiveness of each intervention to the null intervention.

However, there are a few limitations: first, costs and resources varied between different countries, different regional settings and over time, making the cost component comparison of cost-effectiveness measures complex to interpret. Moreover, we only focused on EU and OECD analogous high-income countries excluding Japan and South Korea, and hence our cost-effectiveness analyses are not applicable and generalisable to other countries and particularly middle-income and low-income countries. Additionally, discrepancies in context and populations likely affect the implementation and efficacy of interventions, undermining even the effectiveness elements comparability in the cost-effectiveness measures,

especially in complex multi-component public health interventions. In addition, our study did not include studies published before 2003 or after 2019. Also, it should be noticed that publication bias may exist due to the English language restriction applied. Another limitation to be noted is that this review excluded seasonal influenza outbreaks since these occur on a yearly basis. Furthermore, this study was performed before the impact of COVID-19 and hence reflects the published knowledge before the current pandemic. Thus, the results cannot be directly extrapolated to the COVID-19 pandemic.

CONCLUSION

The value of this systematic review of econometric studies is to provide a synthesis of the evidence of the cost of respiratory infectious disease outbreaks and the cost-effectiveness of specific interventions that can be applied in response. Furthermore, our assessment identifies a minimum number of econometric measures which should be recorded during the reporting of respiratory infectious disease outbreaks that would aid future decision making. Our cost analysis results give evidence to public health policymakers, primarily in the EU or the USA, as to the cost-effectiveness of a range of pharmaceutical and non-pharmaceutical intervention strategies which may be applied to mitigate or respond to infectious respiratory disease outbreaks.

Author affiliations

¹School of Medicine, University of Crete, Heraklion, Greece

²Department of Public Health Policy, School of Public Health, University of West Attica, Egaleo, Greece

³Division of Epidemiology and Public Health, University of Nottingham School of Medicine, Nottingham, UK

⁴Epidemiological Methods, European Centre for Disease Prevention and Control, Solna, Sweden

⁵Emergency Preparedness and Response Support, European Centre for Disease Prevention and Control, Solna, Sweden

Acknowledgements We would like to thank Ioanna Lagou and Chrysa Chatzopoulou for contributing to data management.

Contributors CV, JL-B, ST and JES designed the study. KN and KZ undertook the literature review and extracted the data with help from JL-B and RP. JL-B and RP developed the search code. KZ, KN and KA analysed and interpreted the econometrics data. HJ and MC participated in data evaluation and interpretation along with CV, JL-B, RP, JES, KN, KZ, KA and ST. CV and KN wrote the first draft of the manuscript with input from all authors. All authors reviewed and revised subsequent drafts.

Funding This report was commissioned by the European Centre for Disease Prevention and Control (ECDC), to the PREP-EU Consortium, coordinated by Dr Vardavas under specific contract No. 1 ECD.9630 within Framework contract ECDC/2019/001.

Disclaimer The information and views in this manuscript are those of the authors and do not necessarily reflect the official opinion of the Commission/Agency. The Commission/Agency do not guarantee the accuracy of the data included in this study. Neither the Commission/Agency nor any person acting on the Commission's/Agency's behalf may be held responsible for the use which may be made of the information contained therein.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data sharing not applicable as no datasets generated and/or analysed for this study.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Katerina Nikitara <http://orcid.org/0000-0002-7270-6278>

Helen Johnson <http://orcid.org/0000-0002-5730-2638>

Jonathan E Suk <http://orcid.org/0000-0003-4689-4583>

REFERENCES

- European Respiratory Society. *Forum of international respiratory societies the global impact of respiratory disease*, 2017: 1–43.
- European Centre for Disease Prevention and Control (ECDC). *Factsheet about seasonal influenza*, 2017: 1–7.
- Nielsen J, Vestergaard LS, Richter L, *et al*. European all-cause excess and influenza-attributable mortality in the 2017/18 season: should the burden of influenza B be reconsidered? *Clin Microbiol Infect* 2019;25:1266–76.
- WHO. *Definitions: emergencies*. World Health Organization, 2016.
- Meltzer MI, Gambhir M, Atkins CY, *et al*. Standardizing scenarios to assess the need to respond to an influenza pandemic. *Clin Infect Dis* 2015;60:S1–8.
- Mitigation of pandemic influenza. *Review of cost-effectiveness studies*, 2009.
- Drolet M, Bénard Élodie, Jit M, *et al*. Model comparisons of the effectiveness and cost-effectiveness of vaccination: a systematic review of the literature. *Value Health* 2018;21:1250–8.
- Drake TL, Chalabi Z, Coker R. Cost-effectiveness analysis of pandemic influenza preparedness: what's missing? *Bull World Health Organ* 2012;90:940–1.
- Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- Husereau D, Drummond M, Petrou S, *et al*. Consolidated health economic evaluation reporting standards (cheers) statement. *BMJ* 2013;346:f1049.
- Consortium YHE. Perspective - YHEC - York Health Economics Consortium.
- Lu H, Zhao Y, Zhang J, *et al*. Date of origin of the SARS coronavirus strains. *BMC Infect Dis* 2004;4:3.
- Evers S, Goossens M, de Vet H, *et al*. Criteria list for assessment of methodological quality of economic evaluations: consensus on health economic criteria. *Int J Technol Assess Health Care* 2005;21:240–5.
- CCEMG. *EPPI-centre cost converter v.1.4*, 2015.
- IMF. World economic and financial surveys. world economic outlook database.
- Institute JB. The systematic review of economic evaluation evidence. *Implement Sci* 2014:1–24.
- Prager F, Wei D, Rose A. *Total economic consequences of an influenza outbreak in the United States. risk analysis*. Blackwell Publishing Inc, 2017: 4–19.
- Morales-Suárez-Varela M, Llopis-González A, González-Candela F, *et al*. Economic evaluation of health services costs during pandemic influenza A (H1N1) pdm09 infection in pregnant and non-pregnant women in Spain. *Iran J Public Health* 2016;45:423–34.
- Silva ML, Perrier L, Späth H-M, *et al*. Economic burden of seasonal influenza B in France during winter 2010–2011. *BMC Public Health* 2014;14:56.



- 20 Higgins AM, Pettilä V, Harris AH. *The critical care costs of the influenza A/H1N1 2009 pandemic in Australia and New Zealand. Anaesthesia and intensive care*. London: SAGE Publications, 2011: 384–91.
- 21 Wilson N, Nhung N, Higgins A, *et al*. A national estimate of the hospitalisation costs for the influenza (H1N1) pandemic in 2009. *N Z Med J* 2012;125:16–20.
- 22 Rodríguez-Rieiro C, Carrasco-Garrido P, Hernández-Barrera V, *et al*. Pandemic influenza hospitalization in Spain (2009): incidence, in-hospital mortality, comorbidities and costs. *Hum Vaccin Immunother* 2012;8:443–7.
- 23 Lankelma JM, Hermans MHA, Hazenberg E. Implementation of point-of-care testing and a temporary influenza ward in a Dutch Hospital. *Netherland J Med* 2019:109–15.
- 24 Sadique MZ, Adams EJ, Edmunds WJ. Estimating the costs of school closure for mitigating an influenza pandemic. *BMC Public Health* 2008;8:135.
- 25 Orset C. People's perception and cost-effectiveness of home confinement during an influenza pandemic: evidence from the French case. *Eur J Health Econ* 2018;19:1335–50.
- 26 Saunders-Hastings P, Quinn Hayes B, Smith R, *et al*. Modelling community-control strategies to protect hospital resources during an influenza pandemic in Ottawa, Canada. *PLoS One* 2017;12:e0179315.
- 27 Halder N, Kelso JK, Milne GJ. Cost-Effective strategies for mitigating a future influenza pandemic with H1N1 2009 characteristics. *PLoS One* 2011;6:e22087.
- 28 Tracht SM, Del Valle SY, Edwards BK. Economic analysis of the use of facemasks during pandemic (H1N1) 2009. *J Theor Biol* 2012;300:161–72.
- 29 Yarmand H, Ivy JS, Roberts SD. *Cost-effectiveness analysis of vaccination and self-isolation in case of H1N1*, 2010.
- 30 Sander B, Nizam A, Garrison LP, *et al*. Economic evaluation of influenza pandemic mitigation strategies in the United States using a stochastic microsimulation transmission model. *Value Health* 2009;12:226–33.
- 31 Khazeni N, Hutton DW, Garber AM, *et al*. Effectiveness and cost-effectiveness of expanded antiviral prophylaxis and adjuvanted vaccination strategies for an influenza A (H5N1) pandemic. *Ann Intern Med* 2009;151:840–53.
- 32 Balicer RD, Huerta M, Davidovitch N, *et al*. Cost-benefit of stockpiling drugs for influenza pandemic. *Emerg Infect Dis* 2005;11:1280–2.
- 33 Medema JK, Zoellner YF, Ryan J, *et al*. Modeling pandemic preparedness scenarios: health economic implications of enhanced pandemic vaccine supply. *Virus Res* 2004;103:9–15.
- 34 Pérez Velasco R, Praditsitthikorn N, Wichmann K, *et al*. Systematic review of economic evaluations of preparedness strategies and interventions against influenza pandemics. *PLoS One* 2012;7:e30333.

Appendix

Cost-effectiveness of emergency preparedness measures in response to infectious respiratory disease outbreaks: a systematic review and econometric analysis: Supplementary Information

Appendix 1. Search concept construction

OVID MEDLINE

Date of Search	OVID Medline		
	#	Search Terms	Hits
29/7/2019	1	Economics/	27061
	2	"costs and cost analysis"/	47439
	3	Cost allocation/	1997
	4	Cost-benefit analysis/	77184
	5	Cost control/	21373
	6	Cost savings/	11287
	7	Cost of illness/	25397
	8	Cost sharing/	2443
	9	"deductibles and coinsurance"/	1716
	10	Medical savings accounts/	529
	11	Health care costs/	37249
	12	Direct service costs/	1171
	13	Drug costs/	15395
	14	Employer health costs/	1088
	15	Hospital costs/	10427
	16	Health expenditures/	18983
	17	Capital expenditures/	1987
	18	Value of life/	5653
	19	exp economics, hospital/	23708
	20	exp economics, medical/	14108
	21	Economics, nursing/	3989
	22	Economics, pharmaceutical/	2874
	23	exp "fees and charges"/	29802
	24	(low adj cost).mp.	51166
	25	(high adj cost).mp.	13286
	26	(health?care adj cost\$.mp.	10352
	27	(fiscal or funding or financial or finance).tw.	136748
	28	(cost adj estimate\$.mp.	2132
	29	(cost adj variable).mp.	42
	30	(unit adj cost\$.mp.	2368
	31	(economic\$ or pharmacoeconomic\$ or price\$ or pricing).tw.	277405
	32	Economic evaluation.mp.	9144
	33	(Cost?effectiveness analysis or CEA).mp.	21933
	34	(Cost?utility analysis or CUA).mp.	1153
	35	(Cost?benefit analysis or CBA).mp.	26471
	36	(Cost?consequence analysis or CCA).mp.	7687
	37	(Cost?minimi?sation analysis or CMA).mp.	3583
	38	(cost?outcome or marginal analysis).mp.	204
	39	exp Cost benefit analysis/ or exp budgets/	90077
	40	investment\$.mp. or investments/	39609
	41	or/1-40	769608
	42	exp Emergency Preparedness/	2678
	43	exp Preparedness, Emergency/	2678
	44	(Community Preparedness or Community Recovery or Emergency Operations Coordination or (Emergency Public Information and Warning) or Fatality Management or Information Sharing or Mass Care or Medical Countermeasure Dispensing or (Medical Materiel Management and Distribution)	2614

	or Medical Surge or Non-Pharmaceutical Interventions or Public Health Laboratory Testing).mp.	
45	exp Public Health Surveillance/	2623
46	(Epidemiological Investigation or (Responder Safety adj Health) or Volunteer Management).mp.	2867
47	(disaster preparedness or public health emergencies).mp.	2012
48	((Detection adj assessment) or policy development or policy implementation or policy adaptation or health services or (coordination adj communication) or emergency risk communication or personal preparedness).mp.	393599
49	((state or local or national or legal or business or healthcare) and preparedness).mp.	4359
50	(vaccination or immuni?ation or anti?viral medication or personal hygiene or hand hygiene or household ventilation or ((food and safety) or storage) or food hygiene or respiratory etiquette or (washing and saniti?ing) or social distancing or triage or food security or (emergency adj3 food) or (school adj3 closure) or public gathering* or public meeting* or household isolation or quarantine or PPE or personal protective equipment or (environmental adj3 cleaning)).mp.	545475
51	or/42-50	940853
52	exp disease outbreak/ or exp communicable diseases/	119498
53	(disease outbreak or outbreak or epidemic or pandemic or public health emergency).mp.	150101
54	(avian flu or abola or EVD or H1N1 or H5N1 or infectious disease or influenza or swine flu or flu or MERS or Middle East Respiratory Syndrome).mp.	165207
55	(SARS or Severe Acute Respiratory syndrome or measles or zika or cholera or H7N9 or dengue or fever or plague or fever or malaria or polio).mp.	381260
56	(Bacillus cereus or Campylobacter jejuni or Clostridium or Cryptosporidium or Cyclospora cayetanensis or (E adj coli) or Hepatitis A or Listeria monocytogenes or Noroviruses or Salmonella or Shigella or Staphylococcus aureus or Staphylococcus or Vibrio parahaemolyticus or Vibrio vulnificus).mp.	493801
57	(Diphtheria or Haemophilus influenzae type b or Hib or Hepatitis B or Human Papillomavirus or HPV).mp.	165778
58	((Meningococcal adj Infection\$) or Mump\$ or Pertussis or Whooping Cough or Pneumococcal Infection\$ or Polio or Rotavirus or Rubella or German Measles or Tetanus or varicella or chicken pox or vectorborne diseases or vector?borne disease\$ or waterborne diseases or water?borne disease\$ or Cholera or Diarrhea or diarrhoea).mp.	255768
59	(Typhoid fever or Giardiasis or Schistosomiasis or Dracunculiasis or Dysentery or Cryptosporidiosis or amoebiasis or Traveler\$s diarrhea or travelers diarrhoea).mp.	68954
60	exp infectious disease medicine/ or exp malaria/ or exp influenza, human/ or SARS virus/ or exp norovirus/ or exp coronavirus infections/ or exp measles/ or exp poliomyelitis/ or exp chickenpox/	165698
61	(anthrax or botulism or brucellosis or campylobacter enteritis or chikungunya or chlamydia\$ or CJD or Creutzfeldt?Jakob).mp.	63613
62	(diphtheria or echinococcosis or gonococcal or haemophilus influenzae or hepatitis or HIV or AIDS or human immunodeficiency virus or acquired immunodeficiency syndrome).mp.	714394
63	(legionnaires?disease or leptospirosis or listeriosis or lyme or streptococcus pneumoniae or Q fever or rabies or congenital rubella or salmonella or shiga toxin or verocytotoxin?producing E?coli or STEC or VTEC or HUS or haemolitic?uraemic or hemolitic?uremic).mp.	182938
64	(shigellosis or smallpox or syphilis or congenital syphilis or tick?borne viral encephalitis or congenital toxoplasmosis or	341238

		trichinellosis or tuberculosis or TB or typhoid or paratyphoid or VHF or viral hemorrhagic fever\$ or viral haemorrhagic fever\$ or West Nile virus or Yellow fever or (enteritis adj3 yersinia).mp.	
	65	or/52-64	2357602
	66	41 and 51 and 65 (studies before 2003 excluded)	18127

EMBASE

Date of Search	EMBASE		
	#	Search Terms	Hits
29/7/2019	1	Socioeconomics/	133589
	2	Cost benefit analysis/	81690
	3	Cost effectiveness analysis/	143890
	4	Cost of illness/	18428
	5	Cost control/	65812
	6	Economic aspect/	110246
	7	Financial management/	110636
	8	Health care cost/	181209
	9	Health care financing/	13089
	10	Health economics/	32080
	11	Hospital cost/	20343
	12	(fiscal or financial or finance or funding).tw.	178545
	13	Cost minimization analysis/	3375
	14	(cost adj estimate\$.mp.	3181
	15	(cost adj variables\$.mp.	188
	16	(unit adj cost\$.mp.	4210
	17	investment\$.mp. or investments/	49607
	18	or/1-17	906830
	19	"Emergency Preparedness".tw.	1780
	20	(Community Preparedness or Community Recovery or Emergency Operations Coordination or (Emergency Public Information and Warning) or Fatality Management or Information Sharing or Mass Care or Medical Countermeasure Dispensing or (Medical Materiel Management and Distribution) or Medical Surge or Non-Pharmaceutical Interventions or Public Health Laboratory Testing).mp.	3358
	21	exp Public Health Surveillance/	210835
	22	(Epidemiological Investigation or (Responder Safety adj Health) or Volunteer Management).mp.	3638
	23	(disaster preparedness or public health emergencies).mp.	2176
	24	((Detection adj assessment) or policy development or policy implementation or policy adaptation or health services or (coordination adj communication) or emergency risk communication or personal preparedness).mp.	124325
	25	((state or local or national or legal or business or healthcare) and preparedness).mp.	5302
	26	(vaccination or immuni?ation or anti?viral medication or personal hygiene or hand hygiene or household ventilation or ((food and safety) or storage) or food hygiene or respiratory etiquette or (washing and saniti?ing) or social distancing or triage or food security or (emergency adj3 food) or (school adj3 closure) or public gathering* or public meeting* or household isolation or quarantine or PPE or personal protective equipment or (environmental adj3 cleaning)).mp.	703894
	27	or/19-26	1031364


28	exp disease outbreak/ or exp communicable diseases/	119087
29	(disease outbreak or outbreak or epidemic or pandemic or public health emergency).mp.	207717
30	(avian flu or abola or EVD or H1N1 or H5N1 or infectious disease or influenza or swine flu or flu or MERS or Middle East Respiratory Syndrome).mp.	198862
31	(SARS or Severe Acute Respiratory syndrome or measles or zika or cholera or H7N9 or dengue or fever or plague or fever or malaria or polio).mp.	561066
32	(Bacillus cereus or Campylobacter jejuni or Clostridium or Cryptosporidium or Cyclospora cayetanensis or (E adj coli) or Hepatitis A or Listeria monocytogenes or Noroviruses or Salmonella or Shigella or Staphylococcus aureus or Staphylococcus or Vibrio parahaemolyticus or Vibrio vulnificus).mp.	579612
33	(Diphtheria or Haemophilus influenzae type b or Hib or Hepatitis B or Human Papillomavirus or HPV).mp.	241684
34	((Meningococcal adj Infection\$) or Mump\$ or Pertussis or Whooping Cough or Pneumococcal Infection\$ or Polio or Rotavirus or Rubella or German Measles or Tetanus or varicella or chicken pox or vectorborne diseases or vector?borne disease\$ or waterborne diseases or water?borne disease\$ or Cholera or Diarrhea or diarrhoea).mp.	421975
35	(Typhoid fever or Giardiasis or Schistosomiasis or Dracunculiasis or Dysentery or Cryptosporidiosis or amoebiasis or Traveler\$s diarrhea or travelers diarrhoea).mp.	56723
36	exp infectious disease medicine/ or exp malaria/ or exp influenza, human/ or SARS virus/ or exp norovirus/ or exp coronavirus infections/ or exp measles/ or exp poliomyelitis/ or exp chickenpox/	213621
37	(anthrax or botulism or brucellosis or campylobacter enteritis or chikungunya or chlamydia\$ or CJD or Creutzfeldt?Jakob).mp.	78479
38	(diphtheria or echinococcosis or gonococcal or haemophilus influenzae or hepatitis or HIV or AIDS or human immunodeficiency virus or acquired immunodeficiency syndrome).mp.	938033
39	(legionnaires?disease or leptospirosis or listeriosis or lyme or streptococcus pneumoniae or Q fever or rabies or congenital rubella or salmonella or shiga toxin or verocytotoxin?producing E?coli or STEC or VTEC or HUS or haemolitic?uraemic or hemolitic?uremic).mp.	206207
40	(shigellosis or smallpox or syphilis or congenital syphilis or tick?borne viral encephalitis or congenital toxoplasmosis or trichinellosis or tuberculosis or TB or typhoid or paratyphoid or VHF or viral hemorrhagic fever\$ or viral haemorrhagic fever\$ or West Nile virus or Yellow fever or (enteritis adj3 yersinia)).mp.	313309
41	or/28-40	2651637
42	18 and 27 and 41 (studies before 2003 excluded)	14223


ECONLIT


Date of Search	EconLit		
	#	Search Terms	Hits
30/8/2019	1	cost OR (deductibles and coinsurance) OR Medical savings accounts OR health expenditure OR economic OR (fees and charges) OR Economic evaluation OR cost effectiveness analysis OR Cost utility analysis OR cost benefit analysis OR Cost consequence analysis OR Investment	1,344,466
30/8/2019	2	(Emergency Preparedness) OR Preparedness OR emergency OR Surveillance OR disaster OR (detection or diagnosis or identification or early detection) OR screening OR vaccination OR hygiene OR school closure OR quarantine	48,619
30/8/2019	3	disease outbreak OR disease OR infectious diseases OR communicable diseases OR outbreak OR pandemic OR epidemic	9,194
31/8/2019	4	S1 AND S2 AND S3 (limitation: from 2003 to 2019)	965

IDEAS REPEC

Date of Search	IDEAS REPEC			
	Search Term 1	Search Term 2	Search Term 3	Results (n)
28/7/2019	cost-effective	infectious		139
28/7/2020	Emergency Public Information and Warning	cost		8
28/7/2021	Health Surveillance	infectious	cost	10
30/7/2022	economics	health preparedness		39
30/7/2023	cost-effectiveness	cost effectiveness	health preparedness	8
30/7/2024	prevention	cost	disease outbreaks	42
30/7/2025	economic evaluation	Public health surveillance		12
30/7/2026	investment	Infectious disease	outbreak	16
30/7/2027	economics	H1N1		16
30/7/2028	economics	flu	outbreak	25
1/8/2019	Cost-effectiveness	ebola		4
1/8/2019	economics	disease threats		171
1/8/2019	pandemic	economic	cost	42

Appendix 2. Total quality appraisal score (in percentages) for all included studies (n=17)

Appendix 3. Quality appraisal score by item/question for the partial economic evaluation studies (n=6)

Appendix 4. Quality appraisal by item/question of the full evaluation studies (n=11)

Appendix 5. Characteristics of cost analyses studies of influenza outbreaks

Study, (Publication Year)	Setting, year	Study population (n)	Economic Evaluation	Approach	Perspective	Time Horizon	Type of Sensitivity analysis
Prager et al. (2017)	USA, n/a	The population of the USA	Cost of illness (although mentions about cost-effective of vaccination in the pandemic scenario, No CEA outcomes)	Simulation model	Healthcare system, Governmental, Societal	Not clearly stated	Performed, unclear
Morales-Suárez-Varela et al. (2016)	Spain, 2009-2010	Unvaccinated women of childbearing age with influenza A (H1N1)	Partial Economic Evaluation (Cost of illness)	Observational	Healthcare system, Societal	4 months	Not performed
Silva et al. (2014)	France, 2010-2011	Population with Influenza B (201)	Partial Economic Evaluation (cost of illness)	Observational	Payer, Societal	3 months	One-way sensitivity analysis and probabilistic analysis
Higgins et al. (2011)	Australia and New Zealand, 2009	All Influenza cases (H1N1) in New Zealand and Australia (762)	Partial Economic Evaluation Cost of illness	Observational	Healthcare system	3 months	Not performed
Wilson et al. (2009)	New Zealand, 2009	All Influenza hospitalisations in New Zealand) 1224 – 1122 hospitalizations and + 122 ICU	Partial Economic Evaluation, Cost of illness (incl. hypothetical cost-effectiveness analysis)	Observational	Healthcare system	12 months	Probabilistic sensitivity analysis
Rodríguez-Rieiro et al. (2009)	Spain, 2009	All Spanish patients with H1N1 (11,449)	Partial Economic Evaluation (Cost of illness)	Observational	Healthcare system	12 months	Not performed

Appendix 6. Characteristics of studies on preparedness and response measures of influenza outbreaks

Study, (Publication year)	Setting, Year	Population (n)	Economic Evaluation Approach	Perspective	Timeframe	Discount	Sensitivity analysis
Observational studies							
Lankelma et al. (2019)	Netherlands, 2017-2018	Patients with acute RTI at the emergency department (1546 tests, 624 cases)	Partial Economic Evaluation	Healthcare system	4,5 months	N/A	Not performed
Sadique et al. (2008)	UK, 2005	Working parents with depending children	Partial Economic Evaluation	Societal	1 year	N/A	Scenarios
Simulation or mathematical models							
Orset (2018)	France, 2014	200 participants, data extrapolated	Both cost-benefit and cost-effectiveness analysis	Public health and societal	1 year	1% for costs	Not performed
Saunders-Hastings et al. (2017)	Canada, n/a	A simulation of Ottawa, Canada (1.2 million)	cost-effectiveness analysis	Healthcare system	Lifetime	1.5%	Multivariate sensitivity analyses
Halder et al. (2011)	Australia, 2009	A community in Western Australia (30,000)	cost-effectiveness analysis	Healthcare and Societal	Lifetime	3%	Scenarios
Tracht et al. (2012)	USA, (2009-2010 influenza season)	Simulation of the US (302 million people:73 million children, 191 million adults, and 38 million seniors)	cost-effectiveness analysis	Healthcare system and societal	1 year	N/A	Multivariate sensitivity analyses
Yarmand et al. (2010)	USA, (2009-2010 influenza season)	North Carolina State University undergraduate students (23,087)	cost-effectiveness analysis	Healthcare system	5 months	N/A	One-way and two-way sensitivity analyses
Sander et al. (2009)	USA, n/a	Residents of a 1 632-million-person city	Cost Utility Analysis	Societal	6 months	3%	Multivariate sensitivity analyses
Khazeni et al. (2009)	USA, n/a	A U.S. metropolitan city (8.3 million)	cost-effectiveness analysis	Societal	Lifetime	3% for benefits/ costs	Monte Carlo probabilistic sensitivity analysis
Balicer et al. (2005)	Israel, n/a	Population of Israel (1 618 200 cases)	Cost-benefit analysis	Healthcare system Societal	Lifetime	Not specified	Multivariate sensitivity analyses
Medema et al. (2004)	UK, Germany, Netherlands, (2004)	Developed Countries (1 Billion people)	Cost-effectiveness analysis	Healthcare system	Not clearly stated	5%	Performed unclear

RTI: Acute respiratory tract infection, N/A: Not applicable, UK: United Kingdom, US: United States

Appendix 7. Comparative analysis of health indexes when averting/responding to respiratory disease outbreaks

Study / year	Intervention(s)/Screening methods vs. comparators	Outcomes/benefits
Prager et al. 2016	Case 1: No Vaccination, Seasonal Outbreak Case 2: No Vaccination, Pandemic Outbreak Vs. Case 3: Vaccination, Seasonal Outbreak Case 4: Vaccination, Pandemic Outbreak	<p>Productivity loss and behavioural response In the case of a pandemic influenza outbreak Vaccination:</p> <ol style="list-style-type: none"> 1. Reduces illness-related workday losses from 83.3 million days to 61.1 million days (a reduction of 22.2 million days). 2. Causes 7.4 million days of workday losses due to the time that people spend on getting the vaccination doses. 3. Can reduce public avoidance behaviours by 25%. <p>In the case of a seasonal influenza outbreak Vaccination:</p> <ol style="list-style-type: none"> 1. Reduces illness-related workday losses from 18.7 million days to 13.9 million days (a reduction of 4.8 million days). 2. Causes 6.7 million days of workday losses due to the time that people spend obtaining vaccinations. 3. Can reduce public avoidance behaviours by 25%.
Medema et al. 2004	<ol style="list-style-type: none"> 1. Egg-based vaccine manufacture 2. Cell culture-based vaccine manufacture <p>Vs.</p> <p>No intervention</p>	<p>Cases, PCP consultations and hospitalizations prevented Cell culture-based intervention vs no intervention: Cell culture-based intervention avoids 75 million influenza cases, 3.78 million PCP consultations for influenza treatment and, respectively, 5.81 million and 1.21 million influenza-related hospitalizations and excess deaths.</p> <p>Egg-based vaccine intervention vs no intervention: Egg-based vaccine intervention leads to vaccination of 17% of the population, which avoids 29.8 million influenza cases, 1.74 million PCP visits, 2.67 million hospitalizations and 556 000 deaths</p> <p>Cell culture-based intervention vs egg-based vaccine intervention with 17% vaccine coverage: Cell culture-based intervention strategy leads to vaccination of 37% of the population, avoiding an additional 35 million influenza cases, 2.04 million PCP consultations for influenza treatment, 3.14 million influenza-related hospitalizations and 654 500 excess deaths</p> <p>Years of life lost (YLL) Cell culture-based intervention strategy: 2.56 million</p>

<p>Saunders-Hastings et al. 2017</p>	<p>1. Vaccination and antiviral treatment 2. Vaccination, antiviral treatment and antiviral prophylaxis 3. Community-contact reduction, personal protective measures and voluntary isolation 4. Community-contact reduction, personal protective measures, voluntary isolation and antiviral treatment 5. School closure, community-contact reduction, personal protective measures, voluntary isolation and quarantine 6. All interventions</p> <p>Vs.</p> <p>7. No intervention</p>	<p>Hospitalizations In case of no intervention, a total of 2 472 pandemic-associated hospitalizations have been estimated. Following no intervention, vaccination interventions (combined with other interventions) contributed to 765-815 hospitalizations. Last, school closure, combined with other interventions, contributed to 108-550 hospitalizations.</p> <p>YLL</p> <ol style="list-style-type: none"> 1. 3,026 2. 2,801 3. 1,767 4. 1,607 5. 1,393 6. 267 7. 9,421 <p>Reductions of illness (H2N2 cases) Vaccination, personal protective measures, combined voluntary isolation and quarantine procedures resulted in the greatest reductions, producing attack rates of 50.0%, 45.5% and 33.9%, respectively. Antiviral treatment, antiviral prophylaxis, school closure and community-contact reduction produced only small reductions in illness attack rate, whether implemented alone or in combination with other interventions. Even in the absence of any pharmaceutical intervention, adherence to rigorous non-pharmaceutical protocols -school closure, community-contact reduction, personal protective measures, voluntary isolation and quarantine-resulted in a reduction of the illness attack rate to 15.2%.</p>
<p>Khazeni et al. 2009</p>	<p>1) Vaccination and antiviral pharmacotherapy in quantities similar to those currently available in the U.S. stockpile (stockpiled strategy), 2) Stockpiled strategy but with the expanded distribution of antiviral agents (expanded prophylaxis strategy), and 3) Stockpiled strategy but with the adjuvanted vaccine (expanded vaccination strategy).</p> <p>Vs.</p> <p>no intervention</p>	<p>Clinical attack rate The clinical attack rate has been 11%, 17%, 19% and 33% for expanded adjuvanted vaccination, expanded antiviral prophylaxis, Stockpiled strategy and for no intervention, respectively.</p> <p>Deaths averted Expanded adjuvanted vaccination – 45 941 deaths averted Expanded antiviral prophylaxis – 32 745 deaths averted Stockpiled strategy – 29 761 deaths averted No intervention - No deaths averted</p>
<p>Sander et al.</p>	<ol style="list-style-type: none"> 1. HTAP25 with a stockpile for 25% of the population 2. HTAP50 with a stockpile for 50% of the population 3. HTAP with an unlimited stockpile 	<p>QALYs gained, total Expanded adjuvanted vaccination – 404 030 total QALYs gained Expanded antiviral prophylaxis – 282 329 total QALYs gained</p>

<p>2009</p>	<p>4. School closure for 26 weeks 5. Pre vaccination 70% of the population with a low efficacy vaccine 6. HTAP25 + school closure: 7. HTAP50 + school closure: 8. HTAP + school closure: 9. Pre vaccination + school closure: Pre vaccinating 70% population with the low-efficacy vaccine, plus closing all schools for 26 weeks 10. Treatment only: Treating all cases with antivirals 11. FTAP25 for household contacts and 60% of work/school contacts, stockpile for 25% of the population 11. FTAP50 for household contacts and 60% of work/school contacts, stockpile for 50% of population 12. FTAP for household contacts and 60% of work/school contacts, stockpile unlimited 14. FTAP25 + school closure 13. 15. FTAP50 + school closure 14. 16. FTAP + school closure</p>	<p>Stockpiled strategy – 258 342 total QALYs gained No intervention - No QALYs gained</p> <p>QALYs per 1000 population, total All interventions gained a similar amount of QALYs, with some differences between them (21,141 for no intervention to 21 403 for pre vaccination and school closure). Compared to FTAP not involving school closure, FTAP plus school closure or pre vaccination plus school closure gains 51 QALYs</p> <p>QALYs per 1000 population, incremental FTAP and school closure and the intervention of pre vaccination and school closure contributed to the most incremental QALYs (262)</p> <p>Deaths per 1000 population Pre-vaccination intervention was the most effective strategy. Only 1 death/1000 population occurred via this strategy. On the other side, most deaths have been seen in case of no intervention (13 deaths/1000 population) and FTAP25 with 12 deaths.</p> <p>Number of cases Full TAP is the most effective single strategy, reducing the number of cases by 54% Pre-vaccination reduces the number of cases by 48% Adding school closure to full TAP or pre-vaccination further improves health outcomes</p>
<p>Yarm and et al. 2010</p>	<p>Self-isolation and mandatory quarantine Vs. vaccination</p>	<p>Effectiveness in low-levels of interventions Vaccination is more effective than self-isolation.</p> <p>Effectiveness in high-levels of interventions Self-isolation is more effective than vaccination. This has been shown due to weaknesses of vaccinations, such as delays in effectiveness.</p>
<p>Halder et al. 2011</p>	<p>Antiviral drugs combined with limited duration school closure Vs. 1. School closure as a sole intervention alone and as dual, triple, quadruple strategy 2. Other social distancing strategies, such as reduced workplace attendance</p>	<p>The illness attack rate of interventions (symptomatic) The illness attack rate ranges from 2.4% (SD 0.37) to 8.5% (SD 1.1) while that of the unmitigated attack rate is 13% (SD 0.9). The individual school closure for 2 weeks along with the continuous – 50% workplace closure, antiviral treatment, household antiviral prophylaxis and extended antiviral prophylaxis showed the lowest illness attack rate (2.4%). This combination is the most effective intervention. Short-duration school closure is less effective (6.5 to 8.2 illness attack rate) Continuous school closure is more effective, with an attack rate of 3.2.</p>

<p>Orset 2018</p>	<p>Home confinement</p> <p>Vs.</p> <p>No intervention</p>	<p>Incidence rate reduction by the home confinement intervention</p> <p>There are studies that indicate the higher the compliance rate regarding home confinement, the higher the reduction of the incidence rate of influenza will be. More particularly:</p> <p>In case of a 70% compliance rate: 83% reduction of incidence rate In the case of 80% compliance rate: 91% reduction of incidence rate</p> <p>The compliance rate with home confinement is between 75.90 and 94.44%, for this study.</p> <p>Rate reduction threshold in the incidence due to intervention</p> <p>The higher the proportion of all cases complying with home confinement, the higher the reduction rate of the threshold for VSL will be.</p> <p>For example: In case of 49.24% of all cases complying with home confinement: €7.65 million Threshold for VSL In case of 51.39% of all cases complying with home confinement: €5.06 million Threshold for VSL</p>
<p>Tracht et al. 2012</p>	<p>Mask wearing group</p> <p>Vs.</p> <p>No intervention</p>	<p>When there are no interventions (no masks worn)</p> <p>Cumulative number of cases/ based on three scenarios - R avg/unc*</p> <p>In the case of 1.25; A total of 101,424,384 cases. Most of them identified at 18-64 age group.</p> <p>In the case of 1.3; A total of 117 673 024 cases. Most of them identified at 18-64 age group.</p> <p>In the case of 1.35; A total of 130 043 351 cases. Most of them identified at 18-64 age group.</p> <p>Hospitalizations</p> <p>Based on three different scenarios - R avg/unc: 1.25, 1.3, and 1.35</p> <p>In the case of 1.25: For all age groups, a total of 3 275 616 hospitalizations have been estimated. 75.8% of them found to be in 18-64 ages</p> <p>In the case of 1.3: For all age groups, a total of 3 793 350 hospitalizations have been estimated. 74.8% of them found to be in 18-64 ages</p> <p>In the case of 1.35: For all age groups, a total of 4 184 352 hospitalizations have been estimated. 73.7% of them found to be in 18-64 ages</p> <p>Deaths</p> <p>More deaths have been found in ages 18-64, both in three scenarios, and more than 90% of the total deaths (281 319-349 578)</p>

		As a result, the model showed that in case of 10% of the population wearing masks with an effectiveness of 20% in reducing susceptibility and infectivity, there is a large reduction in the cumulative number of cases.
--	--	--

PCP: Primary care physician, YLL: Years of life lost, VSL: Value of statistical life, QALY: Quality-adjusted life year, FTAP: Full-targeted antiviral prophylaxis, SD: Standard deviation,

* Average effective reproduction number (uncontrolled)