



apjtm.org



Perspective

Asian Pacific Journal of Tropical Medicine

doi: 10.4103/apjtm.apjtm_100_24

Addressing schistosomiasis in the Philippines: Need for a coordinated intersectoral effort

Alvin I. Abeleda¹✉, Don Eliseo Lucero-Prisno III^{2,3}, Jerico B. Ogaya⁴¹College of Public Health, University of the Philippines Manila, Philippines²Department of Global Health and Development, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London, United Kingdom³Faculty of Management and Development Studies, University of the Philippines (Open University), Los Baños, Laguna, Philippines⁴Department of Medical Technology, Institute of Health Sciences and Nursing, Far Eastern University, Manila, Philippines

The World Health Organization (WHO) recognized schistosomiasis as one of the neglected tropical diseases (NTDs) caused by parasitic blood flukes. With a 4.68% national prevalence in 2017, schistosomiasis is endemic in 28 provinces in the Philippines[1,2]. Having agriculture as a common source of livelihood, the country is vulnerable to schistosomiasis as this disease is transmitted when a larva, released by freshwater snails, penetrates the skin of a susceptible individual. In fact, it was estimated that 12.4 million Filipinos are at risk, while 2.7 million are directly exposed to the parasite[2]. This remains to be a major public health concern as it contributes to a considerable morbidity and mortality. In children, this disease causes malnutrition and impaired cognitive development. Globally, schistosomiasis as an NTD puts the burden on poverty-stricken areas, especially in low- and middle-income countries, exacerbating the health inequality within a country and across the globe.

To address this, the Philippines implements the National Schistosomiasis Control and Elimination Program, highlighting mass drug administration and preventive chemotherapy as the main strategies for managing morbidity. While this program is a positive step, the government needs to make additional efforts to effectively control schistosomiasis.

In particular, improvement of surveillance of schistosomiasis must be prioritized. The most common diagnostic tests done for schistosomiasis in the country are direct fecal smear and Kato-Katz Technique, both of which have low sensitivities[3]. This leads to underreporting of schistosomiasis cases, thus not seeing the true magnitude of the disease. It manifests the dire need for diagnostic advancements, shifting from microscopic examinations to serologic

tests such as the detection of circulating anodic and cathodic antigen (CCA/CAA) and polymerase chain reaction (PCR), which, although more expensive, shows a significantly higher sensitivity and specificity[4,5]. By investing in making these serologic tests available and accessible in endemic areas and capacitating healthcare workers in doing such tests with higher precision, the true prevalence of schistosomiasis in the Philippines will be determined and appropriate measures will be recommended accordingly.

Aside from this, a strong collaboration with the water, sanitation, and hygiene (WASH) sector must be reinvigorated since open defecation contributes to the transmission of schistosomiasis[4]. In fact, infected individuals eliminate *Schistosoma* eggs with feces or urine[6]. To reduce its transmission, it is necessary to achieve zero open defecation in the country, especially in the endemic areas.

Along with the WASH sector, the collaboration with the Bureau of Animal Industry of the Department of Agriculture, together with other veterinary sectors, must be solidified[4]. In particular, *Oncomelania quadrasi*, a freshwater snail, serves as the main intermediate host for the growth and development of the parasite[5]. Thus, appropriate interventions, such as the use of molluscicides or

✉To whom correspondence may be addressed. E-mail: a.beleda32@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2024 Asian Pacific Journal of Tropical Medicine Produced by Wolters Kluwer-Medknow.

How to cite this article: Abeleda AI, Lucero-Prisno DE III, Ogaya JB. Addressing schistosomiasis in the Philippines: Need for a coordinated intersectoral effort. Asian Pac J Trop Med 2024; 17(11): 479–480.

Article history: Received 10 February 2024
Accepted 22 September 2024

Revision 20 September 2024
Available online 14 October 2024

the proliferation of competitor snails as biological controls, must be made to control the population of these intermediate hosts.

Overall, despite the implementation of the National Schistosomiasis Control and Elimination Program, the Philippines should show more commitment to do better. There shall be a coordinated intersectoral effort, including the surveillance and epidemiology, WASH, and veterinary sector, for schistosomiasis control.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Funding

The authors received no extramural funding for the study.

Authors' contributions

A.A. and L-P.D. conceived the research. A.A. led the literature review and drafted the initial version of the manuscript. L-P.D. and J.O. provided critical revision of the manuscript. A.A., L-P.D., and J.O. contributed to the final version of the manuscript.

References

- [1] Gong YF, Feng JX, Luo ZW, Xue JB, Guo ZY, Zhang LJ, et al. Spatiotemporal heterogeneity of schistosomiasis in mainland China:

Evidence from a multi-stage continuous downscaling sentinel monitoring.

Asian Pac J Trop Med 2022; **15**(1): 26-34.

- [2] Tenorio JC, Molina E. *Schistosoma japonicum* in the Philippines: Its epidemiology, diagnostics, control, and elimination. *J Agric Res Dev Ext Technol* 2021; **3**(1): 71-87.
- [3] Belizario Jr. V, Plan A, de Leon W. Laboratory diagnosis of selected neglected parasitic diseases in the Philippines: Can we do better? *Acta Med Philipp* 2014; **48**(4): 4-10.
- [4] Belizario Jr. V, de Cadiz A, Navarro R, Flores MJ, Molina V, Dalisay SN, et al. The status of schistosomiasis japonica control in the Philippines: The need for an integrated approach to address a multidimensional problem. *Int J One Health* 2022; **8**(1): 8-19. <https://doi.org/10.14202/IJOH.2022.8-19>.
- [5] Hoekstra P, Madinga J, Lutumba P, van Grootveld R, Brienen E, Corstjens P, et al. Diagnosis of schistosomiasis without a microscope: Evaluating circulating Antigen (CCA, CAA) and DNA detection methods on banked samples of a community-based survey from DR Congo. *Trop Med Infect Dis* 2022; **7**(10): 315. <https://doi.org/10.3390/tropicalmed7100315>.
- [6] Nelwan M. Schistosomiasis: Life cycle, diagnosis, and control. *Curr Ther Res Clin Exp* 2019; **91**: 5-9.

Publisher's note

The Publisher of the *Journal* remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Edited by Zhang Q, Lei Y, Pan Y