

OPEN

Youth-friendly sexual health services and peer support for improved sexual and reproductive health outcomes among adolescents and young adults in South Africa: results of a factorial randomized controlled trial

Jana Jarolimova, MD¹, Jacob Busang, MSc², Natsayi Chimbindi, PhD^{2,3,4}, Nonhlanhla Okesola, BSN², Theresa Smit, PhD², Guy Harling, ScD^{2,3,4,5}, Nuala McGrath, ScD^{2,4,6}, Andrew Copas, PhD³, Janet Seeley, PhD^{2,4,7}, Kathy Baisley, MSc^{2,7}, Maryam Shahmanesh, PhD^{2,3,4*}, and the Isisekelo Research Group (Carina Herbst, MSc², Jaco Dreyer, NDipIT², Thembelihle Zuma, PhD^{2,3,4}, Thandeka Khoza, MBChB², Ngundu Behuhuma, MBChB², Ingrid V. Bassett, MD^{1,2}, Lorraine Sherr, PhD³)

¹Massachusetts General Hospital, Boston, USA

²Africa Health Research Institute, Durban, South Africa

³University College London, London, United Kingdom

⁴University of KwaZulu-Natal, Durban, South Africa

⁵University of the Witwatersrand, Johannesburg, South Africa

⁶University of Southampton, Southampton, United Kingdom

⁷London School of Hygiene & Tropical Medicine, London, United Kingdom

*Corresponding author:

Maryam Shahmanesh, MD PhD, UCL, Institute for Global Health, 3rd floor Mortimer Market Centre, Capper Street, London WC1E 6JB, UK. Tel: +44 7776185572; e-mail: m.shahmanesh@ucl.ac.uk

Funding:

This trial was made possible through funding from the US National Institutes of Health (R01MH114560); Wellcome Trust (082384/Z/07/Z); 3ie international initiative for impact evaluation, and the Bill and Melinda Gates Foundation (INV-033650). MS is a National Institute for Health and Care Research Professor (NIHR 301634). GH is supported by a fellowship from the Wellcome Trust and Royal Society (210479/Z/18/Z). NMcG is supported by an NIHR Research Professorship award (Ref:RP:2017-08-ST2-008) using UK aid from the UK Government to support global health research. JJ is supported by NIH K23AI174917. IVB is supported by NIH K24AI141036. The views expressed in this publication are those of the authors and not necessarily those of the NIHR, the UK government, or the US National Institutes of Health.

Declaration of interests

JJ has received in-kind research support from binx health. All other authors declare no competing interests.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Short summary

Peer support and integrated SRH each had only small effects on STIs, contraceptive use, and pregnancy among young people in rural South Africa; combined or more intensive interventions are needed.

ACCEPTED

Abstract

Background

Adolescents and young adults in South Africa have high burdens of STIs and unintended pregnancy. We evaluated the impact of peer support and/or expanded sexual and reproductive health (SRH) services on STIs, contraception, and pregnancy in rural KwaZulu-Natal, South Africa.

Methods

We analyzed secondary outcomes from a 2x2 factorial randomized controlled trial conducted March 2020-August 2022 among 16-29-year-olds, comparing: 1) enhanced Standard of Care (SoC): access to mobile youth-friendly HIV prevention (AYFS); 2) SRH: self-collected STI testing and referral to AYFS with expanded SRH services; 3) Peer-support: peer navigator facilitation of AYFS attendance; 4) SRH+peer-support. At 12 months all participants were offered STI testing; female participants self-reported contraceptive use and pregnancy.

Results

Among 1743 trial participants (51% female), 927 (53%) had 12-month STI results; 209 (22.5%) tested positive: 163 (17.6%) chlamydia, 54 (5.8%) gonorrhea, 44 (4.8%) trichomoniasis. STI prevalence was somewhat lower among those exposed to peer-support (aOR adjusted for sex, age, location: 0.77, 95%CI 0.56-1.06) or SRH (aOR 0.74, 0.56-1.06) and, compared to SoC, was reduced in those exposed to both (aOR 0.59, 0.38-0.94). In SRH arms, 64/469 (13.6%) had a new STI at 12 months, with no difference by peer-support ($p=0.97$). Among females, 336/634 (53.0%) reported using contraception and 47/667 (7.1%) pregnancy, with little difference by study arm.

Conclusions

Peer support and STI testing with expanded SRH each had no more than small effects on STIs, contraception, or pregnancy. Combined or more intensive interventions, e.g., repeat screening, enhanced partner notification, and deeper understanding of structural drivers, are needed.

ACCEPTED

Introduction

Adolescents and young adults (AYA) in South Africa experience a high burden of poor sexual and reproductive health (SRH) outcomes. Over 40% of the one million new daily global infections with gonorrhea, chlamydia, trichomoniasis, and syphilis occur in sub-Saharan Africa.¹ In South Africa specifically, sexually transmitted infection (STI) prevalence rates are estimated to be as high as 14.7% among adult women and 6.0% among men for chlamydia, and 6.6% among women and 3.5% among men for gonorrhea,² and are highest in youth aged 15-24.^{3,4} In South Africa, STIs are syndemic with HIV, increasing risks for HIV transmission and acquisition when untreated and sharing common risk factors.^{5,6} In the province of KwaZulu-Natal, population-based studies have shown curable STI prevalence as high as 20% among young women and 10% among young men.^{3,4} This same age group also experiences a high rate of unintended and teenage pregnancies and unmet contraceptive need.⁷

Untreated curable STIs can lead to substantial morbidity, particularly for women, including pelvic inflammatory disease, tubal infertility, and pregnancy complications.⁸⁻¹¹ However, multiple barriers limit young people's attainment of optimal SRH outcomes. STI diagnosis and management is limited by a lack of accessible and affordable diagnostic testing, leading to a reliance on syndromic management and absence of asymptomatic screening for high-risk populations.¹² This results in the undertreatment of STIs, the majority of which are asymptomatic, particularly in women.^{13,14} Access to comprehensive SRH care in clinics and other healthcare facilities for young adults is limited by stigma, lack of privacy, judgmental approaches from healthcare workers, and logistical barriers.^{15,16} The World Health Organization, in the 2022-2030 Global Health Sector Strategic Plan on HIV, Viral Hepatitis, and Sexually Transmitted Infections,

thus recommends both an expansion in diagnostic testing and screening for STIs and the development of age-appropriate and positive sexual health education and services.¹⁷

There is growing evidence on the effectiveness of integrated, community-based SRH services. A population-based study in KwaZulu-Natal found that home-based STI specimen collection was highly acceptable among AYA,⁴ and mobile-clinic based STI testing has found high rates of untreated STIs in rural South Africa.¹⁸ Community-based STI testing could both decrease the prevalence and complications of curable STIs and create demand for further SRH services. In some settings, STI diagnosis has been associated with subsequent uptake of HIV pre-exposure prophylaxis (PrEP).¹⁹ Further, there is growing evidence for peer-led interventions to support HIV prevention²⁰, and peer navigation and support could similarly enable linkages to comprehensive SRH care including STI testing. We hypothesized that integrated SRH services including home-based self-sampling for STI testing and referral to youth-friendly clinics, along with peer support, could improve SRH outcomes in rural South Africa, including STI positivity, pregnancy, and contraceptive use at 12 months.

Methods

Study design

This secondary outcome analysis reports results from a 2x2 factorial randomized controlled trial (Isisekelo Sempilo, NCT 04532307) evaluating acceptability, feasibility, and preliminary population-level impact of peer navigator support, with or without comprehensive SRH services, on the prevalence of transmissible HIV. The four-arm trial enrolled AYA ages 16-29 in

uMkhanyakude district in rural KwaZulu-Natal, South Africa. Full trial details are described in the protocol paper.²¹

Study setting

The study was conducted within a Health and Demographic Surveillance Site (HDSS), where the Africa Health Research Institute (AHRI) has been conducting annual household-based surveys since 2000. The HDSS covers 845km² with ~140,000 individuals in 20,000 households, including over 20,000 AYA aged 16-29.²² The mostly rural study area has high unemployment (62% of adults without formal employment) and HIV prevalence of 19% among men and 40% among women aged 15-54 years.²²

Study procedures

Using the HDSS as a sampling frame, 3000 AYA, stratified by sex and area, were randomly selected to be assessed for eligibility. Men and women aged 16-29 years, residing in the HDSS area, willing and able to provide informed consent, and willing to be contacted at 12 months, were eligible. At enrollment, participants were randomized between four study arms (**Figure S1**): a) enhanced standard of care (SoC) (referral to adolescent and youth friendly services [AYFS] comprised of clinic-based, nurse-led HIV-focused services), b) SRH (home-based self-collection of STI specimens with referral to AYFS for STI results management with integrated, expanded SRH services), c) peer support (referral to peer navigator to assess social, health, and educational needs and provide risk-informed HIV prevention and referral to AYFS),²³ or d) combined SRH and peer support. Participants randomized to the two SRH arms were offered home-based STI self-sampling at enrollment, as well as expanded counseling and service provision related to sexual

health, fertility, and family planning, provided through the AYFS, beyond that offered in the SoC.²¹ For the STI testing, participants were instructed to self-collect vaginal swab or urine specimens during a study staff home visit, and were given an invitation card to attend the AYFS after 7 days to receive SRH services and the test results. Participants with positive STI results who did not present to the AYFS clinics within 7 days were contacted individually for treatment. In the peer support arms, participants were offered support of named peer navigators residing in their area, trained to provide one-to-one HIV prevention counseling, health promotion, and support in accessing clinical services. Peer support activities, which normally occur in person, were temporarily transitioned to virtual support March 24, 2020 – November 24, 2020 due to the COVID-19 pandemic.

AYFS, the enhanced SoC upon which the other interventions were layered, were gender-neutral and HIV status-neutral HIV prevention and treatment services and basic primary care SRH services delivered by study nurses in two primary health clinics and two mobile clinics that visited fixed sites across the surveillance area every 2 weeks. For participants in the two SRH arms, expanded SRH services to be provided by the AYFS were emphasized at time of referral. All AYFS clinic attendees were offered HIV counselling and testing, with immediate ART initiation if found to be living with HIV, or PrEP if negative and eligible (i.e., HIV negative on rapid antibody testing, no symptoms of acute HIV, willing and able to take PrEP as prescribed, no contraindications to oral PrEP, $\geq 35\text{kg}^{24}$). Further, all clinic attendees were offered counseling around U=U (undetectable=untransmittable); pregnancy testing and basic family planning support; syphilis, hepatitis B testing and vaccines; and STI syndromic management. Participants were encouraged to attend the services three-monthly for HIV testing, PrEP and contraception refills,

and SRH services. Attendance at AYFS services was recorded for all enrolled participants; initial visits were tracked with a barcoded referral slip.

All participants were contacted 12 months after enrollment. Participants completed a questionnaire that included uptake and experience of HIV services through the AYFS, use of contraception, and pregnancy incidence (among females), and were offered STI testing by home-based self-collection. For female participants, research staff described the procedure to self-collect a vaginal swab. Menstruating females provided urine specimens. Male participants were instructed to collect a first-catch urine specimen. All STI specimens collected either at enrollment (in the two SRH arms) or during the 12-month survey were transported to the AHRI central laboratory in Durban, where testing for *Neisseria gonorrhoeae*, *Chlamydia trachomatis*, and *Trichomonas vaginalis* was conducted by real-time polymerase chain reaction using GeneXpert (Cepheid, Sunnyvale, CA, USA). Participants with positive results were contacted for treatment according to South African national guidelines and counseled on partner notification.²⁵

Measures

Valid STI test results were recorded as ‘detected’ or ‘not detected’; invalid as ‘invalid’ or ‘error’ based on test output. New STI was defined as testing positive for one of the three STIs at 12 months after testing negative for the same STI at baseline. Use of contraception at 12 months was assessed by the question, “are you currently using any contraceptive methods to prevent pregnancy?”. New pregnancy was assessed by the question, “are you currently pregnant?”. Socio-demographic data, including education (whether still in school, years completed) and current employment (employed,

not employed, studying), were obtained from the 12-month survey. Residence (urban/peri-urban, rural) was derived from linking study participants to the HDSS household-level survey.

Statistical analysis

We summarized demographic data using medians and interquartile ranges (IQR) for continuous variables and frequency counts and percentages for categorical variables. We fitted logistic regression models to jointly estimate the odds ratio (OR) and 95% confidence interval (CI) for the main effects of peer navigator support and the SRH intervention on STI prevalence, contraceptive use, and pregnancy at 12 months, assuming no interaction, in an intention-to-treat analysis. As a secondary approach we also fitted a 4-level categorical variable to estimate the effect of each trial arm, i.e., peer support alone, SRH alone, and peer support combined with SRH, all relative to SoC. We additionally conducted a per-protocol adjusted main effects analysis limited to those participants who attended at least one AYFS clinic visit. We tested for an interaction between peer-support and SRH interventions on 12-month STI positivity. We compared new STIs at 12 months between the SRH arms using a Chi-square test. Characteristics of participants who were reached for the endline survey and consented to STI testing were compared to all enrolled participants using univariate and multivariable logistic regression to assess for any differences. Missing data were not imputed and participants with missing data were not included in multivariable analyses. All reported p-values were two-tailed; $p < 0.05$ was considered statistically significant. Analyses were conducted using Stata version 16.1 (Stata Corp, College Station, TX, USA).

Role of the funding source

The funder had no role in study design, data collection, data analysis, data interpretation, or writing of the report. All authors had full access to all study data and accept responsibility for the decision to submit for publication.

Results

Between 02 March 2020-18 May 2021, 2627 (88%) of the 3000 AYA sampled were contacted, among whom 2301 (88%) were eligible and 1743 (76%) consented to enrollment (**Figure 1**). Primary outcomes of the trial, which showed that the SRH intervention increased linkage to AYFS but neither intervention reduced transmissible HIV, have been previously reported.²⁶ There were no important differences in baseline characteristics by arm. Within 60 days of enrollment, 755 participants (43%) linked to clinical services and attended at least one AYFS visit; significantly higher in the SRH study arms.²⁶ Overall, 519 (29.8%) of 1743 participants attended AYFS more than once; this was highest in the combined SRH and peer-support arm (34%).^{26,27}

Among the 1743 enrolled participants, 1300 (75%) were reached at 12 months for the endline survey, of whom 987 (76%) consented to STI testing. In adjusted analyses, those reached at 12 months and consented to STI testing were more likely to have been randomized to one of the SRH arms, to be female, to reside in a rural area, and to be unemployed (**Table S1**). In total, 927/987 (94%) participants provided adequate STI specimens and received valid test results and thus contributed data to analyses of 12-month STI outcomes. Contraceptive use and pregnancy data in the 12-month survey were available for 634/687 (92%) and 667/687 (98%) female participants, respectively.

Among the 927 participants with complete 12-month STI results, 209 (22.5%) tested positive for at least one STI: 163 (17.6%) chlamydia, 54 (5.8%) gonorrhea, and 44 (4.8%) trichomoniasis (**Table 1**). STI positivity by sex and age is shown in **Figure 2**. In the primary main effects analysis, adjusting for sex, age, and location, 12-month STI positivity was somewhat lower in the SRH intervention arms compared to non-SRH arms (20.4% vs 24.9%, aOR 0.74, 95%CI 0.56-1.06) and in peer-support arms (20.7% vs 24.5%, aOR 0.77, 95%CI 0.56-1.06, **Table 1, Figure 3**). There was little evidence of interaction between peer-support and SRH ($p=0.978$). There was not clear variability in STI positivity between the four trial arms in a ‘global’ test ($p=0.16$) after adjustment. However, when compared to SoC, there was some evidence of lower STI positivity, particularly in the combined SRH and peer-support arm (aOR 0.59, 95%CI 0.38-0.94, **Table 1, Figure 3**). In a per-protocol adjusted main effects analysis limited to participants who attended at least one AYFS visit ($n=856$), 12-month STI positivity was lower among those assigned to SRH intervention arms (20.4% vs 27.6%; aOR 0.69, 95%CI 0.50-0.96, $p=0.026$) and those assigned to peer-support arms (20.9% vs 26.7%; aOR 0.70, 95%CI 0.50-0.97, $p=0.030$) (**Table S2**).

Among the 469 participants with both baseline and endline STI test results, 64 (13.6%) acquired a new STI during the 12-month study period; this rate did not differ by assignment to the peer-support intervention (**Table 2**). Thirty-nine participants (8.3%) tested positive for the same STI at baseline and endline, demonstrating either persistent infection or re-infection; this rate also did not differ by assignment to peer-support (**Table 2, S3**).

At study end, 336/634 (53.0%) female participants reported using any method of contraception, with no difference by SRH ($p=0.301$) or peer-support intervention ($p=0.797$) (**Table S4**). In a per-protocol analysis limited to female participants who attended at least one AYFS clinic visit ($n=428$) and adjusted for age and urban vs rural residence, report of any contraceptive use at the end of the study did not differ by assignment to the SRH ($p=0.830$) or peer navigator intervention ($p=0.798$) (**Table S5**). At the same time point, 47/667 (7.1%, 95%CI 5.2-9.3%) female participants reported being pregnant; among women ages 16-19, 14/237 (5.9%, 95%CI 3.3-9.7%) reported being pregnant (**Table S6**). Pregnancy did not differ by assignment to the peer navigator intervention ($p=0.67$), but there was some evidence of fewer pregnancies under assignment to the SRH intervention ($p=0.064$, **Table S7**).

Discussion

To our knowledge, this is one of the first trials to examine the effectiveness of comprehensive SRH services including home-based self-sampling for STI testing and peer navigator support for STIs and SRH outcomes among a representative sample of AYA in southern Africa. We did not find clear evidence that either access to home-based, self-collected STI screening with expanded SRH services or peer support individually reduced curable STIs or pregnancy, or increased contraception use, after 12 months of follow-up when compared to an enhanced SoC. However, we did find some evidence of a potential ‘additive’ effect of the SRH and peer support interventions in combination to reduce prevalent STIs, and an effect for both interventions individually among those participants who did attend youth-friendly clinic services during the study period.

We found high STI prevalence in this cohort, both at baseline (22.4%)²⁸ and at 12 months (22.5%). Among participants with STI results at both timepoints, 13.6% acquired a new STI during the study period. While 12-month STI prevalence did not clearly differ by assignment to either intervention, there was evidence of a potential additive effect (additive on the log-odds scale, without interaction) when interventions were combined. Additionally, in a per-protocol analysis limited only to participants who attended at least one AYFS clinic visit during the 12-month study period, both the SRH and peer support interventions were significantly associated with a lower 12-month STI prevalence. The primary outcomes analysis of the trial demonstrated that combined peer support and SRH interventions increased retention in AYFS services.²⁶ This suggests that the two interventions may bolster the effect of the AYFS services on STI risk, in part by improving retention in services, but the effect of each intervention on its own, without associated and ongoing connection to services, may be modest. Combined and more intensive approaches may thus be needed in this setting, such as repeat STI testing, point-of-care testing, test of cure, and enhanced partner services including expedited treatment. Further, structural drivers such as poverty, gender inequality, and lack of healthcare access likely play a significant role in the high STI prevalence in this population.²⁹ Thus, conceptually different approaches are likely also needed to impact STI rates, such as a focus on broader social determinants of sexual health behaviors, focus on the efficacy of counseling delivered, and evaluation of ways to enhance engagement, uptake, and continuity of attendance to supplement interventions and maximize efficacy and reach.

Among participants found to have the same STI at baseline and 12 months, the majority were treated following their baseline testing, suggesting a high rate of reinfections. Screening for curable STIs needs to be incorporated into a broader intervention and delivery mechanism, with effective

tools to prevent subsequent reinfection. Partner services, including partner notification, assisted partner services, and expedited partner therapy are all aimed at decreasing STI reinfection.³⁰ However, uptake of partner notification has been low among AYA in sub-Saharan Africa, and efforts to strengthen partner notification services have not consistently led to decreased incidence rates.³⁰ Additional strategies, such as improving the acceptability of partner notification services, and enhanced approaches such as expedited partner therapy, may be needed to strengthen the impact of screening and treatment.

Among those testing positive for STIs at baseline, over 40% were unable to be treated within four weeks of specimen collection,²⁸ primarily due to difficulty contacting participants. High rates of incomplete or delayed STI treatment were also found in a study in Zimbabwe providing community-based STI testing to AYA.³¹ These findings highlight the need for point-of-care tests (POCTs) to enable treatment within the same clinic- or community-based encounter. Development of POCTs that meet WHO REASSURED criteria is a priority for STI control worldwide.³² POCTs could both optimize rates of STI treatment and expand STI testing capacity outside of clinical settings to increase access. Accessible, cost-effective tests could also enable more frequent STI screening, in line with recommendations for repeated STI screening for individuals at increased-risk³³ and findings that frequent screening is likely needed to lower prevalence.³⁴

We found high uptake of home-based STI specimen self-collection among this cohort. Moreover, those who had been offered home-based testing at baseline were more likely to accept it at study end. In the clinical trial, the SRH intervention increased linkage to AYFS and retention in risk-differentiated HIV prevention and care.²⁶ These results suggest STI testing is acceptable and has

the potential to increase awareness and engagement in STI, SRH, and HIV care. SRH services including STI testing and treatment may thus serve as an acceptable pathway to HIV testing and risk-differentiated prevention for AYA in these settings. They may also be a mechanism for prevention of reinfection.

We additionally found that rates of self-reported contraceptive use and pregnancy at 12 months did not differ by study arm. This lack of difference may reflect the strength of the ‘enhanced SoC, as the accessibility and acceptability of the AYFS may have led to similar exposure to family planning education and access across study arms, irrespective of randomization. The findings may also reflect the fact that this age group encompasses the most common age for childbirth in this society and the balance between STI prevention and pregnancy intentions may need to be balanced. Overall rates of reported contraceptive use were similar to those estimated nationally for South Africa ³⁵. The teenage pregnancy rate of 5.9% overall was lower than data from this setting previously (11%) (unpublished data), suggesting that accessible SRH services may be a useful tool in reducing the persistently high rates of teenage and unintended pregnancy in South Africa.

This study has several limitations. Just over half of trial participants could be followed up and provided consent for STI testing at 12 months and thus included in our analyses. Rates of new or repeat STIs could only be determined for participants randomized to study arms with baseline STI testing; these numbers also limited our power to detect a difference in STI acquisition rates by peer navigator support. Participants randomized to SRH study arms and females were more likely to accept STI testing at 12 months, potentially introducing bias in the 12-month STI prevalence rates. The small absolute number of reported pregnancies at study end may have limited our power to

detect a difference in pregnancy rate by arm; further, self-reported pregnancy at study end would have missed new pregnancies during the study period that terminated or came to term prior to the 12-month endline. Confining pregnancy measures to females furthermore provides little insight into impacts on male reproductive behavior. Finally, the study period spanned the highest level of lockdowns due to COVID-19 (beginning late March 2020), when study activities paused and peer support moved to a virtual format; the subsequent process evaluation suggested that young people experienced limitations in phone and data access during this time.²⁷ Peer navigator support then remained virtual after the remainder of study activities resumed, affecting the fidelity of the peer support intervention and potentially limiting the ability to detect an impact of this intervention.

In conclusion, we found a high prevalence of STIs and new infections over 12 months in a cohort of AYA in South Africa, with no clear effect from exposure to baseline STI testing or peer navigator support, but a potential additive effect of the two interventions on 12-month STI positivity. Similarly, there was no difference between arms in use of contraception or reported pregnancy. While the ability to detect differences between arms may have been impacted by the enhanced SoC available to all and changes in the peer navigator intervention due to the COVID-19 pandemic, these results suggest that more intensive, combination interventions may impact STI and other SRH outcomes among AYA in rural South Africa.

Acknowledgments

The authors thank all the participants who contributed data to this study, the AHRI Clinical Laboratory for the laboratory services provided for this study, as well as the entire Africa Health Research Institute (AHRI) research team.

Declaration of interests

JJ has received in-kind research support from binx health. All other authors declare no competing interests.

This research was funded in whole, or in part, by the Wellcome Trust [Grant numbers 201433/Z/16/Z and 210479/Z/18/Z]. For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

Contributors

MS conceived the study. MS, TZ, CH, NO, JD, AC, KB, NM, JS, NC, GH, and LS designed and implemented the study. TS oversaw laboratory testing procedures. TK, NB, CH, and NO provided project supervision. JD was responsible for data management. JD, JB, and KB verified the data. JB and KB conducted the quantitative data analysis. JJ wrote the first draft of the manuscript. JB, NC, NO, JD, TS, GH, CH, NM, TZ, TK, NB, IVB, LS, AC, JS, KB, and MS read and critically revised the manuscript. All authors read and approved the final manuscript.

Data sharing

All datasets generated from this study are publicly available through the Africa Health Research Institute data repository site (<https://data.ahri.org/index.php/home>). The full study protocol, study data collection tools, and consent forms are available from the corresponding author (m.shahmanesh@ucl.ac.uk).

ACCEPTED

References

- 1 Rowley J, Vander Hoorn S, Korenromp E, *et al.* Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016. *Bull World Health Organ* 2019; **97**: 548-562P.
- 2 Kularatne RS, Niit R, Rowley J, *et al.* Adult gonorrhea, chlamydia and syphilis prevalence, incidence, treatment and syndromic case reporting in South Africa: Estimates using the Spectrum-STI model, 1990-2017. *PLOS ONE* 2018; **10**: e0205863.
- 3 Kharsany ABM, McKinnon LR, Lewis L, *et al.* Population prevalence of sexually transmitted infections in a high HIV burden district in KwaZulu-Natal, South Africa: Implications for HIV epidemic control. *Int J Infect Dis* 2020; **98**: 130–7.
- 4 Francis SC, Mthiyane TN, Baisley K, *et al.* Prevalence of sexually transmitted infections among young people in South Africa: A nested survey in a health and demographic surveillance site. ; : e1002512.
- 5 Johnson LF, Lewis DA. The effect of genital tract infections on HIV-1 shedding in the genital tract: a systematic review and meta-analysis. *Sex Transm Dis* 2008; **35**: 946–59.
- 6 Sexton J, Garnett G, Rottingen J-A. Metaanalysis and metaregression in interpreting study variability in the impact of sexually transmitted diseases on susceptibility to HIV infection. *Sex Transm Dis* 2005; **32**: 351–7.
- 7 Barron P, Subedar H, Letsoko M, Makua M, Pillay Y. Teenage births and pregnancies in South Africa, 2017 - 2021 – a reflection of a troubled country: Analysis of public sector data. *SAMJ* 2022; **112**: 252–8.

- 8 den Heijer CDJ, Hoebe CJP, Driessen JHM, *et al.* Chlamydia trachomatis and the Risk of Pelvic Inflammatory Disease, Ectopic Pregnancy, and Female Infertility: A Retrospective Cohort Study Among Primary Care Patients. *Clin Infect Dis* 2019; **69**: 1517–25.
- 9 Reekie J, Donovan B, Guy R, *et al.* Risk of Ectopic Pregnancy and Tubal Infertility Following Gonorrhea and Chlamydia Infections. *Clin Infect Dis* 2019; **69**: 1621–3.
- 10 Warr AJ, Pintye J, Kinuthia J, *et al.* Sexually transmitted infections during pregnancy and subsequent risk of stillbirth and infant mortality in Kenya: a prospective study. *Sex Transm Infect* 2019; **95**: 60–6.
- 11 Vallely LM, Egli D, Wand H, *et al.* Adverse pregnancy and neonatal outcomes associated with *Neisseria gonorrhoeae*: systematic review and meta-analysis. *Sex Transm Infect* 2021; **97**: 104–11.
- 12 World Health Organization. Guidelines for the management of symptomatic sexually transmitted infections. Geneva: World Health Organization, 2021 <https://apps.who.int/iris/handle/10665/342523> (accessed Oct 30, 2021).
- 13 Mlisana K, Naicker N, Werner L, *et al.* Symptomatic vaginal discharge is a poor predictor of sexually transmitted infections and genital tract inflammation in high-risk women in South Africa. *The Journal of Infectious Diseases* 2012; **206**: 6–14.
- 14 Zemouri C, Wi TE, Kiarie J, *et al.* The performance of the vaginal discharge syndromic management in treating vaginal and cervical infection: a systematic review and meta-analysis. *PLoS ONE* 2016; **11**: e0163365.
- 15 Newton-Levinson A, Leichter JS, Chandra-Mouli V. Help and Care Seeking for Sexually Transmitted Infections Among Youth in Low- and Middle-Income Countries. *Sex Transm Dis* 2017; **44**: 319–28.

- 16 Zuma T, Seeley J, Mdluli S, *et al.* Young people's experiences of sexual and reproductive health interventions in rural KwaZulu-Natal, South Africa. *Int J Adolesc Youth* 2020; **25**: 1058–75.
- 17 World Health Organization. Global health sector strategies on, respectively, HIV, viral hepatitis and sexually transmitted infections for the period 2022-2030. Geneva: World Health Organization, 2022 <https://apps.who.int/iris/handle/10665/360348> (accessed Aug 26, 2022).
- 18 Hoffman CM, Mbambazela N, Sithole P, *et al.* Provision of sexually transmitted infection services in a mobile clinic reveals high unmet need in remote areas of South Africa: a cross-sectional study. *Sexually Transmitted Diseases* 2019; **46**: 206–12.
- 19 Oluoch LM, Roxby A, Mugo N, *et al.* Does providing laboratory confirmed STI results impact uptake of HIV pre-exposure prophylaxis (PrEP) uptake among Kenyan adolescents girls and young women? A descriptive analysis. *Sex Transm Infect* 2020; : sextrans-2020-054637.
- 20 Shahmanesh M, Mthiyane TN, Herbst C, *et al.* Effect of peer-distributed HIV self-test kits on demand for biomedical HIV prevention in rural KwaZulu-Natal, South Africa: a three-armed cluster-randomised trial comparing social networks versus direct delivery. *BMJ Glob Health* 2021; **6**: e004574.
- 21 Chidumwa G, Chimbindi N, Herbst C, *et al.* Isisekelo Sempilo study protocol for the effectiveness of HIV prevention embedded in sexual health with or without peer navigator support (Thetha Nami) to reduce prevalence of transmissible HIV amongst adolescents and young adults in rural KwaZulu-Natal: a 2 × 2 factorial randomised controlled trial. *BMC Public Health* 2022; **22**: 454.
- 22 Gareta D, Baisley K, Mngomezulu T, *et al.* Cohort Profile Update: Africa Centre Demographic Information System (ACDIS) and population-based HIV survey. *Int J Epidemiol* 2021; **50**: 33–40.

- 23 Shahmanesh M, Okesola N, Chimbindi N, *et al.* Thetha Nami: participatory development of a peer-navigator intervention to deliver biosocial HIV prevention for adolescents and youth in rural South Africa. *BMC Public Health* 2021; **21**: 1393.
- 24 South Africa Department of Health. 2021 Updated Guidelines for the Provision of Oral Pre-exposure Prophylaxis (PrEP) to Persons at Substantial risk of HIV Infection. South Africa, 2021.
- 25 South Africa Department of Health. Sexually Transmitted Infections Management Guidelines 2018. Pretoria, South Africa, 2018.
- 26 Shahmanesh M, Chimbindi N, Busang J, *et al.* Effectiveness of integrating HIV prevention within sexual reproductive health services with or without peer support among adolescents and young adults in rural KwaZulu-Natal, South Africa (Isisekelo Sempilo): 2 × 2 factorial, open-label, randomised controlled trial. *The Lancet HIV* 2024; **11**: e449–60.
- 27 Zuma T, Busang J, Hlongwane S, *et al.* A mixed methods process evaluation: understanding the implementation and delivery of HIV prevention services integrated within sexual reproductive health (SRH) with or without peer support amongst adolescents and young adults in rural KwaZulu-Natal, South Africa. *Trials* 2024; **25**: 448.
- 28 Jarolimova J, Chidumwa G, Chimbindi N, *et al.* Prevalence of Curable Sexually Transmitted Infections in a Population-Representative Sample of Young Adults in a High HIV Incidence Area in South Africa. *Sexual Trans Dis* 2023; **50**: 796–803.
- 29 Mannell J, Willan S, Shahmanesh M, Seeley J, Sherr L, Gibbs A. Why interventions to prevent intimate partner violence and HIV have failed young women in southern Africa. *J Int AIDS Soc* 2019; **22**: e25380.

- 30 Taleghani S, Joseph-Davey D, West SB, Klausner HJ, Wynn A, Klausner JD. Acceptability and efficacy of partner notification for curable sexually transmitted infections in sub-Saharan Africa: A systematic review. *Int J STD AIDS* 2019; **30**: 292–303.
- 31 Martin K, Olaru ID, Buwu N, *et al.* Uptake of and factors associated with testing for sexually transmitted infections in community-based settings among youth in Zimbabwe: a mixed-methods study. *Lancet Child Adolesc Health* 2021; **5**: 122–32.
- 32 Land KJ, Boeras DI, Chen X-S, Ramsay AR, Peeling RW. REASSURED diagnostics to inform disease control strategies, strengthen health systems and improve patient outcomes. *Nat Microbiol* 2019; **4**: 46–54.
- 33 Peters RPH, Garrett N, Chandiwana N, *et al.* Southern African HIV Clinicians Society 2022 guideline for the management of sexually transmitted infections: Moving towards best practice. *South Afr J HIV Med* 2022; **23**: 1450.
- 34 Hui BB, Hocking JS, Braat S, *et al.* Intensified partner notification and repeat testing can improve the effectiveness of screening in reducing *Chlamydia trachomatis* prevalence: a mathematical modelling study. *Sex Transm Infect* 2022; **98**: 414–9.
- 35 Haakenstad A, Angelino O, Irvine CMS, *et al.* Measuring contraceptive method mix, prevalence, and demand satisfied by age and marital status in 204 countries and territories, 1970–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* 2022; **400**: 295–327.

Figure legends.

Figure 1. CONSORT Diagram. ¹Enhanced Standard of Care - adolescent & youth friendly services (AYFS). ²Sexual and reproductive health component. ³Peer navigator component

Figure 2. STI positivity at 12 months (n=927)

Figure 3. STI positivity at 12 months by trial interventions, compared to non-receipt of each intervention (A) and by trial arm, compared to enhanced standard of care (arm 1), adjusted for age, sex, and urban vs rural residence) (B).

Tables

Table 1. Prevalence of any STI, chlamydia, gonorrhea, and trichomoniasis at 12 months (N=927)

Table 2. New sexually transmitted infections (STIs) at 12 months (N=469)

Supplemental Digital Content

1. Isisekelo SRH outcomes_supp material.docx contains:

Table S1. Results from unadjusted and adjusted models for characteristics associated with being reached at 12 months and consenting to testing for STIs among the 1743 study participants

Table S2. Prevalence of any STI at 12 months among those who linked to care during the study period

Table S3. New infections by individual STI at 12 months (N=469)

Table S4. Contraception use at 12 months among all females who linked to care during the study period

Table S5. Unweighted prevalence of self-reported pregnancy among adolescent and young women at 12 months (N=667)

Figure 1




Figure 2




Figure 3




Table 1. Prevalence of any STI, chlamydia, gonorrhea, and trichomoniasis at 12 months (N=927)*

	Number with outcome/ total (%)	Unadjusted OR (95% CI)	p-value	Adjusted OR ¹ (95% CI)	p-value
Prevalence of any STI at 12 months					
Overall SRH²	209/927 (22.5)		P=0.100		P=0.115
No	109/437 (24.9)	1		1	
Yes	100/490 (20.4)	0.77 (0.57, 1.05)		0.74 (0.56, 1.06)	
Peer support			P=0.168		P=0.104
No	110/449 (24.5)	1		1	
Yes	99/478 (20.7)	0.81 (0.59, 1.10)		0.77 (0.56, 1.06)	
Trial arm			P=0.200		P=0.162
Enhanced SoC ³	57/211 (27.0)	1		1	
SRH	53/238 (22.3)	0.77 (0.50, 1.19)		0.78 (0.50, 1.22)	
Peer support	52/226 (23.0)	0.81 (0.52, 1.25)		0.77 (0.49, 1.21)	
SRH + peer support	47/252 (18.7)	0.62 (0.40, 0.96)		0.59 (0.38, 0.94)	
Prevalence of chlamydia at 12 months					
Overall SRH²	163/927 (17.6)		P=0.080		P=0.099
No	87/437 (19.9)	1		1	
Yes	76/490 (15.5)	0.74 (0.53, 1.04)		0.75 (0.53, 1.06)	
Peer support			P=0.165		P=0.116
No	87/449 (19.4)	1		1	
Yes	76/478 (15.9)	0.79 (0.56, 1.10)		0.76 (0.54, 1.07)	
Trial arm			P=0.157		P=0.145
Enhanced SoC	45/211 (21.3)	1		1	
SRH	42/238 (17.7)	0.79 (0.49, 1.26)		0.80 (0.50, 1.29)	
Peer support	42/226 (18.6)	0.84 (0.53, 1.35)		0.81 (0.50, 1.31)	
SRH + peer support	34/252 (13.5)	0.58 (0.35, 0.94)		0.56 (0.34, 0.92)	
Prevalence of gonorrhea at 12 months					
Overall SRH	54/927 (5.8)		P=0.204		P=0.256
No	30/437 (6.9)	1		1	
Yes	24/490 (4.9)	0.70 (0.40, 1.21)		0.72 (0.41, 1.27)	
Peer support			P=0.813		P=0.723
No	27/449 (6.0)	1		1	

Yes	27/478 (5.7)	0.94 (0.54, 1.62)		0.90 (0.52, 1.58)	
Trial arm			P=0.640		P=0.698
Enhanced SoC	15/211 (7.1)	1		1	
SRH	12/238 (5.0)	0.69 (0.32, 1.52)		0.71 (0.32, 1.56)	
Peer support	15/226 (6.6)	0.93 (0.44, 1.95)		0.89 (0.42, 1.88)	
SRH + peer support	12/252 (4.8)	0.65 (0.44, 1.43)		0.65 (0.30, 1.44)	
Prevalence of trichomoniasis at 12 months					
Overall	44/927 (4.8)				
SRH			P=0.697		P=0.598
No	22/437 (5.0)	1		1	
Yes	22/490 (4.5)	0.89 (0.48, 1.62)		0.85 (0.45, 1.58)	
Peer support			P=0.407		P=0.353
No	24/449 (5.4)	1		1	
Yes	20/478 (4.2)	0.77 (0.42, 1.42)		0.74 (0.40, 1.39)	
Trial arm			P=0.740		P=0.724
Enhanced SoC	13/211 (6.2)	1		1	
SRH	11/238 (4.6)	0.74 (0.32, 1.68)		0.75 (0.32, 1.74)	
Peer support	9/226 (4.0)	0.63 (0.26, 1.51)		0.65 (0.26, 1.58)	
SRH + peer support	11/252 (4.4)	0.70 (0.30, 1.59)		0.64 (0.28, 1.50)	

* Includes data from participants who provided STI specimens and had valid test results for all three STIs during the 12-month survey

¹Adjusted for sex, age group, and location of residence

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table 2. New sexually transmitted infections (STIs) at 12 months (N=469)*

	Overall N=469	SRH¹ N=226	SRH + peer support N=243	p-value²
Any STI at baseline, n (%)	111 (23.7)	55 (24.3)	56 (23.1)	0.742
Any STI at endline, n (%)	95 (20.3)	49 (21.7)	46 (18.9)	0.459
Any new STI, n (%) ³	64 (13.6)	31 (13.7)	33 (13.7)	0.966
Diagnosed with same STI at endline as at baseline, n (%)	39 (8.3)	21 (9.3)	18 (7.4)	0.460

*Includes data from participants with valid STI results at baseline and at 12 months (limited to participants in the IS/SRH arms only)

¹Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

²Chi-square test of independence

³Participants who tested negative for any individual STI at baseline and positive for that STI at 12 months

Table 3. Contraception use among all females at the end of the study (N=634)

	Number self-reporting contraception use/total (%)	Unadjusted OR (95% CI)	Adjusted OR ¹ (95% CI)
Overall	336/634 (53.0)		
SRH²		P=0.152	P=0.301
No	150/300 (50.0)	1	1
Yes	186/148 (44.3)	1.26 (0.92, 1.72)	1.19 (0.86, 1.65)
Peer support		P=0.940	P=0.797
No	167/316 (52.9)	1	1
Yes	169/318 (53.1)	1.01 (0.74, 1.38)	1.04 (0.75, 1.45)
Trial arm		P=0.560	P=0.716
Enhanced SoC ³	76/152 (50.0)	1	1
SRH	91/164 (55.5)	1.25 (0.80, 1.94)	1.29 (0.81, 2.05)
Peer support	74/148 (50.0)	1.00 (0.64, 1.57)	1.13 (0.70, 1.82)
SRH + peer support	95/170 (44.1)	1.27 (0.82, 1.96)	1.24 (0.78, 1.97)

¹Adjusted for, age group, and location of residence

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table 4. Female participants who self-reported pregnancy at the end of the study (N=667)¹

	Number self-reporting contraception use/total (%)	Unadjusted OR (95% CI)	Adjusted OR ² (95% CI)
Overall	47/667 (7.1)		
SRH³		P=0.093	P=0.064
No	28/318 (8.8)	1	1
Yes	19/349 (5.4)	0.60 (0.33, 1.09)	0.57 (0.31, 1.04)
Peer support		P=0.689	P=0.668
No	22/331 (6.7)	1	1
Yes	25/336 (7.4)	1.13 (0.62, 2.05)	1.14 (0.63, 2.07)
Trial arm		P=0.560	P=0.297
Enhanced SoC ⁴	13/159 (8.2)	1	1
SRH	9/172 (5.2)	0.62 (0.26, 1.49)	0.61 (0.25, 1.46)
Peer support	15/159 (9.4)	1.17 (0.54, 2.55)	1.22 (0.56, 2.67)
SRH + peer support	10/177 (5.7)	0.67 (0.29, 1.58)	0.65 (0.27, 1.52)

¹Includes data from female participants who completed the 12-month survey and answered 'yes' or 'no' regarding self-reported pregnancy

²Adjusted for age group, and location of residence

³Sexual and Reproductive Health: Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

⁴Enhanced standard of care

Youth-friendly sexual health services and peer support for improved sexual and reproductive health outcomes among adolescents and young adults in South Africa: results of a factorial randomized controlled trial

Jarolimova et al.

Supplementary materials

ACCEPTED

Table of contents

Figure S1. Schematic of study procedures.....	3
Table S1. Results from unadjusted and adjusted models for characteristics associated with being reached at 12 months and consenting to testing for STIs among the 1743 study participants.....	4
Table S2. Prevalence of any STI at 12 months among those who linked to care during the study period.	5
Table S3. New infections by individual STI at 12 months (N=469)*.	6
Table S4. Contraception use among all females at the end of the study (N=634).....	7
Table S5. Contraception use at 12 months among all females who linked to care during the study period.	8
Table S6. Unweighted prevalence of self-reported pregnancy among adolescent and young women at 12 months (N=667) ¹	9
Table S7. Female participants who self-reported pregnancy at the end of the study (N=667).....	10

Figure S1. Schematic of study procedures.




Table S1. Results from unadjusted and adjusted models for characteristics associated with being reached at 12 months and consenting to testing for STIs among the 1743 study participants.

Characteristic	Overall, n (%)	Proportion reached and consented to STI testing, n/N (%)	Unadjusted OR (95% CI)	P-value	Adjusted ¹ OR (95% CI)	P-value
Overall	1743	987/1743 (56.6)				
SRH²						
No	880 (50.5)	465/880 (52.8)	Ref	0.001	Ref	0.002
Yes	863 (49.5)	522/863 (60.5)	1.37 (1.13, 1.65)		1.35 (1.11, 1.64)	
Peer support						
No	858 (49.2)	476/858 (55.5)	Ref	0.487	Ref	0.381
Yes	885 (50.8)	511/885 (57.7)	1.10 (0.91, 1.33)		1.09 (0.90, 1.32)	
Study arm						
Enhanced SoC ³	435 (25.0)	221/435 (50.8)	Ref	0.008	-	
SRH	423 (24.3)	255/423 (60.3)	1.47 (1.12, 1.93)			
Peer support	445 (25.5)	244/445 (54.8)	1.18 (0.90, 1.53)			
SRH + peer support	440 (25.2)	267/440 (60.7)	1.49 (1.14, 1.95)			
Sex						
Male	847 (48.6)	459/847 (54.2)	Ref	0.046	Ref	0.039
Female	896 (51.4)	528/896 (58.9)	1.21 (1.00, 1.47)		1.23 (1.01, 1.49)	
Age group						
16-19	639 (36.7)	359/639 (56.2)	Ref	0.767	Ref	0.838
20-25	625 (35.9)	361/625 (57.8)	1.07 (0.85, 1.33)		0.95 (0.73, 1.25)	
25-29	479 (27.5)	267/479 (55.7)	0.98 (0.77, 1.25)		0.91 (0.66, 1.25)	
Area of residence						
Rural	1,082 (62.1)	639/1,082 (59.1)	Ref	0.009	Ref	0.049
Urban/Peri-Urban	661 (37.9)	348/661 (52.6)	0.77 (0.6, 0.94)		0.82 (0.67, 1.00)	
Highest level of education attained;						
Primary	644 (37.0)	370/644 (57.5)	Ref	0.363	Ref	0.411
Secondary	963 (55.3)	534/963 (55.5)	0.92 (0.75, 1.13)		0.93 (0.72, 1.19)	
Post-secondary	135 (7.7)	83/135 (61.5)	1.18 (0.81, 1.73)		1.20 (0.81, 1.78)	
Employment						
Not employed	1,239 (71.9)	745/1,239 (60.1)	Ref	0.001	Ref	0.001
Employed	154 (8.9)	73/154 (47.4)	0.60 (0.43, 0.84)		0.66 (0.47, 0.94)	
Studying	331 (19.2)	169/331 (51.1)	0.69 (0.54, 0.88)		0.65 (0.49, 0.86)	

¹Model adjusted for all characteristics excluding 'study arm' (due to collinearity)

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table S2. Prevalence of any STI at 12 months among those who linked to care during the study period.

	Number with outcome/total (%)	Unadjusted OR (95% CI)	Adjusted OR ¹ (95% CI)
Overall	201/856 (23.5)		
SRH²		P=0.014	P=0.026
No	102/370 (27.6)	1	1
Yes	99/486 (20.4)	0.67 (0.49 -0.92)	0.69 (0.50 -0.96)
Peer support		P=0.046	P=0.030
No	103/386 (26.7)	1	1
Yes	98/470 (20.9)	0.72 (0.53 -0.99)	0.70 (0.50 -0.97)
Trial arm		P=0.009	P=0.012
Enhanced SoC ³	51/203 (25.1)	1	1
SRH	52/300 (17.3)	0.57 (0.36 -0.89)	0.59 (0.37 -0.94)
Peer support	51/311 (16.4)	0.60 (0.38 -0.96)	0.59 (0.37 -0.95)
SRH + peer support	47/331 (14.2)	0.45 (0.29 -0.72)	0.45 (0.28 -0.73)

¹Adjusted for sex, age group, and location of residence

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table S3. New infections by individual STI at 12 months (N=469)*.

	New diagnosis of each STI at endline, n/N (%) ¹	Same STI positive at baseline and endline, n/N (%)	Repeat positive tests in those treated (reinfections)	Repeat positive tests in those not treated (persistent infections)
Chlamydia	42/469 (9.0)	31/469 (6.6)	27/31 (87.1)	4/31 (12.9)
Gonorrhea	18/469 (3.8)	4/467 (0.9)	3/4 (75.0)	1/4 (25.0)
Trichomoniasis	13/469 (2.8)	7/469 (1.5)	7/7 (100)	0

*Includes data from participants with valid STI results at both baseline and 12 months (limited to participants in the IS/SRH arms only)

¹Participants who tested negative for each STI at baseline and positive for that STI at 12 months

Table S4. Contraception use among all females at the end of the study (N=634)

	Number self-reporting contraception use/total (%)	Unadjusted OR (95% CI)	Adjusted OR ¹ (95% CI)
Overall	336/634 (53.0)		
SRH²		P=0.152	P=0.301
No	150/300 (50.0)	1	1
Yes	186/148 (44.3)	1.26 (0.92, 1.72)	1.19 (0.86, 1.65)
Peer support		P=0.940	P=0.797
No	167/316 (52.9)	1	1
Yes	169/318 (53.1)	1.01 (0.74, 1.38)	1.04 (0.75, 1.45)
Trial arm		P=0.560	P=0.716
Enhanced SoC ³	76/152 (50.0)	1	1
SRH	91/164 (55.5)	1.25 (0.80, 1.94)	1.29 (0.81, 2.05)
Peer support	74/148 (50.0)	1.00 (0.64, 1.57)	1.13 (0.70, 1.82)
SRH + peer support	95/170 (44.1)	1.27 (0.82, 1.96)	1.24 (0.78, 1.97)

¹Adjusted for, age group, and location of residence

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table S5. Contraception use at 12 months among all females who linked to care during the study period.

	Number self-reporting contraception use/total (%)	Unadjusted OR (95% CI)	Adjusted OR ¹ (95% CI)
Overall	235/428 (54.9)		
SRH²		P=0.552	P=0.830
No	98/370 (26.5)	1	1
Yes	137/486 (28.2)	1.12 (0.76 -1.65)	1.05 (0.70 -1.56)
Peer support		P=0.919	P=0.798
No	106/386 (27.5)	1	1
Yes	129/470 (27.4)	1.02 (0.70 -1.50)	1.05 (0.71 -1.57)
Trial arm		P=0.933	P=0.986
Enhanced SoC ³	43/203 (21.2)	1	1
SRH	63/300 (21.0)	1.06 (0.60 -1.89)	1.09 (0.60 -1.97)
Peer support	55/311 (17.7)	0.97 (0.54 -1.73)	1.10 (0.60 -2.01)
SRH + peer support	74/331 (22.4)	1.14 (0.65 -1.99)	1.11 (0.62 -1.99)

¹Adjusted for sex, age group, and location of residence

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table S6. Unweighted prevalence of self-reported pregnancy among adolescent and young women at 12 months (N=667)¹.

Study arm	Overall	16-19 years	20-29 years
	Number pregnant/total (%)		
Overall	47/667 (7.1)	14/237 (5.9)	33/430 (7.7)
SRH²			
No	28/318 (8.8)	8/120 (6.7)	20/198 (10.1)
Yes	19/349 (5.4)	6/117 (5.1)	13/232 (5.6)
Peer support			
No	22/331 (6.7)	6/112 (5.4)	16/219 (7.3)
Yes	25/336 (7.4)	8/125 (6.4)	17/211 (8.1)
Trial arm			
Enhanced SoC ³	13/159 (8.2)	3/52 (5.6)	10/107 (9.4)
SRH	9/172 (5.2)	3/60 (5.0)	6/112 (5.4)
Peer support	15/159 (9.4)	5/68 (7.4)	10/91 (11.0)
SRH + peer support	10/177 (5.7)	3/57 (5.3)	7/120 (5.8)

¹Includes data from female participants who completed the 12-month survey and answered 'yes' or 'no' regarding self-reported pregnancy

²Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

³Enhanced standard of care

Table S7. Female participants who self-reported pregnancy at the end of the study (N=667)¹

	Number self-reporting contraception use/total (%)	Unadjusted OR (95% CI)	Adjusted OR ² (95% CI)
Overall	47/667 (7.1)		
SRH³		P=0.093	P=0.064
No	28/318 (8.8)	1	1
Yes	19/349 (5.4)	0.60 (0.33, 1.09)	0.57 (0.31, 1.04)
Peer support		P=0.689	P=0.668
No	22/331 (6.7)	1	1
Yes	25/336 (7.4)	1.13 (0.62, 2.05)	1.14 (0.63, 2.07)
Trial arm		P=0.560	P=0.297
Enhanced SoC ⁴	13/159 (8.2)	1	1
SRH	9/172 (5.2)	0.62 (0.26, 1.49)	0.61 (0.25, 1.46)
Peer support	15/159 (9.4)	1.17 (0.54, 2.55)	1.22 (0.56, 2.67)
SRH + peer support	10/177 (5.7)	0.67 (0.29, 1.58)	0.65 (0.27, 1.52)

¹Includes data from female participants who completed the 12-month survey and answered 'yes' or 'no' regarding self-reported pregnancy

²Adjusted for age group, and location of residence

³Sexual and Reproductive Health: Adolescent and youth friendly sexual and reproductive health services including home-based STI self-sampling

⁴Enhanced standard of care