

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Nascimento, MC; Sumita, LM; Souza, VU; Weiss, HA; Oliveira, J; Mascheretti, M; Quiroga, M; Vela, RA; Sabino, E; Pannuti, CS; +1 more... Mayaud, P; (2009) Seroprevalence of Kaposi sarcoma-associated herpesvirus and other serologic markers in the Brazilian Amazon. *Emerging infectious diseases*, 15 (4). pp. 663-7. ISSN 1080-6040 DOI: <https://doi.org/10.3201/eid1504.081488>

Downloaded from: <http://researchonline.lshtm.ac.uk/5663/>

DOI: <https://doi.org/10.3201/eid1504.081488>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Seroprevalence of Kaposi Sarcoma-associated Herpesvirus and Other Serologic Markers in the Brazilian Amazon

Maria C. Nascimento, Laura M. Sumita, Vanda U. Souza, Helen A. Weiss, Juliane Oliveira, Melissa Mascheretti, Mariana Quiroga, Rodrigo A.R. Vela, Ester Sabino, Claudio S. Pannuti, and Philippe Mayaud

To determine the presence of Kaposi sarcoma-associated herpesvirus (KSHV) and other serologic markers, we tested serum specimens of 339 Amerindians, 181 rural non-Amerindians, and 1,133 urban blood donors (13 Amerindians) in the Brazilian Amazon. High KSHV seroprevalence in children and inverse association with herpes simplex virus type 2 indicates predominant nonsexual transmission among Amerindians.

Kaposi sarcoma-associated herpesvirus (KSHV) is the cause of Kaposi sarcoma (KS) and certain lymphoproliferative diseases (1). KSHV seroprevalence is low (<5%) in most Western populations (1) and reaches 50% in some African populations (2), mirroring KS incidence rates (3). However, the highest KSHV seroprevalences worldwide (>80% in adults) have been reported in Amerindian tribes from the Amazon regions of Brazil (4,5) and Ecuador (6), despite the apparently low KS incidence in these populations (7). KSHV is thought to be transmitted through saliva between young siblings in disease-endemic areas such as French Guiana (8) or Africa (9), whereas sexual transmission in low-prevalence countries occurs within risk groups such as men who have sex with men (10). Modes of transmission have not been clearly determined in Amerindian populations.

Author affiliations: Universidade de São Paulo, São Paulo, Brazil (M.C. Nascimento, L.M. Sumita, V.U. Souza, J. Oliveira, M. Mascheretti, M. Quiroga, R.A.R. Vela, E. Sabino, C.S. Pannuti); and London School of Hygiene and Tropical Medicine, London, UK (M.C. Nascimento, H.A. Weiss, P. Mayaud)

DOI: 10.3201/eid1504.081488

The Study

We conducted a cross-sectional study during February 2003–April 2004 to investigate the seroprevalence and factors associated with KSHV infection in Amerindian and non-Amerindian populations living in 2 regions of the Brazilian Amazon: a remote rural region of Para State (Mapuera, on the banks of the Trombetas River) and Manaus, the capital city of Amazonas State (Figure). Serologic markers of fecal–oral (hepatitis A virus [HAV]), blood-borne (hepatitis B and C viruses [HBV, HCV]) and sexually transmitted infections (*Treponema pallidum* [syphilis] and herpes simplex virus type 2 [HSV-2]) were used as proxies to identify possible routes of KSHV transmission in these populations.

A convenience sample of unselected Amerindians and non-Amerindians living in the Mapuera area and a consecutive sample of nonpaid first-time blood donors from the Manaus blood bank (HemoAm) consented to collection of blood samples, as previously reported (4,11) Ethical approval was obtained from the institutional review board of HemoAm, the ethical board of the Brazilian Ministry of Health, and the ethics committee of the London School of Hygiene and Tropical Medicine.

In the absence of a definitive test to determine KSHV infection, all serum specimens were tested by using a previously validated in-house whole-virus KSHV ELISA (12) and 2 immunofluorescence assays (IFAs) that detected antibodies against lytic (IFA-lytic) and latent-associated nuclear antigens (IFA-LANA) (12). KSHV infection was defined as positivity by any of these serologic assays. Serum specimens were also tested for the agent of syphilis by us-



Figure. Map of Brazil showing study area (black box) in Amazonas (Manaus) and Para (Mapuera region) States. Printed with permission of the Instituto Brasileiro de Geografia e Estatística.

ing a *T. pallidum*-specific assay (Enzygnost Syphilis; Dade Behring, Marburg, Germany); for HSV-2 antibodies by using the type-specific HerpeSelect gG2 ELISA (Focus Technologies, Cypress Hill, CA, USA), with a higher cut-off (>3.5) to increase specificity (13); and for HAV antibodies by using BioELISA HAV (Biokit, Barcelona, Spain). Presence of HBV anti-core antibodies was determined by using Ortho HBc ELISA (Ortho Diagnostics, Raritan, NJ, USA) in Mapuera serum specimens and Hepanostika anti-HBc Uni-Form (Organon-Teknika, Boxtel, the Netherlands) in Manaus serum specimens. HCV antibodies were detected by using Ortho HCV 3.0 ELISA (Ortho Diagnostics) in Mapuera serum specimens and Murex Anti-HCV version 4.0 ELISA (Murex Biotech S.A., Kyalami, South Africa) in Manaus serum specimens.

KSHV seroprevalence was calculated separately for men and women and directly age-standardized to the Mapuera Amerindian population. The risk associated with KSHV infection was estimated with prevalence ratios (PRs) and 95% confidence intervals (CIs), adjusted for sex and age group (18–24 years, 25–34 years, and ≥ 35 years for the blood donor population; 0–9 years, 10–17 years, 18–24 years, 25–34 years, and ≥ 35 years for both Mapuera populations). The associations of KSHV with sociodemographic variables, indicators of socioeconomic status, and other serologic markers were estimated with odds ratios (ORs) and 95% CIs. Variables associated with a significant increased risk for KSHV ($p < 0.05$) in univariable analysis were included in a multivariable logistic regression model adjusted for age and sex.

We recruited 339 Amerindians (median age 22 years, interquartile range [IQR] 13–37 years; 57.5% female) and 181 non-Amerindians (median age 17 years, IQR 9–35 years; 58.6% female) in the Mapuera communities and 1,133 blood donors (median age 25 years, IQR 21–32 years; 22.9% female) in Manaus. The blood donor population had a similar age distribution to that of the adult population in Manaus in the 2000 regional census (14).

Among Mapuera Amerindians, KSHV seroprevalence was 65.0% in those 0–9 years, increasing to 92.9% in those ≥ 35 years. In contrast, among Mapuera non-Amerindians, KSHV seroprevalence was 9.8% in those 0–9 years of age, increasing to 50.0% in those ≥ 35 years of age. Among blood donors, KSHV seroprevalence was 31.3% in those ≥ 35 years of age and 53.8% in the 13 who were of Amerindian descent. After age standardization, KSHV seroprevalence remained lower among Mapuera non-Amerindians (30% and 27% among men and women, respectively) and blood donors (16% and 23%, respectively) than among Mapuera Amerindians. When results were compared with those of the Mapuera Amerindians, the age- and sex-adjusted PRs were 0.35 (95% CI 0.28–0.45) and 0.59 (95% CI 0.56–0.63) in Mapuera non-Amerindians and blood donors, respectively.

In each population, KSHV seroprevalence was slightly

higher among females, and increased with age (p for trend < 0.001) in Mapuera Amerindians and non-Amerindians, but not among (adult) blood donors (Table 1). KSHV seroprevalence varied little with house crowding (socioeconomic indicator), and hepatitis infections, but was associated with HSV-2 infection in non-Amerindians (OR 4.2, 95% CI 2.1–8.5) and blood donors (OR 1.3, 95% CI 1.0–1.7). In Amerindians, KSHV infection was not associated with HSV-2 in univariable analysis (OR 0.7, 95% CI 0.3–1.9).

In multivariable analysis (Table 2), KSHV infection remained associated with female sex among blood donors (age- and sex-adjusted OR [aOR] 1.3, 95% CI 1.0–1.7), and increased significantly with age in both Mapuera populations (p for trend < 0.001). KSHV infection was associated with HSV-2 infection among Mapuera non-Amerindians (aOR 2.7, 95% CI 1.2–6.5) and Manaus blood donors (aOR 1.3, 95% CI 1.0–1.6), but was inversely associated with HSV-2 infection in Mapuera Amerindians (aOR 0.3, 95% CI 0.1–0.9).

Conclusions

Our data confirm the high KSHV seroprevalence observed among Amazonian Amerindian populations (5,7). However, the inclusion of convenience samples of remote populations and first-time blood donors, who may not necessarily be representative of the adult general population and notably exclude persons who report a range of potentially high-risk behavior for sexually transmitted and blood-borne infections, may have limited the generalizability of our findings. High KSHV seroprevalence combined with an apparent lack of KS development among Amerindian populations support the theory of genetic predisposition to KSHV acquisition, as hypothesized for other Amazonian populations, in whom segregation genetic analysis has suggested that an unidentified recessive gene may influence KSHV serostatus (15).

The high KSHV seroprevalence (65%) among Mapuera Amerindians < 10 years of age contrasts with the low (9.8%) seroprevalence among non-Amerindians of the same age group living in the same area, which suggests different transmission modes in these neighboring populations. Although we did not collect data on the age of initial sexual experience in either population, the high prevalence in childhood and inverse association with HSV-2 supports nonsexual transmission of KSHV in Amerindians. Conversely, the association of KSHV infection with HSV-2 among Mapuera non-Amerindians and blood donors supports a role for sexual transmission in these groups, although saliva transmission in younger urban inhabitants cannot be ruled out. Universal HAV infection status and low rates of HBV and HCV in all populations precluded any meaningful analysis of transmission routes associated with hepatitis viruses.

In summary, this study contributes data on the epidemiology of KSHV infection and transmission in some Brazilian Amazonian populations. Irrespective of urban or rural setting, our data are consistent with a predominant non-sexual transmission of KSHV (most likely through saliva) in Amerindian tribes compared with a probable combination of sexual and nonsexual modes of transmission among non-Amerindian populations living in the same region.

Acknowledgments

We thank Katia Torres, Adele Schwartz-Benzaken, David Mabey, and Onno Dekker for their support of the study. Data collection was carried out at the Laboratory of Virology, Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, and the Department of Infectious and Parasitic Diseases, Faculdade de Medicina, Universidade de São Paulo; laboratory testing was also performed at the Laboratory of Virology; and data analysis was conducted at the Department of Infectious and Tropical Diseases and the

Table 1. Seroprevalence of KSHV among 3 populations in the Brazilian Amazon*†

Variables	Mapuera Amerindians, n = 339†		Mapuera non-Amerindians, n = 181†		Manaus blood donors, n = 1,133†	
	% KSHV positive (no. tested)	OR (95% CI)	% KSHV positive (no. tested)	OR (95% CI)	% KSHV positive (no. tested)	OR (95% CI)
Sex						
Male	79.2 (144)	1	26.7 (75)	1	28.6 (874)	1
Female	82.6 (195)	1.2 (0.7–2.1)	27.4 (106)	1.0 (0.5–2.0)	34.4 (259)	1.3 (1.0–1.7)
p value		0.4		0.1		0.08
Age group, y						
0–9	65.0 (43)	0.1 (0.05–0.4)	9.8 (51)	0.1 (0.03–0.3)	–	–
10–17	70.0 (93)	0.2 (0.07–0.4)	22.5 (40)	0.3 (0.1–0.7)	–	–
18–34	86.5 (104)	0.5 (0.2–1.3)	27.3 (44)	0.4 (0.1–0.9)	29.6 (916)	0.9 (0.7–1.3)
≥35	92.9 (99)	1	50.0 (46)	1	31.3 (217)	1
p for trend		<0.001		<0.001		
Crowding‡						
1–2	93.7 (16)	1	55.6 (9)	1	32.6 (175)	1
3	91.3 (23)	0.7 (0.06–8.4)	33.3 (15)	0.4 (0.07–2.2)	29.9 (941)	0.9 (0.6–1.2)
≥4	79.7 (300)	0.3 (0.03–2.0)	24.8 (145)	0.3 (0.07–1.3)	6.2 (16)	0.4 (0.2–1.0)
p value		0.1		0.1		0.1
Ethnicity						
African	–	–	–	–	29.6 (743)	1
Caucasian	–	–	–	–	30.5 (308)	1.0 (0.8–1.4)
Indigenous	100 (339)	–	–	–	53.8 (13)	2.8 (0.9–8.3)
Other	–	–	100 (181)	–	25.8 (66)	0.8 (0.5–1.5)
p value						0.08
Hepatitis A virus						
Negative	83.3 (6)	1	12.5 (16)	1	42.9 (7)§	1
Positive	81.1 (333)	0.9 (0.1–7.5)	28.5 (165)	2.8 (0.6–12.7)	28.6 (154)§	0.5 (0.1–2.5)
p value		0.9		0.2		0.4
Hepatitis B virus						
Negative	81.6 (315)	1	32.0 (75)§	1	30.2 (1,075)	1
Positive	73.9 (23)	0.6 (0.2–1.7)	53.3 (15)§	2.4 (0.8–7.5)	25.0 (56)	0.8 (0.4–1.4)
p value		0.4		0.1		0.4
Hepatitis C virus						
Negative	81.0 (338)		36.0 (90)†		29.9 (1,129)	1
Positive	0		0		25.0 (4)	0.8 (0.1–7.5)
p value						0.8
HSV-2						
Negative	81.5 (314)	1	18.1 (127)	1	27.8 (715)	1
Positive	76.0 (25)	0.7 (0.3–1.9)	48.1 (54)	4.2 (2.1–8.5)	33.2 (406)	1.3 (1.0–1.7)
p value		0.5		<0.001		0.06
<i>Treponema pallidum</i>						
Negative	81.0 (338)	–	26.3 (171)		29.9 (1,122)	1
Positive	0		40.0 (10)	1	36.4 (11)	1.2 (0.6–2.3)
p value				1.9 (0.5–6.9)		0.7
				0.3		

*Seroreactivity by any serologic assay, whole virus. KSHV, Kaposi sarcoma-associated herpesvirus; OR, odds ratio; CI, confidence interval; HSV-2, herpes simplex virus type 2.

†Some figures do not add up to the total because of missing data.

‡Number of residents living in the house.

§Only a random subsample tested.

Table 2. Multivariable analysis of risk factors for KSHV infection among 3 populations in the Brazilian Amazon*

Variables	aOR (95% CI)		
	Mapuera Amerindians, n = 339	Mapuera non-Amerindians, n = 181	Manaus blood donors, n = 1,133
Sex			
Male	1	1	1
Female	1.2 (0.7–2.2)	1.0 (0.5–2.1)	1.3 (1.0–1.7)
p value	0.5	0.9	0.08
Age group, y			
0–9	0.1 (0.05–0.4)	0.1 (0.04–0.3)	
10–17	0.2 (0.07–0.4)	0.3 (0.1–0.7)	
18–34	0.5 (0.2–1.2)	0.4 (0.1–0.9)	0.9 (0.7–1.3)
≥35	1	1	1
p value	<0.001	<0.001	0.6
HSV-2			
Negative	1	1	1
Positive	0.3 (0.1–0.9)	2.7 (1.2–6.5)	1.3 (1.0–1.6)
p value	0.03	0.02	0.09

*Seroreactivity by any serologic assay (whole virus ELISA, IFA-LANA, or IFA-lytic) in multivariable analysis. KSHV, Kaposi sarcoma–associated herpesvirus; IFA-LANA, immunofluorescence assay that detected latent-associated nuclear antigens; IFA-lytic, IFA that detected lytic-associated nuclear antigens; aOR, age- and sex-adjusted odds ratio; CI, confidence interval; HSV-2, herpes simplex virus type-2.

Department of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK (LSHTM).

The study in blood banks was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico, an agency of the Brazilian Ministry of Science and Technology (grants 304879/2003-7, 305258/2006-0 and 473867/2006-0). The study among Mapuera populations was supported by grants from the Wellcome Trust (grant 075454/B/04/Z), CNPq 300317/97-2, and Fundação Faculdade de Medicina, University of São Paulo, Brazil. Additional financial support was provided by the United Kingdom Department for International Development–funded Research Programme Consortium on Research and Capacity Building on Sexual and Reproductive Health and HIV in Developing Countries of the LSHTM.

Dr Nascimento is a research associate scientist in the Laboratory of Virology, Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, Brazil. Her main interests are in the epidemiology and molecular epidemiology of viral infections that cause cancer, in particular, the serologic diagnosis and epidemiology of Kaposi sarcoma–associated herpesvirus infection in various populations in Brazil.

References

- Boshoff C, Weiss RA. Epidemiology and pathogenesis of Kaposi's sarcoma–associated herpesvirus. *Philos Trans R Soc Lond B Biol Sci.* 2001;356:517–34. DOI: 10.1098/rstb.2000.0778
- Newton R, Ziegler J, Bourbouli D, Casabonne D, Beral V, Mbidde E, et al. The sero-epidemiology of Kaposi's sarcoma–associated herpesvirus (KSHV/HHV-8) in adults with cancer in Uganda. *Int J Cancer.* 2003;103:226–32. DOI: 10.1002/ijc.10817
- Newton R, Ziegler J, Bourbouli D, Casabonne D, Beral V, Mbidde E, et al. Infection with Kaposi's sarcoma–associated herpesvirus (KSHV) and human immunodeficiency virus (HIV) in relation to the risk and clinical presentation of Kaposi's sarcoma in Uganda. *Br J Cancer.* 2003;89:502–4. DOI: 10.1038/sj.bjc.6601113
- de Souza VA, Sumita LM, Nascimento MC, Oliveira J, Maschertti M, Quiroga M, et al. Human herpesvirus-8 infection and oral shedding in Amerindian and non-Amerindian populations in the Brazilian Amazon region. *J Infect Dis.* 2007;196:844–52. DOI: 10.1086/520549
- Cunha AM, Caterino-de-Araujo A, Costa SC, Santos-Fortuna E, Boa-Sorte NC, Goncalves MS, et al. Increasing seroprevalence of human herpesvirus 8 (HHV-8) with age confirms HHV-8 endemicity in Amazon Amerindians from Brazil. *J Gen Virol.* 2005;86:2433–7. DOI: 10.1099/vir.0.81087-0
- Whitby D, Marshall VA, Bagni RK, Wang CD, Gamache CJ, Guzman JR, et al. Genotypic characterization of Kaposi's sarcoma–associated herpesvirus in asymptomatic infected subjects from isolated populations. *J Gen Virol.* 2004;85:155–63. DOI: 10.1099/vir.0.19465-0
- Mohanna S, Maco V, Bravo F, Gotuzzo E. Epidemiology and clinical characteristics of classic Kaposi's sarcoma, seroprevalence, and variants of human herpesvirus 8 in South America: a critical review of an old disease. *Int J Infect Dis.* 2005;9:239–50. DOI: 10.1016/j.ijid.2005.02.004
- Plancoulaine S, Abel L, van Beveren M, Tregouet DA, Joubert M, Torreyvoye P, et al. Human herpesvirus 8 transmission from mother to child and between siblings in an endemic population. *Lancet.* 2000;356:1062–5. DOI: 10.1016/S0140-6736(00)02729-X
- Bourbouli D, Whitby D, Boshoff C, Newton R, Beral V, Carrara H, et al. Serologic evidence for mother-to-child transmission of Kaposi sarcoma–associated herpesvirus infection. *JAMA.* 1998;280:31–2. DOI: 10.1001/jama.280.1.31-a
- Engels EA, Atkinson JO, Graubard BI, McQuillan GM, Gamache C, Mbisa G, et al. Risk factors for human herpesvirus 8 infection among adults in the United States and evidence for sexual transmission. *J Infect Dis.* 2007;196:199–207. DOI: 10.1086/518791
- Nascimento MC, de Souza VA, Sumita LM, Freire W, Weiss HA, Sabino EC, et al. Kaposi's sarcoma–associated herpesvirus (KSHV) infection among blood donors in Brazil: a multi-centre serosurvey. *J Med Virol.* 2008;80:1202–10. DOI: 10.1002/jmv.21188
- Nascimento MC, de Souza VA, Sumita LM, Freire W, Munoz F, Kim J, et al. Comparative study of Kaposi's sarcoma–associated herpesvirus serological assays using clinically and serologically defined reference standards and latent class analysis. *J Clin Microbiol.* 2007;45:715–20. DOI: 10.1128/JCM.01264-06
- Nascimento MC, Ferreira S, Sabino E, Hamilton I, Parry J, Panuti CS, et al. Performance of the HerpeSelect (Focus) and Kalon enzyme-linked immunosorbent assays for detection of antibodies against herpes simplex virus type 2 using monoclonal antibody-

blocking enzyme immunoassay (Mab-EIA) and clinico-virological reference standards in Brazil. *J Clin Microbiol.* 2007;45:2309–11. DOI: 10.1128/JCM.00144-07

14. Contas Regionais do Brasil. [cited 2009 Jan 15]. Instituto Brasileiro de Geografia e Estatística. Available from <http://www.ibge.gov.br>
15. Plancoulaine S, Gessain A, van Beveren M, Tortevoye P, Abel L. Evidence for a recessive major gene predisposing to human herpesvirus 8 (HHV-8) infection in a population in which HHV-8 is endemic. *J Infect Dis.* 2003;187:1944–50. DOI: 10.1086/375345

Address for correspondence: Maria C. Nascimento, Laboratory of Virology, Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, Av Dr Eneas de Carvalho Aguiar 470, 05403 000, São Paulo, SP, Brazil; email: maria.nascimento@lshtm.ac.uk

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the U.S. Department of Health and Human Services.

EMERGING INFECTIOUS DISEASES

A Peer-Reviewed Journal Tracking and Analyzing Disease Trends

Vol.11, No.5, May 2005



Search
past issues

EID
online
www.cdc.gov/eid

Persistent Reemergence of Dengue