

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Carter, V; Shimizu, S; Arai, M; Dessens, JT; (2008) PbSR is synthesized in macrogametocytes and involved in formation of the malaria crystalloids. *Molecular microbiology*, 68 (6). pp. 1560-9. ISSN 0950-382X DOI: <https://doi.org/10.1111/j.1365-2958.2008.06254.x>

Downloaded from: <http://researchonline.lshtm.ac.uk/7691/>

DOI: <https://doi.org/10.1111/j.1365-2958.2008.06254.x>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Fig. S1 Analysis of a second independent clone of parasite line Δ SRCR/EGFP. **A:** GFP fluorescence in an ookinete, showing lack of focal spots corresponding to crystalloids. **B:** Molecular analysis by PCR. Lane 1: 1kb product obtained with primers DHFRCAS3' (TCGTGGGCTACGTCCCGCAC) and SR3'-R (CGCCTTCACGCTGATGT), confirming genomic integration of the selectable marker gene cassette in the *pbsr* locus. Lane 2: 2.2kb product obtained with primers SR1250 (CGGAATTTTCGATTATAGAAGG) and pDNR-PbSR-R (ATGAGGGCCCCTAAGCTTAAGCGTTTCAAAAAGGTAAATGA), confirming the presence of the modified *pbsr* locus (missing the sequence encoding the SRCR domains) and the absence of the wild-type *pbsr* locus (would give band of 2.9kb). Lane M: DNA size markers.

