

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



LSHTM Research Online

Weeks, EN; Córdón-Rosales, C; Davies, C; Gezan, S; Yeo, M; Cameron, MM; (2013) Risk factors for domestic infestation by the Chagas disease vector, *Triatoma dimidiata* in Chiquimula, Guatemala. *Bulletin of entomological research*, 103 (6). pp. 634-43. ISSN 0007-4853 DOI: <https://doi.org/10.1017/S000748531300014X>

Downloaded from: <http://researchonline.lshtm.ac.uk/790307/>

DOI: <https://doi.org/10.1017/S000748531300014X>

**Usage Guidelines:**

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: Copyright the publishers

<https://researchonline.lshtm.ac.uk>

# Risk factors for domestic infestation by the Chagas disease vector, *Triatoma dimidiata* in Chiquimula, Guatemala

E. N. I. Weeks<sup>1\*†</sup>, C. Córdón-Rosales<sup>2</sup>, C. Davies<sup>1†</sup>,  
S. Gezan<sup>3</sup>, M. Yeo<sup>1</sup> and M. M. Cameron<sup>1</sup>

<sup>1</sup>Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, U.K.; <sup>2</sup>Center for Health Studies, Universidad del Valle de Guatemala, Guatemala; <sup>3</sup>School of Forest Resources and Conservation, University of Florida, Gainesville, Florida, U.S.A.

## Abstract

In Guatemala prior to control initiatives, the main vectors of *Trypanosoma cruzi*, the causative agent of Chagas disease, were *Rhodnius prolixus* and *Triatoma dimidiata*. This study conducted in 2006 in the department of Chiquimula recorded a high level of *T. dimidiata* infestation and an absence of *R. prolixus* in all surveyed communities. In Guatemala, the presence of *T. dimidiata* as domestic, peridomestic and sylvatic populations results in control difficulties as houses are re-infested from the surrounding environment. Entomological surveys, the current method used to select houses in need of control efforts, are labour intensive and time consuming. A time- and cost-effective way to prioritize houses for evaluation and subsequent treatment is the stratification of houses based on the risk of triatomine infestation. In the present study, 17 anthropogenic risk factors were evaluated for associations with house infestation of *T. dimidiata* including: wall, floor and roof type. There was an increased likelihood of domestic infestation with *T. dimidiata* associated with the presence of dirt floors (18/29; OR 8.075, 95% CI 2.13–30.6), uncoated bajareque walls (12/17; OR 4.80, 95% CI 1.35–17.1) and triatomine-like faeces on walls (16/26; OR 3.89, 95% CI 1.19–12.7). These factors could be used to target control of *T. dimidiata* to communities with an increased risk of being infested.

**Key words:** Triatominae, *Trypanosoma cruzi*, infestation, surveillance, floor, housing

(Accepted 9 March 2013)

## Introduction

Chagas disease, caused by the parasitic protozoan *Trypanosoma cruzi*, is responsible for a larger disease burden

in Latin America than all other vector-borne pathogens combined (WHO, 2004). It is estimated that in Guatemala two million people are at risk from Chagas disease and more than 2000 people become infected with *T. cruzi* every year (PAHO, 2005). Insect vectors, haematophagous reduviid bugs, comprise the main route for *T. cruzi* transmission accounting for more than 80% of new cases (PAHO, 2005). In Guatemala, prior to the initiation of the control programme, the main vectors were *Rhodnius prolixus* and *Triatoma dimidiata* (Ponce, 2007). *Rhodnius prolixus* was focalized in a well-defined region in eastern Guatemala, and was always found to be associated with human houses (WHO, 2002). *Triatoma dimidiata*, in contrast was more widely distributed and found in domestic,

---

\*Author for correspondence

Phone: +1 352 273 3954

Fax: +1 352 392 5660

E-mail: eniweeks@ufl.edu

†Deceased.

‡Present address: Entomology and Nematology Department, University of Florida, Gainesville, Florida, U.S.A.

peridomestic and sylvatic environments (WHO, 2002). Consequently, different approaches were launched to control the two species; complete elimination of *R. prolixus*, and domiciliary control of *T. dimidiata* (<5% infestation rate). Elimination of *T. dimidiata* was not believed to be possible due to the risk of house re-invasion from extra-domiciliary populations with potential for subsequent reproduction inside houses. In addition to vector control measures, the Central American Initiative for Chagas Disease Control (IPCA) also ordered the screening of all blood donors (WHO, 2002). Extensive implementation of house spraying with residual insecticides in Guatemala, has placed the country ahead in the goal to eliminate *R. prolixus* (PAHO, 2006). Reports suggest that following the intensive vector control campaign in the last decade *R. prolixus* has been virtually eliminated (Petherick, 2010; King *et al.*, 2011). Although, intradomestic infestations of *T. dimidiata* have been greatly reduced, there are persistent foci in several departments. In Guatemala, 39% of sampled *T. dimidiata* were infected with *T. cruzi*, and vectorial transmission has been demonstrated even when the insect is present at low infestation densities (Schofield & Dujardin, 1997; Monroy *et al.*, 1998a, b; PAHO, 2006). Consequently, it is now considered the most important vector of Chagas disease in Guatemala.

*Triatoma dimidiata* is native and widespread throughout the country in extradomestic habitats and so its elimination is virtually impossible. Sustained monitoring and cost-effective targeting of control will be necessary to maintain houses free of infestation. Entomological surveys, the current method used to monitor for domestic triatomines, are labour intensive and time consuming. Therefore, in countries where Chagas disease is endemic, entomological evaluations are targeted at communities that are believed to be at increased risk of triatomine infestation. Risk is evaluated by quantifying factors that influence the probability of domestic infestation with triatomine bugs e.g., houses possessing mud walls and palm roofs, the results of anecdotal surveys, suspected infestations or houses previously infested (Nakagawa *et al.*, 2003; Campbell-Lendrum *et al.*, 2007; King *et al.*, 2011). However, targeted sampling, using risk factors, was recently found to be less effective than random sampling at detecting villages with *T. dimidiata* prevalence above the 5% threshold (King *et al.*, 2011). The failure of this method was believed to be due to the assumption of a similarity in risk factors across geographic space. However, *T. dimidiata* prevalence is patchy and appears to be dependent upon different risk factors and environmental variables in different areas (King *et al.*, 2011). In addition, one of the risk factors used for targeting sampling was the presence of a palm roof, a characteristic that is highly correlated with *R. prolixus* infestation (Cordon-Rosales & Pennington, 2007), but *T. dimidiata* is rarely associated with palm roofing materials. Therefore, risk factors specific to *T. dimidiata* need to be identified to increase the effectiveness of control based on priority stratification of communities and houses. In domestic habitats, *T. dimidiata* are often found within cracks, in the floor or walls, concentrated around the sleeping areas (Zeledon *et al.*, 1973; Schofield & Dujardin, 1997; Monroy *et al.*, 1998a, b; WHO, 2002). However, *T. dimidiata* exhibits a preference for peridomestic refuges, such as piles of roof tiles or firewood (Zeledon *et al.*, 1973; Starr *et al.*, 1991). Consequently, despite living outside the house, the insects are close enough to enter houses and feed on human hosts.

Differences between geographic regions, such as variation in vector distribution due to environmental factors, e.g.,

temperature, humidity and vegetation type, also have an effect on risk. For example, the risk of *T. dimidiata* infestation is elevated in certain life zones, in particular subtropical forests (Cordon-Rosales & Pennington, 2007). To stratify an area based on risk of triatomine infestation, locally important variables need to be established. Risk factors may vary spatially between regions due to variation in human behaviour, vector behaviour and ecology and environmental factors. Local knowledge can then be used to more effectively target houses for vector control and also to determine house characteristics that could be prioritized in house improvement programmes (Campbell-Lendrum *et al.*, 2007).

The present study aimed to identify factors associated with domestic infestation in Chiquimula, Guatemala, a department with high levels of triatomine infestation and infection with *T. cruzi* (Tabaru *et al.*, 1999a, b; Monroy *et al.*, 2003). More specifically, variables could be used to stratify houses for vector control and prioritize improvements in housing to reduce suitability for *T. dimidiata* infestation.

## Materials and methods

### Study sites

The survey was performed from June to August 2006 in the eastern highlands in the Chagas disease endemic department of Chiquimula, Guatemala (fig. 1). Chiquimula is at a latitude of 14°48'00"N and a longitude of 89°33'00"W, with an elevation of 420 m above sea level.

A total of 50 houses were sampled representing six communities from two municipalities. Chancó, Corral de Piedra, Río Arriba, Los Encuentros in San Juan Ermita (654 m.a.s.l.), and Pedregalito and El Limón in Quezaltepeque (685 m.a.s.l.). The communities were chosen based on infestation data and intervention history collected by the Universidad del Valle de Guatemala. All study communities had baseline infestation rates greater than 15% and had no previous record of control intervention. Within each community, a probabilistic sample of houses was selected by systematically choosing every fifth house from a census map of the community, provided by the national vector control programme. The number of houses to be sampled per community was estimated using census data provided by the Guatemalan Ministry of Health (Ministerio de Salud Pública y Asistencia Social-MSPAS) with the following parameters: 30% estimated infestation level, 5% precision, 95% CI and 80% statistical power (Hashimoto *et al.*, 2006). Therefore, if the total number of houses was 0–30 then 50% were sampled, 33% for 31–59 houses and 10% for 60–100 houses.

### Risk factors

Householders were shown photographs and pinned specimens of *R. prolixus* and *T. dimidiata* and asked about triatomine activity using four questions; (1) Have you seen a triatomine bug in the last month?, (2) Was it inside or outside your house?, (3) Have you been bitten by a triatomine bug in the last month?, and (4) Have you sprayed insecticide to try and kill triatomine bugs or other insects? Information was recorded about the house structure and potential refuges in the domestic environment and these data formed the basis for the 17 risk-factor measures (table 1). With regard to the more subjective variables: organization and lighting, categories

Table 1. House characteristics by community in the municipalities of San Juan Ermita and Quezaltepeque, Chiquimula, Guatemala.

Risk factor			Chancó	Corral de Piedra	Río Arriba	Los Encuentros	Pedregalito	El Limón
			Percentage of houses with characteristics					
House construction	Walls	Bajareque uncoated	42.86	37.50	87.50	37.50	8.33	0.00
		Bajareque coated	57.14	62.50	12.50	0.00	0.00	14.29
		Adobe uncoated	0.00	0.00	0.00	37.50	50.00	57.14
		Adobe coated	0.00	0.00	0.00	25.00	41.67	28.57
	Roof	Tile	28.57	0.00	0.00	0.00	41.67	28.57
		Palm	28.57	37.50	62.50	12.50	0.00	0.00
		Metal	42.86	62.50	37.50	87.50	58.33	71.43
	Floor	Earth/Dirt	42.86	62.50	87.50	87.50	33.33	42.86
		Concrete or Tile	57.14	37.50	12.50	12.50	66.67	57.14
	Refuges	Cracks	Present	57.14	37.50	87.50	87.50	58.33
Boxes			Present	57.14	75.00	100.00	75.00	58.33
Shrine		Present	100.00	50.00	12.50	12.50	16.67	14.29
		Pictures	Present	85.71	62.50	25.00	62.50	75.00
Plastic		Present	42.86	25.00	0.00	12.50	33.33	0.00
		Sacks	Present	57.14	50.00	100.00	50.00	58.33
Clothes		Present	85.71	75.00	100.00	37.50	91.67	100.00
		Wood	Present	14.29	37.50	87.50	25.00	0.00
Chicken nests		Present	14.29	0.00	12.50	0.00	8.33	14.29
		Overall house organization <sup>1</sup>	Disorganized	71.43	50.00	75.00	50.00	58.33
Other	Lighting <sup>1</sup>	Dark	85.71	75.00	100.00	75.00	58.33	28.57
	Insecticide	Not used	57.14	75.00	37.50	37.50	58.33	57.14
	Triatomine-like faeces	Present	71.43	25.00	75.00	50.00	50.00	42.86

<sup>1</sup> Categorized by visual inspection (interviewer). Organized=belongings tidied away or intentionally positioned. Disorganized=no obvious intentional positioning of belongings, with items such as food, clothing or firewood, on the floor providing potential triatomine refuges. Light=natural light in bedroom, Dark=no natural light in bed room.



Fig. 1. Maps of Guatemala, showing the department of Chiquimula (A) and the sampled municipalities in Chiquimula (B), San Juan Ermita and Quezaltepeque, with the sample houses marked by circles (black).

were assigned after visual inspection (E. Weeks). In 'organized' houses belongings were tidied away or intentionally positioned, whereas in 'disorganized' houses there was no obvious intentional positioning of belongings, with items such as food, clothing or firewood, on the floor providing potential triatomine refuges. Lighting was classified based on the presence of natural light during the day, sleeping areas with

natural light entering through windows or doors were 'light' and those with no natural light were 'dark'. Only natural light was considered as only two of the six communities had an electricity supply.

A global positioning system (GPS) device was used to record the latitude, longitude and altitude of each sample house.

### Triatomine detection methods

A combination of three detection methods was used during the study to identify properties infested with triatomines. These included a passive sensor box method and two active methods. Active methods were householder collection and manual inspection by trained personnel. All three methods were applied to each house. Sensor boxes were installed and householder collection pots were distributed at the beginning of the study. After three weeks the pots and boxes were collected and a manual inspection was completed. The sensor boxes were checked for any signs of triatomine infestation: faeces, eggs, exuviae, nymphs or adults. All triatomines collected were classified morphologically into species using identification keys (Lent & Wygodzinsky, 1979). Sensor boxes were of the Gomez Nunez design (Gomez Nunez, 1965), which were made of cardboard and measured 31 × 25 × 5 cm. On one side of the box, positioned in contact with the wall, there were 12 holes; each 2 cm in diameter. Inside each box, stiff brown paper sheet was folded and perforated with holes to form a lattice-like structure for the triatomines to rest on. Two sensor boxes were positioned in the principal bedroom, on the wall nearest the bed, one each at 50 and 150 cm. The householder collection pots were adapted 50 ml conical tubes; the lid of the tube was perforated to hold a piece of mesh in place for ventilation. Inside the tube was a piece of folded brown resting paper. Protective gloves were provided for use when catching triatomines and the use of the pot was demonstrated. A manual inspection of the domestic area was conducted for 15 min or 0.5 man hours using forceps and flashlights (Nakagawa *et al.*, 2003). A 15 min search of any potential peridomestic refuges, e.g., wood piles, chicken coops, within the house boundaries (maximum distance 50 m) followed the intradomestic search. The distance constraint was set within 50 m or the change to farmland or sylvatic areas, whichever was met first.

### Data analyses

Triatomine distribution was measured using four conventional entomological indices at the department, municipality and community level (WHO, 2002; Nakagawa *et al.*, 2005). The infestation, colonization and crowding index were defined as the proportion of houses that have *T. dimidiata*, that have immature stages and the number of triatomines per infested property, respectively. The infestation density was defined as the number of triatomines per sample house.

Risk factor analyses were performed in STATA (StataCorp, TX) using multivariate stepwise logistic regression with the dependant variable being houses found infested with *T. dimidiata* by any method. Independent variables, or risk factors, were interpreted as categorical variables. The maximal model included all variables with a z-test *P* value of less than 0.2 from the univariate analyses. Variables with a z-test *P* value of greater than 0.05 were removed in a stepwise method from the maximal model until only variables significant by the z-test remained. Finally, each variable that had been excluded was added back in to the model individually, to confirm that it added no explanatory power to the model. Variables with more than two categories were examined for association with the likelihood ratio (LR) test in STATA. In addition, a logistic regression was also performed to test the significance of geographical variables, i.e., altitude, latitude and longitude,

over the presence of *T. dimidiata* infestation using GenStat (VSN International, U.K.).

### Ethics

The study was reviewed and approval was granted by the Research Ethics Committee at the London School of Hygiene and Tropical Medicine, U.K. The first visit to each house began with an explanation of the study to gain verbal consent from the inhabitants.

## Results

### Infestation

A total of 130 *T. dimidiata* and one *Triatoma nitidia* were collected. No samples of *R. prolixus* were collected by any of the methods. Of the *T. dimidiata* collected, 50 specimens were adults and 80 were nymphs (table 2). The single *T. nitidia* specimen, collected in Los Encuentros (San Juan Ermita), was a nymph; the house was not co-infested with *T. dimidiata*. In all houses where nymphs were found, adults were also collected (table 2; fig. 2). In Chiquimula the *T. dimidiata* infestation index was found to be 44% (22/50; 95% CI 30.3–58.6), with the two municipalities surveyed, San Juan Ermita and Quezaltepeque, having indices of 52% (16/31; 95% CI 33–70) and 32% (6/19; 95% CI 9–54), respectively. All surveyed communities had greater than 15% infestation with *T. dimidiata*. The average number of *T. dimidiata* collected per infested house in Chiquimula was 6 (±4.6 s.d.; crowding index) and the average number of *T. dimidiata* collected per sample house was 2.6 (±4.22 s.d.; infestation density). The infestation density and crowding indices by municipality were similar. The highest number of triatomines collected from one house was 17 specimens, consisting of six adults and 11 nymphs, in Chancó (San Juan Ermita).

### Risk factor analyses

Risk factor analyses were performed to identify variables associated with *T. dimidiata* infestation (table 3; the single house positive for *T. nitidia* was excluded). The probability that a house was infested with *T. dimidiata* increased significantly with the presence of uncoated bajareque walls (*P*=0.015; OR 0.66, 95% CI 0.18–2.40) and dirt floors (*P*=0.002; OR 8.08, 95% CI 2.13–30.60). Furthermore, the presence of triatomine-like faeces on interior walls, was associated with an increase in infestation rate (16/26, 62% infested; *P*=0.025; OR 3.89, 95% CI 1.19–12.70). Although only 17 of the 50 houses sampled possessed uncoated bajareque walls, 71% (12/17) of these houses were infested with *T. dimidiata*. The largest odds ratio was associated with floor type, signifying a strong association of this variable with the probability of a house being infested by *T. dimidiata*. Of the houses with dirt floors, 66% (18/29) were infested with *T. dimidiata*.

The analysis produced a maximal model with nine risk factors ( $\chi^2=13.60$ ; *P*=0.0587). The presence of uncoated bajareque walls, a dirt floor, cracks in the walls (*P*=0.182; OR=2.27, 95% CI=0.068–7.53), firewood (*P*=0.197; OR=2.35, 95% CI=0.64–8.58) and triatomine-like faeces increased the probability of an infestation. Disorganized (*P*=0.053; OR=3.81, 95% CI=0.99–11.10) and dark houses (*P*=0.154; OR=2.67, 95% CI=0.69–10.30) were also more likely to be infested with *T. dimidiata*. The presence of coated adobe walls

Table 2. *Triatoma dimidiata* captured by community.

Department	Municipality	Community	Houses sampled (total houses in community)	Total triatomines collected	Total adults collected	Total nymphs collected	Infestation index % (95% CI)	Colonization index % (95% CI)	Infestation density (SD)	Crowding index (SD)
Chiquimula	San Juan Ermita		50	130	50	80	44 (30–59)	36 (22–50)	2.60 (4.22)	5.91 (4.61)
		Chancó	31	93	37	56	52 (33–70)	45 (27–63)	3.00 (4.45)	5.81 (4.71)
		Corral de Piedra	7 (67)	31	15	16	57	43	6.57	7.75
		Río Arriba	8 (77)	6	2	4	25	13	1.00	7.8
		Los Encuentros	8 (81)	39	14	25	63	63	6.63	3.4
	Quezaltepeque		8 (83)	17	6	11	63	63	2.88	4.60
		Pedregalito	19	37	13	24	32 (9–54)	21 (1–41)	1.95 (3.87)	6.17 (4.75)
		El Limón	12 (45)	21	6	15	17	17	2.25	10.5
			7 (91)	16	7	9	57	29	2.86	4

( $P=0.130$ ; OR=0.272, 95% CI=0.050–1.47) and plastic on the walls ( $P=0.168$ ; OR=0.356, 95% CI=0.082–1.55) decreased the likelihood that a house was found to be infested. Multivariate stepwise logistic regression analyses produced a minimal model where only the presence of a dirt floor remained significant ( $\chi^2=13.36$ ;  $P=0.0013$ ). In comparison with any other type of floor (concrete or tile), dirt floors increased the probability that a house was infested with *T. dimidiata*.

Analysis of the geographical variables by logistic regression revealed a significant linear relationship of altitude with the presence of *T. dimidiata* infestation (fig. 3;  $P=0.027$ ; range 791–1101 m.a.s.l.). Within this range, every 100 m increase in altitude resulted in an increase in the community infestation rate by 25%. No significant relationship was found between *T. dimidiata* infestation and the latitude ( $P=0.116$ ) or longitude ( $P=0.675$ ) of the sample houses.

## Discussion

Historically, Chiquimula was an area with a high level of domestic triatomine infestation (Tabaru *et al.*, 1999a, b). In a previous survey completed in 1999, more than 25% of houses within the department were infested with *T. dimidiata* or *R. prolixus* (Tabaru *et al.*, 1999a, b). All six of the study communities in the municipalities of San Juan Ermita and Quezaltepeque (Chiquimula, Guatemala) were infested with and colonized by triatomines. The majority of the triatomines collected were identified as *T. dimidiata*. Previous studies in Guatemala found that at a *T. dimidiata* infestation rate of approximately 25% (in the absence of other vectors) there was a 8.9% *T. cruzi* seropositivity among inhabitants (Paz-Bailey *et al.*, 2002). Our study indicated that in five of the six communities that were sampled the infestation rate was equal to or greater than this level. Chiquimula was, therefore, in urgent need of control of *T. dimidiata*.

The use of house characteristics to stratify communities based on risk is one method of targeting control on those houses that are most likely to be infested. In the current study, the presence of statistically significant associations between the probability of infestation by *T. dimidiata* and several anthropogenic characteristics has been demonstrated. The probability of infestation increased significantly with the presence of dirt floors, uncoated bajareque walls and triatomine-like faeces, risk factors that are consistent with previous studies and existing knowledge of triatomine ecology. Dirt floors are thought to provide refuge and the opportunity to use dirt for camouflage, a distinctive behaviour of *T. dimidiata* (Zeledon *et al.*, 1969). In a previous study in Guatemala, the majority of houses were found to have dirt floors thus limiting the power to detect a significant association; the results indicated no significant correlation (Cordon-Rosales & Pennington, 2007). In the communities visited in the present study, 21 out of 50 houses had modified concrete or tile floors so it was possible to study the association between floor type and the probability of infestation with *T. dimidiata*. A study in Jutiapa, Guatemala, found that intradomestic prevalence of *T. dimidiata* in villages was more likely to be above the 5% threshold in communities with a higher proportion of houses with dirt floors (King *et al.*, 2011). Communities with a higher proportion of alternative floor types, e.g., tile, had a lower likelihood of village infestation (King *et al.*, 2011). The results are consistent with studies conducted in other countries into the ecology of *T. dimidiata*. For example, in two studies in Costa Rica the




Fig. 2. Infestation maps for municipalities and communities in Chiquimula, Guatemala. Maps show the sampled properties (circles) in Quezaltepeque (A, B, C) and San Juan Ermita (D, E, F), infested properties have filled in circles (black).

species was found to have an association with dirt floors (Zeledon & Vargas, 1984; Starr *et al.*, 1991). Although, Starr *et al.* (1991) did not find a significant association between dirt floors and the risk of being infested with *T. dimidiata*, the factor was assumed to be important due to high odds ratios. Other studies have not detected the relationship with dirt floors. For example, a study in Colombia identified a reduction in triatomine infestation in general with the presence of other floor types, but the species specific analysis found no relationship of *T. dimidiata* infestation and floor type (Campbell-Lendrum *et al.*, 2007). Another study completed in Guatemala found no significant association between *T. dimidiata* infestation and floor type, the authors hypothesized that the result was due to the infestations being well established (Bustamante *et al.*, 2009). The variation between studies completed in different areas highlights the need to understand the risk factors in each epidemiological setting before the factors are used to prioritize houses for control.

The other risk factors identified, including the presence of uncoated bajareque walls and triatomine-like faeces, are known indicators of potential triatomine infestation. *T. dimidiata* is known to live in cracks within walls (Zeledon *et al.*, 1973; WHO, 2002), and previous studies also have recorded this association with unplastered walls (Monroy

*et al.*, 1998a, b; Tabaru *et al.*, 1999a, b; Campbell-Lendrum *et al.*, 2007; Cordon-Rosales & Pennington 2007; Bustamante *et al.*, 2009). As there are several wall dwelling species, the presence of uncoated bajareque walls or the absence of wall coating is likely to be a more general risk factor indicative of triatomines in general. Coated walls were found to protect against several other species of Triatominae including *Triatoma infestans* and *Triatoma pallidipennis* (Enger *et al.*, 2004; Levy *et al.*, 2006). In contrast, dirt floors are likely to be a more specific risk factor for *T. dimidiata* infestation, as dirt camouflage is a behaviour specific to this species. Triatomine-like faeces on walls is often used as indirect evidence of triatomine presence during manual inspection and has been found to be a good indicator of domestic infestation for *T. dimidiata* and *R. prolixus* (Monroy *et al.*, 1998a, b). However, the identification of triatomine-like faeces is not as simple as recording the floor and wall type of a house and will involve a more invasive assessment of the property by an experienced technician.

Following further analysis, only the presence of a dirt floor was included in the minimal model, in this area of Guatemala, this factor alone is likely to be the most reliable indicator of *T. dimidiata* infestation risk. Consequently, we propose the use of the presence of dirt floors and uncoated bajareque walls as factors to stratify houses in Chiquimula, Guatemala based on for risk of *T. dimidiata* infestation. At the same time, it would be

Table 3. Univariate analysis of house-level risk factors, effect on the probability that a house was found infested by *T. dimidiata*.

Risk factor		<i>n</i> <sup>1</sup>	% positive <sup>2</sup>	OR (95% CI)	Z	P-value	
House construction	Walls	Bajareque uncoated	17	71	4.80 (1.35–17.10)	2.42	0.015 <sup>3</sup>
		Bajareque coated	11	36	0.60 (0.15–2.39)	–0.72	0.470
		Adobe uncoated	13	39	0.66 (0.18–2.40)	–0.63	0.527
	Roof	Adobe coated	9	22	0.27 (0.05–1.47)	–1.51	0.130 <sup>3</sup>
		Tile	9	33	0.43 (0.01–1.90)	–1.12	0.264
		Palm	11	46	0.97 (0.25–3.73)	–0.04	0.967
	Floor	Metal	30	50	1.74 (0.56–5.46)	0.95	0.342
		Dirt	29	66	8.08 (2.13–30.60)	3.07	0.002 <sup>3</sup>
		Concrete or Tile	21	19	0.15 (0.04–0.55)	n/a	n/a
Refuges	Cracks	Present	31	55	2.27 (0.68–7.53)	1.34	0.182 <sup>3</sup>
		Boxes	36	47	1.19 (0.34–4.14)	0.28	0.781
	Shrine	Present	16	56	1.84 (0.55–6.10)	0.99	0.321
		Pictures	33	46	0.94 (0.29–3.03)	–0.11	0.914
	Plastic	Present	10	30	0.36 (0.08–1.55)	–1.38	0.168 <sup>3</sup>
		Sacks	31	48	1.50 (0.48–4.72)	0.69	0.488
	Clothes	Present	41	46	1.36 (0.33–5.55)	0.42	0.671
		Wood	13	62	2.35 (0.64–8.58)	1.29	0.197 <sup>3</sup>
	Chicken nests	Present	4	75	3.90 (0.38–40.40)	1.14	0.254
Overall house organization <sup>4</sup>		Disorganized	28	61	3.81 (0.99–11.10)	1.94	0.053 <sup>3</sup>
Other	Natural illumination <sup>5</sup>	Dark	35	54	2.67 (0.69–10.30)	1.42	0.154 <sup>3</sup>
	Insecticide	Not used	23	48	1.25 (0.40–3.86)	0.39	0.390
	Triatomine-like faeces	Present	26	62	3.89 (1.19–12.70)	2.25	0.025 <sup>3</sup>

<sup>1</sup> Number of houses out of 50 where the factor was present.

<sup>2</sup> Percentage of houses with factors that were positive for *T. dimidiata* (house positive for *T. nitidia* excluded from the analysis).

<sup>3</sup> Indicates inclusion in the maximal model.

<sup>4</sup> Categorized by visual inspection (interviewer). Organized=belongings tidied away or intentionally positioned. Disorganized=no obvious intentional positioning of belongings, with items such as food, clothing or firewood, on the floor providing potential triatomine refuges.

<sup>5</sup> Categorized by visual inspection (interviewer). Light=natural light in bedroom, Dark=no natural light in bed room.




Fig. 3. Relationship between altitude and infestation rate (%) with *T. dimidiata*. Bars (grey) represent the arithmetic mean altitude in meters above sea level (m.a.s.l.;  $\pm 95\%$  CI) of the sample houses within the six communities, CP=Corral de Piedra, P=Pedregalito, C=Chancó, LE=Los Encuentros, RA=Río Arriba and EL=El Limón. Triangles (black) represent the community infestation rate (%).

necessary to identify the resilient *T. dimidiata* infestation areas to ascertain focalized risk factors.

Coated adobe walls and the presence of plastic on the walls were also included in the maximal model for potentially decreasing the odds of infestation in this study. Coating walls would restrict access to cracks reducing potential refuges, and plastic on the walls may have a similar effect while also making thorough manual inspection more difficult.

The use of plastic material on walls was associated with a low likelihood of infestation (30%), however, the number of houses with this modification was low ( $n=10$ ). If the negative association with plastic is true, then plastic-coated walls could provide an innovative method of controlling *T. dimidiata* infestations, which may be enhanced by the use of insecticide impregnated plastics.

In addition to those factors that were found to significantly increase the probability of infestation in the present study, other variables in the maximal model could be used to modify the risk of *T. dimidiata* infestation, e.g., the presence of cracks in walls and wood piles, disorganization and poor natural illumination. Cracks in walls and wood piles are known refuges of *T. dimidiata* (Zeledon & Vargas, 1984; Dias & Schofield, 1999; WHO, 2002). While there was no significant association detected with any other house content factor in the present study, e.g., boxes, sacks; the majority of homes sampled had uncoated walls (56%) with cracks (60%), so there was no shortage of preferred *T. dimidiata* refuges available. A disorganized, poorly illuminated house could potentially have a greater number of refuges for triatomines and permit an extended period of activity for these nocturnal insects. Previous studies have recorded a negative association of *T. dimidiata* with interdomestic light intensity (Tabaru *et al.*, 1999a, b), and number of windows in the bedroom (Bustamante *et al.*, 2009). Several studies have found an association with the presence of domestic animals and *T. dimidiata* and other Chagas disease vectors (Enger *et al.*, 2004; Levy *et al.*, 2006; Bustamante *et al.*, 2009). During the current study, few animals were observed within the sample properties. Only four of the 50 houses sampled had any evidence of animals living indoors, this was always in the form



of indoor chicken nests and of these four properties three were infested with *T. dimidiata*.

While the identification of house-level risk factors for infestation would decrease the cost of control, the community would need to be visited at least once in order for these risk factors to be recorded. However, unlike entomological evaluation, this assessment could be done by a non-specialist or someone within the community. In addition, it would be desirable to test if remotely sensed environmental variables could help stratify the risk of triatomine infestation in large areas. In Guatemala, *T. dimidiata* can be found throughout a range, from sea level to 1600 m and has been found to be most prevalent between 800 and 1000 m.a.s.l. (Tabaru *et al.*, 1999a, b). Our study describes a relationship of increasing infestation with increasing altitude within this range in Chiquimula. However, the relationship may vary across the county so nationwide studies are needed to determine how altitude affects the infestation rate by department across the whole of Guatemala. Variation in infestation due to changes in altitude is unlikely to be independent of other environmental and climatic variables. Future work could incorporate such factors to identify zones of increased risk within the country. As environmental variables can be determined remotely, such a method for assessing risk would be probably more cost effective than surveys, enabling at risk houses to be predetermined without any field visits.

The identification of house-level risk factors may also be used to inform house improvement schemes of the factors most likely to reduce triatomine infestation. Home improvements have the potential to greatly reduce the risk of triatomine infestation by reducing the availability of refuges for resting and breeding. Campbell-Lendrum *et al.* (2007) predicted a 60% decrease in risk of infestation by triatomines associated with improvements in basic house structure. The model used assumed the complete renovation of all houses within a village through plastering of walls, cementing of dirt floors and replacement of natural roofing materials such as palm or thatch, with metal or tile (Campbell-Lendrum *et al.*, 2007). Costs associated with complete renovation would exceed spraying costs in the short term. For example, a study in Paraguay priced a renovation at US\$600 and a round of insecticide fumigation at US\$29 (Rojas de Arias *et al.*, 1999). However, the results of house improvements are longer lasting and have other benefits due to the increased level of construction and hygiene. Owing to the costs associated with a complete house renovation it is important to prioritize changes in house structure based on locally important risk factors. For example, following the identification of uncoated walls as a risk factor for *T. dimidiata* infestation in Jutiapa, Guatemala (Bustamante *et al.*, 2009), a trial house improvement scheme was established involving wall plastering with traditional materials and methods by inhabitants (Monroy *et al.*, 2009). The single improvement, costing just US\$30 per house, reduced domestic triatomine infestation in all communities equal to a traditional spraying programme but with longer lasting effects (Monroy *et al.*, 2009).

The current study also found uncoated walls to increase the risk of *T. dimidiata* infestation, although the risk factor with the strongest association was found to be the presence of a dirt floor. A survey of school age children completed in Guatemala found a significant association with seropositivity for *T. cruzi* and floor type (Rizzo *et al.*, 2003). Children living in a house with a modified floor had a lower seropositivity rate (3.9%), compared with those children living in houses with traditional

dirt floors (6.5%) (Rizzo *et al.*, 2003). Therefore, the modification of dirt floors could decrease *T. cruzi* transmission through long-term reduction in contact between the insect and the human host. Currently, 42% of houses in Chiquimula have a cement floor so it is a socially acceptable flooring material and would be feasible to implement providing costs were subsidized. The replacement of dirt floors with those made of concrete or tile could have an impact on the transmission of other diseases, for example, tungiasis caused by *Tunga penetrans* (sandflea). Unsurprisingly in areas where tungiasis occurs, there is an association with people that are heavily infested and those who live in houses with dirt floors (Muehlen *et al.*, 2006).

The presence of *T. dimidiata* in extradomiciliary populations is believed to be the main factor that will prevent elimination of the species within its natural range. Following treatment, peridomestic and sylvatic reservoirs of triatomines are able to recolonize houses (Schofield, 1994). As the first point of contact during invasion is likely to be the flooring material, the changing of floor construction, from dirt, that is preferred by this species, to a concrete or tile floor may prevent re-colonization of homes (Zeledon & Vargas, 1984).

This study is not without limitations. For example, one limitation of this study is the sample size. While guidelines were followed with regards to sampling for surveillance of triatomines in Guatemala, a total of 50 houses was relatively low for identification of risk factors. Despite this limitation, the study identified several factors that were statistically significant in their association with *T. dimidiata* infestation with clear biological relevance. Extending the study to more houses in communities across Guatemala could improve the power of the study and enable mapping of the risk factors across the country. Not all possible variables were considered in this study. However, the majority of those with a previously demonstrated association with *T. dimidiata* (e.g., uncoated walls) and *R. prolixus* (e.g., palm roof material), the main vectors in Guatemala, were investigated. Furthermore, two of the variables, organization and light, were highly subjective and classified by one person. Future studies could incorporate a standardization of the method or classification to reduce the effect of a personal bias. Finally, the study was completed several years ago, in 2006, since then there have been many changes in Guatemala. Particularly, the great progress in Chagas disease vector control in Guatemala with the first goal of the Central American Initiative, *R. prolixus* elimination, almost achieved. However, the study remains relevant and its findings important, as although a reduction in domestic infestation with *T. dimidiata* has been observed, in accordance with the second goal, there are areas in Guatemala that remain infested with *T. dimidiata* despite repeated treatment by residual insecticide spraying. Not only is surveillance essential to assess the impact of vector control and to target interventions but also epidemiological studies are needed to find alternative ways to further reduce infestation and prevent re-colonization from the surrounding environment.

In conclusion, we identified a high level of *T. dimidiata* colonization in Chiquimula. Efforts now need to be directed towards control of this species, especially as Guatemala was recently declared free of *T. cruzi* transmission by *R. prolixus* (Guhl *et al.*, 2009). Indeed during the current study no samples of *R. prolixus* were collected. Identification of factors associated with an increased risk of *T. dimidiata* infestation that could potentially be used to prioritize houses for monitoring and control efforts, could be highly beneficial on an economic

basis. Previously in Guatemala, entomological surveys were based on risk factors obtained from Costa Rica (Starr *et al.*, 1991). However, factors may vary between countries or even departments. The current study indicates that in Chiquimula, Guatemala, stratifying houses by floor and wall type would enable the targeting of control interventions to houses at increased risk of triatomine infestation. In addition, the implementation of a house improvement scheme involving the covering of dirt floors with concrete or tiles and walls with plaster could decrease triatomine infestation and reduce the risk of re-invasion through the removal of potential refuges.

### Acknowledgements

This publication is dedicated to Professor Clive Davies, the inspiration and driving force behind this study. The authors would like to thank the field technicians Pedro Peralta and Alfonso Salam who worked on this project, for their invaluable skills and knowledge. To acknowledge Haroldo Monterroso and the Oficina de Control de Enfermedades Transmitidas por Vectores (Chiquimula) for their cooperation and for allowing their experienced technicians (Alvaro, Mario and Felipe) to work with us. In addition, we would like to thank Gabriel Jaime Parra Henao, Germán Aguilera Jaramillo, Maria Dora Feliciangeli and Mónica Flórez Martínez for advice before and during the field work. Sincere thanks and admiration to Joel Irish for assistance with geographical data and mapping.

### References

- Bustamante, D.M., Monroy, C., Pineda, S., Rodas, A., Castro, X., Ayala, V., Quinones, J., Moguel, B. & Trampe, R. (2009) Risk factors for intradomiciliary infestation by the Chagas disease vector *Triatoma dimidiata* in Jutiapa, Guatemala. *Cadernos de Saude Publica* **25**, S83–S92.
- Campbell-Lendrum, D.H., Angulo, V.M., Esteban, L., Tarazona, Z., Parra, G.J., Restrepo, M., Restrepo, B.N., Guhl, F., Pinto, N., Aguilera, G., Wilkinson, P. & Davies, C. R. (2007) House-level risk factors for triatomine infestation in Colombia. *International Journal of Epidemiology* **36**, 866–872.
- Cordon-Rosales, C. & Pennington, P.M. (2007) Eco-epidemiología de la transmisión vectorial de la enfermedad de Chagas en Guatemala. *Revista de la Universidad del Valle de Guatemala* **16**, 63–84.
- Dias, J.C.P. & Schofield, C.J. (1999) The evolution of Chagas disease (American Trypanosomiasis) control after 90 years since Carlos Chagas discovery. *Memorias do Instituto Oswaldo Cruz* **94**, 103–121.
- Enger, K.S., Ordonez, R., Wilson, M.L. & Ramsey, J.M. (2004) Evaluation of risk factors for rural infestation by *Triatoma pallidipennis* (Hemiptera:Triatominae), a Mexican vector of Chagas disease. *Journal of Medical Entomology* **41**, 760–767.
- Gomez Nunez, J.C. (1965) Desarrollo de un nuevo metodo para evaluar la infestacion intradomiciliaria por *Rhodnius prolixus*. *Acta Cientifica Venezolana* **16**, 26–31.
- Guhl, F., Pinto, N. & Aguilera, G. (2009) Sylvatic triatominae: a new challenge in vector control transmission. *Memorias do Instituto Oswaldo Cruz* **104**, 71–75.
- Hashimoto, K., Cordon-Rosales, C., Trampe, R. & Kawabata, M. (2006) Impact of single and multiple residual sprayings of pyrethroid insecticides against *Triatoma dimidiata* (Reduviidae; Triatominae), the principal vector of Chagas disease in Jutiapa, Guatemala. *American Journal of Tropical Medicine and Hygiene* **75**, 226–230.
- King, R.J., Cordon-Rosales, C., Cox, J., Davies, C.R. & Kitron, U.D. (2011) *Triatoma dimidiata* infestation in Chagas disease endemic regions of Guatemala: comparison of random and targeted cross sectional surveys. *PLOS Neglected Tropical Disease* **5**, 1–9.
- Lent, H. & Wygodzinsky, P. (1979) Revision of the Triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas disease. *Bulletin of the American Museum of Natural History* **163**, 125–520.
- Levy, M.Z., Bowman, N.M., Kawai, V., Waller, L.A., del Carpio, J.G.C., Benzaquen, E.C., Gilman, R.H. & Bern, C. (2006) Periurban *Trypanosoma cruzi*-infected *Triatoma infestans*, Arequipa, Peru. *Emerging Infectious Diseases* **12**, 1345–1352.
- Monroy, C., Mejia, M., Rodas, A., Rosales, R., Horio, M. & Tabaru, Y. (1998a) Comparison of indoor searches with whole house demolition collection of the vectors of Chagas disease and their indoor distribution. *Medical Entomology and Zoology* **49**, 195–200.
- Monroy, C., Rodas, A., Mejia, M. & Tabaru, Y. (1998b) Wall plastering and paints as methods to control vectors of Chagas disease in Guatemala. *Medical Entomology and Zoology* **49**, 187–193.
- Monroy, C., Rodas, A., Mejia, M., Rosales, R. & Tabaru, Y. (2003) Epidemiology of Chagas disease in Guatemala: infection rate of *Triatoma dimidiata*, *Triatoma nitida* and *Rhodnius prolixus* (Hemiptera, Reduviidae) with *Trypanosoma cruzi* and *Trypanosoma rangeli* (Kinetoplastida, Trypanosomatidae). *Memorias do Instituto Oswaldo Cruz* **98**, 305–310.
- Monroy, C., Bustamante, D.M., Pineda, S., Rodas, A., Castro, X., Ayala, V., Quinones, J. & Moguel, B. (2009) House improvements and community participation in the control of *Triatoma dimidiata* re-infestation in Jutiapa, Guatemala. *Cadernos de Saude Publica* **25**, S168–S178.
- Muehlen, M., Feldmeier, H., Wilcke, T., Winter, B. & Heukelbach, J. (2006) Identifying risk factors for tungiasis and heavy infestation in a resource-poor community in northeast Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **100**, 371–380.
- Nakagawa, J., Hashimoto, K., Cordon-Rosales, C., Abraham Juarez, J., Trampe, R. & Marroquin Marroquin, L. (2003) The impact of vector control on *Triatoma dimidiata* in the Guatemalan department of Jutiapa. *Annals of Tropical Medicine and Parasitology* **97**, 288–297.
- Nakagawa, J., Juarez, J., Nakatsuji, K., Akiyama, T., Hernandez, G., Macal, R., Flores, C., Ortiz, M., Marroquin, L., Bamba, T. & Wakai, S. (2005) Geographical characterization of the triatomine infestations in north-central Guatemala. *Annals of Tropical Medicine and Parasitology* **99**, 307–315.
- PAHO (2005) Estimacion cuantitativa de la enfermedad de Chagas en las Americas. *Pan American Health Organisation OPS/HDM/CD/425-06*, 28 pp.
- PAHO (2006) Agreements and recommendations from the IXth Annual IPCA Meeting. *Pan American Health Organisation PAHO/HDM/CD/CHA/422/06*, 5 pp.
- Paz-Bailey, G., Monroy, C., Rodas, A., Rosales, R., Tabaru, R., Davies, C. & Lines, J. (2002) Incidence of *Trypanosoma cruzi* infection in two Guatemalan communities. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **96**, 48–52.
- Petherick, A. (2010) Country by country. *Nature* **465**, S10–S11.

- Ponce, C.** (2007) Current situation of Chagas disease in Central America. *Memorias do Instituto Oswaldo Cruz* **102**, 41–44.
- Rizzo, N.R., Arana, B.A., Diaz, A., Cordon-Rosales, C., Klein, R.E. & Powell, M.R.** (2003) Seropositivity of *Trypanosoma cruzi* infection among school-age children in the endemic area of Guatemala. *American Journal of Tropical Medicine and Hygiene* **68**, 678–682.
- Rojas de Arias, A., Ferro, E.A., Ferreira, M.E. & Simancas, L.C.** (1999) Chagas disease vector control through different intervention modalities in endemic localities of Paraguay. *Bulletin of the World Health Organization* **77**, 331–339.
- Schofield, C.J.** (1994) *Triatominae: Biology and Control*. U.K., Eurocommunica Publications, Zeneca Public Health.
- Schofield, C.J. & Dujardin, J.P.** (1997) Chagas disease vector control in Central America. *Parasitology Today* **13**, 141–144.
- Starr, M.D., Rojas, J.C., Zeledon, R., Hird, D.W. & Carpenter, T.E.** (1991) Chagas disease risk-factors for house infestation by *Triatoma dimidiata*, the major vector of *Trypanosoma cruzi* in Costa Rica. *American Journal of Epidemiology* **133**, 740–747.
- Tabaru, Y., Monroy, C., Rodas, A. & Mejia, M.** (1999a) Chagas' disease vector surveillance in various residences in Santa Maria Ixhuatan, Department of Santa Rosa, Guatemala. *Medical Entomology and Zoology* **50**, 19–25.
- Tabaru, Y., Monroy, C., Rodas, A., Mejia, M. & Rosales, R.** (1999b) The geographical distribution of vectors of Chagas disease and population at risk of infestation in Guatemala. *Medical Entomology and Zoology* **50**, 3–8.
- WHO** (2002) Control of Chagas disease. World Health Organisation Technical Report Series 905. p. 109.
- WHO** (2004) The world health report 2004: changing history. World Health Organisation Reports. p. 168.
- Zeledon, R. & Vargas, L.G.** (1984) The role of dirt floors and of firewood in rural dwellings in the epidemiology of Chagas disease in Costa Rica. *American Journal of Tropical Medicine and Hygiene* **33**, 232–235.
- Zeledon, R., Zuniga, A. & Swartzwelder, J.C.** (1969) The camouflage of *Triatoma dimidiata* and the epidemiology of Chagas disease in Costa Rica. *Boletin Chileno de Parasitologia* **24**, 106–108.
- Zeledon, R., Solano, G., Zuniga, A. & Swartzwe, J.** (1973) Biology and ethology of *Triatoma dimidiata* (Latreille, 1811). 3. Habitat and blood sources. *Journal of Medical Entomology* **10**, 363–370.