



You May Also Like

**Artistic Renaissance in Frontotemporal Dementia**  
*Humanities* | April 3, 2018

**Fever in a Traveler Returning From Africa**  
*Case Report* | April 4, 2017



This Issue Citations 29

Letters

June 2, 2004

More

# High Prevalence of Obesity Among the Poor in Mexico

Lia C. Fernald, PhD, MBA; Juan Pablo Gutierrez, MS; Lynnette M. Neufeld, PhD; et al  
*JAMA*. 2004;291(21):2544-2545. doi:10.1001/jama.291.21.2544

**To the Editor:** The burden of disease in developing countries has traditionally been characterized by undernutrition and infectious diseases. However, lifestyle in many developing countries now parallels that in the developed world, with increasing prevalence of overweight and obesity.<sup>1</sup> It is unclear, however, how the prevalence of obesity varies across levels of socioeconomic status within such societies. We examined the prevalence of overweight and obesity among the rural poor in Mexico in comparison with a national sample.

**Methods**

We obtained data from 2 national surveys in Mexico. The first was conducted in 2000 in 45 260 adults as part of the National Health Survey, which was designed as a multistage, stratified, and clustered survey, nationally representative of the Mexican noninstitutionalized population.<sup>2</sup> The second survey was conducted in 2003 in 12 873 adults as part of the Social Welfare Survey, which was designed to be representative of the poorest (income <20th percentile), rural (towns with <2500 inhabitants) communities in 7 Mexican states.<sup>3</sup> These regions had a mean daily per capita income of US \$2. Women were more likely to be sampled because they were more often at home when the surveys were conducted.

Height and weight in both surveys were measured by trained and standardized personnel using standard techniques. Body mass index (BMI) was calculated as weight in kilograms divided by the square of height in meters. Overweight was defined as a BMI of 25.0 to 29.9 and obesity as a BMI of 30.0 or higher, in accordance with World Health Organization recommendations.<sup>4</sup> Values of BMI greater than 60 were excluded as implausible (representing <0.05% of individuals). Variables of interest for this analysis were available for 42 800 (94.6%) of the 2000 survey respondents and 12 844 (99.7%) of the 2003 survey respondents. For both surveys, sampling weights took into consideration unequal probabilities of selection resulting from sample design and nonresponse. Both studies were approved by the National Institute of Public Health in Mexico Ethical Review Committee and the 2003 survey was also approved by the University of California, Berkeley Committee on Human Research. All participants provided written informed consent.

**Results**

In the 2003 sample from low-income, rural regions of Mexico, the combined prevalence of overweight and obesity was nearly 60% in women and more than 50% in men (Table 1), which was slightly lower than the prevalence found in the nationally representative sample from 2000 (67% in women and 61% in men). However, this difference was not significant.

**Table.** Prevalence of Overweight and Obesity in Women and Men in Mexico From a Nationally Representative Sample and a Sample of Low-Income Rural Mexicans\*

	% (95% CI)			
	National Health Survey, 2000†		Social Welfare Survey, 2003‡	
	Women (n = 29 418)	Men (n = 13 385)	Women (n = 8905)	Men (n = 3964)
Age, yr	40.9 (34.1-47.7)	46.1 (42.7-49.5)	36.1 (37.7-34.6)	47.8 (47.2-48.4)
Underweight (BMI <18.5)	1.8 (1.6-1.9)	1.9 (1.7-2.1)	1.3 (1.1-1.6)	1.9 (1.7-2.2)
Normal weight (BMI 18.5-24.9)	31.1 (30.7-31.5)	36.0 (35.6-36.5)	39.5 (37.4-41.6)	45.3 (44.9-45.8)
Overweight (BMI 25.0-29.9)	36.5 (36.0-37.1)	40.4 (39.9-41.0)	36.8 (35.0-38.6)	38.9 (38.6-41.1)
Obese (BMI ≥30.0)	26.2 (25.8-26.7)	26.4 (26.0-26.9)	22.2 (20.4-24.1)	13.6 (11.9-15.3)

\*BMI, body mass index; prevalence of overweight and obesity defined by World Health Organization criteria. †Designed as a multistage, stratified and clustered survey, nationally representative of the Mexican noninstitutionalized population. ‡Designed to be representative of the poorest income <20th percentile, rural (towns with <2500 inhabitants) communities in 7 states of Mexico. §Data presented as mean (95% CI).

**Comment**

There is a high prevalence of overweight and obesity in the poorest segment of the Mexican population, and the prevalence is only slightly lower than that reported for a nationally representative health survey. The survey does not contain data about differences among regions of Mexico. Although the data do not permit conclusions about the causes of obesity in this population, we note that the prevalence of obesity we found is equal to or higher than in other Latin American countries.<sup>5</sup> Thus, it is likely that the population of the entire region is at risk for the comorbid conditions that accompany overweight and obesity, including hypertension, insulin resistance, and diabetes, all of which substantially increase the risk for cardiovascular disease.<sup>1,4</sup> This high prevalence of obesity coexists with traditional diseases of the poor, such as linear growth retardation, anemia, and deficiency of micronutrients in children.<sup>6</sup>

**Funding/Support:** This study was supported by the National Institute of Child Health and Human Development, the Fogarty International Center at the National Institutes of Health, and the Mexican government.

**Acknowledgment:** We thank Aurora Franco, Andrey Ryo Shiba, and Francisco Papaqui at the Instituto Nacional de Salud Pública in Mexico for their supervision of data collection, and all participants in both studies.

**References**

- Popkin BM. The nutrition transition and obesity in the developing world. *J Nutr*.2001;131:871S-873S. [PubMed](#) | [Google Scholar](#)
- Valdespino JL, Olaiz G, Lopez-Barajas MP, et al. *National Health Survey 2000, I: Housing, Population and Utilization of Health Services* [in Spanish]. Cuernavaca, Morelos, México: Instituto Nacional de Salud Pública; 2003.
- Behrman JR, Todd PE. *A Report on the Sample Sizes Used for the Evaluation of the Education, Health and Nutrition Program (PROGRESA) of Mexico*. Washington, DC: International Food Policy Research Institute; 1999. January 25, 1999.
- World Health Organization. *Physical Status: The Use and Interpretation of Anthropometry*. Geneva, Switzerland: World Health Organization; 1995.
- Filozof C, Gonzalez C, Sereday M, Mazza C, Braguinsky J. Obesity prevalence and trends in Latin-American countries. *Obes Rev*.2001;2:99-106. [PubMed](#) | [Google Scholar](#)
- Rivera JA, Sepulveda Amor J. Conclusions from the Mexican National Survey 1999: translating results into nutrition policy. *Salud Publica Mex*.2003;45(suppl 4):S565-S575. [PubMed](#) | [Google Scholar](#)

You May Also Like

**A 2-Step Cerebrospinal Algorithm for the Selection of Frontotemporal Lobar Degeneration Subtypes**  
*JAMA Neurology* | Research | March 19, 2018

**Prevalence and Causes of Unilateral Vision Impairment and Unilateral Blindness in Australia: The National Eye Health Survey**  
*JAMA Ophthalmology* | Research | March 1, 2018

**Association of Single-Nucleotide Polymorphisms in Age-Related Macular Degeneration With Pseudodrusen: Secondary Analysis of Data From the Comparison of AMD Treatments Trials**  
*JAMA Ophthalmology* | Research | May 3, 2018



<p><b>JAMA®</b></p> <p><b>CONTENT</b></p> <ul style="list-style-type: none"> <li>Home</li> <li>New Online</li> <li>Current Issue</li> </ul> <p><b>JOURNAL INFORMATION</b></p> <ul style="list-style-type: none"> <li>For Authors</li> <li>Editors &amp; Publishers</li> <li>RSS</li> <li>Contact Us</li> </ul> <p><b>JN Learning</b></p> <ul style="list-style-type: none"> <li>Store</li> <li>Apps</li> <li>Jobs</li> <li>Institutions</li> <li>Reprints &amp; Permissions</li> </ul> <p><b>Subscribe</b></p> <p>Go</p>	<p><b>JAMA Network™</b></p> <p><b>JOURNALS</b></p> <ul style="list-style-type: none"> <li>JAMA®</li> <li>JAMA Network Open</li> <li>JAMA Cardiology</li> <li>JAMA Dermatology</li> <li>JAMA Facial Plastic Surgery</li> <li>JAMA Internal Medicine</li> <li>JAMA Neurology</li> <li>JAMA Oncology</li> <li>JAMA Ophthalmology</li> <li>JAMA Otolaryngology-Head &amp; Neck Surgery</li> <li>JAMA Pediatrics</li> <li>JAMA Psychiatry</li> <li>JAMA Surgery</li> <li>Archives of Neurology &amp; Psychiatry (1919-1959)</li> </ul> <p><b>SITES</b></p> <ul style="list-style-type: none"> <li>AMA Manual of Style</li> <li>Art and Images in Psychiatry</li> <li>Breast Cancer Screening Guidelines</li> <li>Colorectal Screening Guidelines</li> <li>Declaration of Helsinki</li> <li>Depression Screening Guidelines</li> <li>Evidence-Based Medicine: An Oral History</li> <li>Fishbein Fellowship</li> <li>Genomics and Precision Health</li> <li>Health Disparities</li> <li>Hypertension Guidelines</li> <li>JAMA Network Audio</li> <li>JAMA Network Conferences</li> <li>Med Men</li> <li>Medical Education</li> <li>Opioid Management Guidelines</li> <li>Peer Review Congress</li> <li>Research Ethics</li> <li>Sepsis and Septic Shock</li> <li>Statins and Dyslipidemia</li> <li>Topics and Collections</li> </ul> <p><b>FEATURED ARTICLES</b></p> <ul style="list-style-type: none"> <li>ACS Breast Cancer Screening Guideline</li> <li>CDC Guideline for Prescribing Opioids</li> <li>Consensus Definitions for Sepsis and Septic Shock</li> <li>Income and Life Expectancy in the US</li> <li>JNC 8 Guideline for Management of High Blood Pressure</li> <li>President Obama on US Health Care Reform</li> <li>Screening for Colorectal Cancer</li> <li>Screening for Depression in Adults</li> <li>Statins for Primary Prevention of Cardiovascular Disease</li> <li>WMA Declaration of Helsinki, 7th Revision</li> </ul> <p><b>BLOGS</b></p> <ul style="list-style-type: none"> <li>The JAMA Forum</li> <li>Topics in Ophthalmology</li> <li>AMA Style Insider</li> </ul>	<p><b>INFORMATION FOR</b></p> <ul style="list-style-type: none"> <li>Authors</li> <li>Institutions &amp; Librarians</li> <li>Advertisers</li> <li>Subscription Agents</li> <li>Employers &amp; Job Seekers</li> <li>Media</li> </ul> <p><b>JAMA NETWORK PRODUCTS</b></p> <ul style="list-style-type: none"> <li>AMA Manual of Style</li> <li>JAMAevidence®</li> <li>JN Challenge™</li> <li>JN Listen™</li> <li>JN Reader™</li> <li>Peer Review Congress</li> </ul> <p><b>JN LEARNING</b></p> <ul style="list-style-type: none"> <li>CME Quizzes</li> <li>About CME &amp; MOC</li> <li>MOC Reporting Preferences</li> </ul>	<p><b>Help</b></p> <ul style="list-style-type: none"> <li>Subscriptions &amp; Renewals</li> <li>Email Subscriptions</li> <li>Update Your Address</li> <li>Contact Us</li> <li>Frequently Asked Questions</li> </ul> <p><b>JAMA CAREER CENTER</b></p> <ul style="list-style-type: none"> <li>Physician Job Listings</li> </ul> <p> </p> <p>Get the latest from JAMA</p> <p> <input type="text" value="Email address"/> </p> <p><b>Sign Up</b></p>
--	---	--	--