

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

van Lettow, M; Harries, AD; Kumwenda, JJ; Zijlstra, EE; Clark, TD; Taha, TE; Semba, RD; (2004) Micronutrient malnutrition and wasting in adults with pulmonary tuberculosis with and without HIV co-infection in Malawi. BMC infectious diseases, 4 (1). p. 61. ISSN 1471-2334 DOI: <https://doi.org/10.1186/1471-2334-4-61>

Downloaded from: <http://researchonline.lshtm.ac.uk/8939/>

DOI: <https://doi.org/10.1186/1471-2334-4-61>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

Research article

Open Access

Micronutrient malnutrition and wasting in adults with pulmonary tuberculosis with and without HIV co-infection in Malawi

Monique van Lettow*¹, Anthony D Harries², Johnny J Kumwenda³, Ed E Zijlstra³, Tamara D Clark¹, Taha E Taha⁴ and Richard D Semba¹

Address: ¹Johns Hopkins University School of Medicine, Baltimore, USA, ²National Tuberculosis Control Programme, Lilongwe, Malawi, ³College of Medicine, Blantyre, Malawi and ⁴Johns Hopkins University School of Public Health, Baltimore, USA

Email: Monique van Lettow* - mvlettow@imailbox.com; Anthony D Harries - adharries@malawi.net; Johnny J Kumwenda - jonnykumwenda@yahoo.com; Ed E Zijlstra - eezijlstra@malawi.net; Tamara D Clark - tclark@itsa.ucsf.edu; Taha E Taha - ttaha@jhsph.edu; Richard D Semba - rdsemba@jhmi.edu

* Corresponding author

Published: 21 December 2004

Received: 04 May 2004

BMC Infectious Diseases 2004, 4:61 doi:10.1186/1471-2334-4-61

Accepted: 21 December 2004

This article is available from: <http://www.biomedcentral.com/1471-2334/4/61>

© 2004 van Lettow et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Wasting and micronutrient malnutrition have not been well characterized in adults with pulmonary tuberculosis. We hypothesized that micronutrient malnutrition is associated with wasting and higher plasma human immunodeficiency virus (HIV) load in adults with pulmonary tuberculosis.

Methods: In a cross-sectional study involving 579 HIV-positive and 222 HIV-negative adults with pulmonary tuberculosis in Zomba, Malawi, anthropometry, plasma HIV load and plasma micronutrient concentrations (retinol, α -tocopherol, carotenoids, zinc, and selenium) were measured. The risk of micronutrient deficiencies was examined at different severity levels of wasting.

Results: Body mass index (BMI), plasma retinol, carotenoid and selenium concentrations significantly decreased by increasing tertile of plasma HIV load. There were no significant differences in plasma micronutrient concentrations between HIV-negative individuals and HIV-positive individuals who were in the lowest tertile of plasma HIV load. Plasma vitamin A concentrations $<0.70 \mu\text{mol/L}$ occurred in 61%, and zinc and selenium deficiency occurred in 85% and 87% respectively. Wasting, defined as $\text{BMI} < 18.5$ was present in 59% of study participants and was independently associated with a higher risk of low carotenoids, and vitamin A and selenium deficiency. Severe wasting, defined as $\text{BMI} < 16.0$ showed the strongest associations with deficiencies in vitamin A, selenium and plasma carotenoids.

Conclusions: These data demonstrate that wasting and higher HIV load in pulmonary tuberculosis are associated with micronutrient malnutrition.

Background

Approximately one-third of the world's population is infected with *Mycobacterium tuberculosis*, and the majority

live in less developed countries where human immunodeficiency virus (HIV) infection is spreading rapidly. The World Health Organization (WHO) estimates that the

number of new cases of tuberculosis and the proportion with coexisting HIV infection will continue to increase [1]. Immunosuppression increases the risk of developing clinical tuberculosis, which contributes to the increased prevalence of tuberculosis in association with HIV infection. Malnutrition and wasting are associated with tuberculosis, and co-infection with HIV and tuberculosis may potentially exacerbate the wasting that occurs in tuberculosis or HIV infection alone [2-5]. Micronutrient deficiencies have been described in individuals with tuberculosis [6-17] and in those with HIV infection [17-23]. Several cross-sectional studies suggest that patients with tuberculosis are at high risk of deficiencies of vitamin A [7,10-12], thiamin [13], vitamin B₆ [14], folate [6,15], vitamin E [16], and zinc [10]. Poor selenium status has recently been shown to increase the risk of developing mycobacterial disease among HIV-infected injection drug users [24], but selenium status among HIV-infected adults with pulmonary tuberculosis has not been well characterized. Selenium plays an important role in the selenoenzyme glutathione peroxidase that protects cells against free radical damage and oxidative stress.

The relationship between severity of HIV disease and micronutrient malnutrition needs further elucidation. Such information would help identify subgroups that might benefit the most from nutritional interventions. Plasma HIV load was used as an indicator of severity of HIV disease, as HIV load tends to be higher in more active HIV disease. We hypothesized that wasting in pulmonary tuberculosis is associated with micronutrient malnutrition and that HIV-infected adults with pulmonary tuberculosis who have more active HIV disease, as reflected by higher HIV load, also have more severe micronutrient malnutrition. To test these hypotheses, we conducted a cross-sectional study to examine the relationship between wasting and micronutrient malnutrition in HIV-positive and HIV-negative adults with pulmonary tuberculosis in Zomba, Malawi.

Methods

The study population consisted of adults who presented with new sputum-positive pulmonary tuberculosis in Zomba Central Hospital between July 1999 and April 2003. Subjects were offered HIV testing and were screened for HIV antibodies after signing a written informed consent form. All subjects were given appropriate pre- and post-test HIV counseling. Subjects commenced treatment after enrollment and received standard short course chemotherapy for tuberculosis as per guidelines of the Malawi National Tuberculosis Program [25]. Adults with a previous history of treated pulmonary tuberculosis were excluded. Three sputum samples from each subject were examined with Auramine-O dark-fluorescent staining method. Sputum positive pulmonary tuberculosis was

considered proven when at least one out of three sputum stains showed acid-fast bacilli. HIV infection was diagnosed on the basis of a positive rapid test (Determine 1/2 Rapid test by Abbott, Abbott Laboratories, Johannesburg, SA) and confirmed by a positive enzyme-linked immunosorbent assay for HIV-1 antibodies (Wellcozyme; Wellcome Diagnostics, Dartford, Kent, UK). Plasma HIV load was measured using quantitative HIV-1 RNA PCR (Roche Amplicor Monitor, version 1.5, Branchburg, NJ, USA) with a sensitivity limit of 400 HIV RNA copies mL. CD4⁺ lymphocyte counts were not conducted due to limited resources. None of the participants were taking antiretroviral treatment.

The protocol was approved by the institutional review boards at the Johns Hopkins School of Medicine (Baltimore, Maryland, USA) and the College of Medicine, University of Malawi (Blantyre, Malawi), with final approval by the Office for Protection from Research Risk of the National Institutes of Health.

Nutritional assessment

Body weight was determined to the nearest 0.1 kg using an adult balance (Seca 700 balance, Seca Corporation, Hanover, MD, USA), and standing height was determined to the nearest cm. Body mass index (BMI) was calculated as body weight/height².

Plasma micronutrient concentrations

A venous blood sample was collected by venipuncture (Sarstedt Monovette, Newton, NC). Blood samples were shielded from bright light and immediately aliquoted and stored in cryotubes at -70°C. α -carotene, β -carotene, β -cryptoxanthin, lycopene, lutein, zeaxanthin, retinol, and α -tocopherol concentrations were measured in 100 μ L of plasma by high performance liquid chromatography using a modified method from the Nutrition Laboratory, Inorganic Toxicology and Nutrition Branch Division of Laboratory Sciences, National Center of Environmental Health, Centers of Disease Control and Prevention (Rosemary Schleicher, personal communication) [27]. Total plasma carotenoids were defined as the sum of α -carotene, β -carotene, β -cryptoxanthin, lycopene, lutein and zeaxanthin in μ mol/L. Plasma trace element concentrations were measured using a Perkin Elmer model AAnalyst 600 atomic absorption spectrometer equipped with Zeeman background correction, a THGA graphite furnace, and an AS800 auto sampler (Perkin Elmer Corp., Norwalk, CT). Quality control was assessed by repeated analysis of pooled human plasma controls run at the beginning and the end of each analysis. Standard curves were run periodically using standard reference material 986C (National Institute of Standards and Technology, Gaithersburg, MD). Throughout all analyses, the plasma samples were run in a masked fashion.

Table 1: Characteristics of adults presenting with pulmonary tuberculosis in Zomba, Malawi – by plasma HIV load

Characteristic ¹	HIV negative	HIV positive* Plasma HIV Load (copies/mL)				P-value**
	0 n = 222	≤ 133 200 n = 185	133 200 – 406 000 n = 186	> 406 000 n = 186		
Sex (% Female)	45.9	63.2	58.1	55.9	0.07	
Age (years)	33 ± 12	32 ± 12	33 ± 12	33 ± 12	0.01	
Body mass index (BMI) (wt/ht ²)	18.6 ± 2.9	19.0 ± 2.6	18.3 ± 3.0	17.3 ± 2.7	0.0001	
Wasting: ² No, BMI ≥ 18.5	42.3	51.4	41.9	30.1	0.005	
Mild, BMI 17.0–18.49 (%)	32.0	27.0	22.6	24.2	0.04	
Moderate, BMI 16.0–16.99 (%)	13.1	10.3	17.2	18.8	0.04	
Severe, BMI <16.0 (%)	12.6	11.4	18.3	26.9	0.001	
Retinol (µmol/L)	0.636 (0.367, 1.104)	0.603 (0.336, 1.085)	0.585 (0.321, 1.066)	0.522 (0.157, 1.738)	0.001	
Vitamin A deficiency, retinol <0.70 µmol/L (%) ³	58.6	57.3	64.0	66.7	0.05	
Total Carotenoids (µmol/L) ⁴	0.846 (0.490, 1.459)	0.795 (0.476, 1.329)	0.700 (0.385, 1.279)	0.509 (0.279, 0.929)	0.0001	
α-tocopherol (µmol/L)	15.18 (11.71, 19.65)	14.90 (11.60, 19.16)	15.66 (11.85, 20.71)	16.07 (11.53, 22.40)	0.02	
Vitamin E def., α-tocopherol <11.6 µmol/L (%) ³	13.1	14.6	11.8	12.4	0.67	
Zinc (µmol/L)	8.95 (7.01, 11.43)	8.83 (6.94, 11.25)	8.49 (6.44, 11.19)	9.15 (6.47, 12.94)	0.82	
Zinc deficiency, zinc <11.5 µmol/L (%) ³	84.2	88.1	87.6	77.4	0.09	
Selenium (µmol/L)	0.687 ± 0.23	0.664 ± 0.22	0.624 ± 0.22	0.559 ± 0.21	0.0001	
Selenium deficiency, selenium <0.89 µmol/L (%) ³	79.7	84.9	90.3	92.5	0.001	

¹ Mean ± SD for continuous variables with normal distribution, geometric mean (lower, upper SD) when distribution was not normal

² Grading based on WHO Expert report, reference 27.

³ Cut-offs based on reference 28.

⁴ α-carotene + β-carotene + β-cryptoxanthin + lycopene + lutein + zeaxanthin

* HIV load could not be determined for 21 individuals.

** ANOVA, linear trend across the 4 categories of plasma HIV load.

Data and statistical analysis

Data and statistical analysis were conducted using SAS 8.01 (SAS Institute Cary, NC, USA) and SPSS 9.0 (SPSS, Inc., Chicago, IL, USA). Comparisons between groups were made using *t*-tests and nonparametric Mann-Whitney *U*-tests. Univariate analysis of variance was used to test for linear trends across categories of plasma HIV load and BMI.

HIV load was categorized into tertiles. HIV negative subjects were assigned a fourth category of HIV load (category 0) when groups were merged for analysis. Nutritional status was assessed in adults with pulmonary tuberculosis with and without HIV co-infection. Subjects were separated into groups according to their extent of wasting. Mild wasting was defined as BMI 17.0–18.49, moderate wasting as BMI 16.0–16.99, and severe wasting as BMI <16.0, conform the WHO strata for BMI grading of severity of malnutrition [27].

Plasma retinol <0.70 µmol/L was considered consistent with vitamin A deficiency [28]. Vitamin E deficiency was defined as plasma α-tocopherol <11.6 µmol/L [28]. Zinc deficiency was defined as plasma zinc <11.5 µmol/L and

selenium deficiency as plasma selenium <0.89 µmol/L [28]. Because there is no standard cut-off for deficiency of carotenoids, we divided total plasma carotenoids into quartiles, with the lowest quartile considered to be the most consistent with deficiency.

To examine the risk of micronutrient deficiencies at different severity level of wasting, logistic regression models were fitted with retinol <0.70, α-tocopherol <11.6, zinc <11.5, selenium <0.89, and the lowest quartile of total carotenoids as the outcome variable. Multivariate logistic regression models were conducted to adjust for sex, age and HIV load. A significance level of *P* < 0.01 was used in this study.

Results

The study population consisted of 579 HIV-positive and 222 HIV-negative adults with sputum-positive pulmonary tuberculosis. Among the total study population, 66% (232/352) of male and 77% (347/449) of female participants were HIV-positive. The mean age among all subjects was 33 years (range 18–59 years). The majority of subjects were wasted, as 59% of subjects had a BMI <18.5, 32% of subjects had a BMI <17.0, and 17% of all subjects were

Table 2: Risk of micronutrient deficiencies at different severity levels of wasting in adults with pulmonary tuberculosis with and without HIV co-infection.

Deficiency	Mild wasting BMI 17.0–18.49		Moderate wasting BMI 16.0–16.99		Severe wasting BMI <16.0	
	O.R. (95% C.I.)*	P-value	O.R. (95% C.I.)*	P-value	O.R. (95% C.I.)*	P-value
Vitamin A deficiency	0.81 (0.58–1.13)	0.20	1.59 (1.03–2.47)	0.03	3.51 (2.19–5.72)	0.0001
Lowest quartile of Total Carotenoids	0.92 (0.62–1.36)	0.67	2.46 (1.57–3.85)	0.0001	1.82 (1.18–2.83)	0.007
Vitamin E deficiency	0.86 (0.54–1.37)	0.54	1.24 (0.70–2.18)	0.46	1.13 (0.64–1.97)	0.68
Zinc deficiency	0.71 (0.45–1.15)	0.17	0.61 (0.37–1.00)	0.05	0.76 (0.46–1.24)	0.27
Selenium deficiency	1.59 (1.04–2.43)	0.03	1.16 (0.62–2.17)	0.65	3.25 (1.38–7.62)	0.007

* Adjusted for sex (male), age (per year) and HIV load (quartiles, where category 0 is HIV negative and category 3 is the highest).

severely wasted as defined by BMI<16.0. Plasma retinol concentrations <0.70 $\mu\text{mol/L}$ occurred in 61% of all subjects. Vitamin E, zinc, and selenium deficiency occurred in 13%, 85% and 87% respectively. Table 1 shows characteristics of study participants, such as sex, age, BMI, and plasma carotenoids, retinol, α -tocopherol, zinc and selenium by categories of plasma HIV load. BMI, plasma retinol, total carotenoids and selenium concentrations decreased by increasing plasma HIV load. Age, the proportion of individuals with BMI <18.5, BMI <16.0 and selenium deficiency were increased with increasing plasma HIV load. Plasma α -tocopherol, zinc or the proportion of individuals with vitamin A, vitamin E, or zinc deficiencies were not significantly different across the categories of plasma HIV load. When exploring across the categories separately, there were no significant differences between HIV-negative individuals compared with HIV-positive individuals in the lowest tertile of viral load.

Table 2 shows adjusted odds ratios (O.R.) and 95% confidence intervals (C.I.) for independent associations between wasting and micronutrient deficiencies. Wasting defined as BMI<18.5 was associated with vitamin A deficiency, low plasma carotenoids and selenium deficiency. The odds ratio for an independent association with vitamin A deficiency was 2.86 (95% C.I. 2.11–3.89) when adjusted for sex, age, and plasma HIV load. The adjusted odds ratio for an independent association with the lowest quartile of total carotenoids was 2.96 (95% C.I. 1.99–4.44). The adjusted odds ratio for an independent association with selenium deficiency was 1.59 (95% C.I. 1.04–2.43). When separating severity levels of wasting; mild wasting did not show association with deficiencies, moderate wasting was associated with vitamin A deficiency and severe wasting was significantly associated with vitamin A deficiency, low plasma carotenoids and selenium deficiency. (Table 2) Wasting was not associated with vitamin E or zinc deficiency.

Figures 1, 2 and 3 show plasma retinol, total plasma carotenoids, and plasma selenium concentrations with 95% C.I by severity of wasting and categories of plasma HIV load.

Plasma retinol concentrations significantly decreased with the increase of plasma HIV load among non-wasted adults with pulmonary tuberculosis ($P = 0.004$). Total carotenoid concentrations significantly decreased with the increase of plasma HIV load among non-wasted, mildly wasted, moderately wasted and severely wasted adults ($P = 0.0001$, $P = 0.002$, $P = 0.001$ and $P = 0.001$, respectively). Selenium concentrations decreased significantly with the increase of plasma HIV load among non-wasted and severely wasted adults with pulmonary tuberculosis ($P = 0.0001$ and $P = 0.03$, respectively). Among the HIV negative adults and those in the 1st and 2nd tertile of HIV load, plasma retinol, total carotenoids and selenium concentrations significantly decrease with the increasing severity of wasting. Among those in the 3rd tertile of HIV load, only plasma retinol concentrations significantly decreased with the increasing severity of wasting. This trend did not reach significance for plasma carotenoid and selenium concentrations.

Discussion

The present study shows that micronutrient malnutrition and wasting are more severe in adults with pulmonary tuberculosis who have higher plasma HIV load. The association between high plasma HIV load and nutrient deficiencies was strongest for the major plasma carotenoids and selenium. Overall in this study population, both HIV-positive and HIV-negative adults with pulmonary tuberculosis were extremely malnourished as indicated by BMI and plasma micronutrient concentrations. About one-third of the adults in this study had a BMI <17.0, a cut-off that is predictive of mortality in adults co-infected with tuberculosis and HIV [29].




Figure 1

Log-transformed mean plasma retinol concentrations with 95% C.I. are depicted by severity of wasting and plasma HIV load. Among the not-wasted adults with pulmonary tuberculosis log mean plasma retinol concentration significantly decrease with the increase of plasma HIV load ($P = 0.004$). Among those with mild, moderate and severe wasting this linear trend did not reach significance. Among the HIV negative adults with pulmonary tuberculosis, log mean plasma retinol concentration significantly decrease with the increasing severity of wasting ($P = 0.0001$). The same trend appears among those in the 1st, 2nd and 3rd tertile of HIV load; $P = 0.0001$, $P = 0.0001$ and $P = 0.01$ respectively.

To our knowledge, this is the first study to demonstrate that selenium status is extremely poor among HIV-infected adults with pulmonary tuberculosis, and that the extent of selenium deficiency is associated with higher plasma HIV load. This observation may be of potential importance because selenium deficiency has been associated with increased mortality during HIV infection [30], and selenium supplementation for HIV-infected adults has been shown to reduce morbidity [31]. In the present study, selenium deficiency occurred in 87% of the participants, which, to our knowledge, may be the highest

prevalence of selenium deficiency reported in an HIV-infected group of patients. It is unknown whether selenium supplementation will reduce morbidity and mortality among HIV-infected adults with pulmonary tuberculosis.

Carotenoids are among the most important dietary antioxidants found in human plasma, and this study shows that poor carotenoid status was associated with higher HIV load and with wasting. Plasma carotenoid concentrations are widely considered to be the most accurate indi-




Figure 2
 Log-transformed mean plasma total carotenoid concentrations with 95% C.I. are depicted by severity of wasting and plasma HIV load. Among not-wasted, mildly wasted, moderately wasted and severely wasted log mean plasma total carotenoid concentrations significantly decrease with the increase of plasma HIV load ($P = 0.0001$, $P = 0.002$, $P = 0.001$ and $P = 0.001$, respectively). Among the HIV negative adults, and those in the 1st and 2nd tertile of plasma HIV load, log mean plasma total carotenoid concentrations significantly decreased with the increasing severity of wasting ($P = 0.007$, $P = 0.002$ and $P = 0.0001$, respectively). This trend did not reach significance among those in the 3rd tertile of plasma HIV load.

cator of dietary carotenoid intake [32]. It is not known whether adults with pulmonary tuberculosis and higher HIV load have lower plasma carotenoid concentrations because of increased oxidative stress, or whether these individuals are sicker and unable to consume enough carotenoid-rich foods. Further studies are needed in the future to address dietary intake of carotenoids in HIV-infected adults with pulmonary tuberculosis.

Low BMI is a known risk factor for mortality [5,29], and the present study showed that the risk of micronutrient deficiencies is highest in those with low BMI.

HIV-infected adults with wasting and high viral load were at the highest risk of more severe micronutrient malnutrition, suggesting that this subgroup might potentially benefit the greatest from nutritional interventions.




Figure 3
 Mean plasma selenium concentrations with 95% C.I. are depicted by severity of wasting and plasma HIV load. Among not-wasted and severely wasted adults, mean plasma selenium concentrations significantly decrease with the increase of plasma HIV load ($P = 0.0001$ and $P = 0.03$, respectively). This trend did not reach significance among those with mild and moderate wasting. Among the HIV negative adults, and those in the 1st and 2nd tertile of plasma HIV load, mean plasma selenium concentrations significantly decreased with the increasing severity of wasting ($P = 0.02$, $P = 0.008$ and $P = 0.0001$, respectively). This trend did not reach significance among those in the 3rd tertile of plasma HIV load.

The cross sectional design of this study restricts our conclusions and does not provide information on whether poor nutritional status is a predictor of more severe pulmonary tuberculosis. It is unknown whether nutritional interventions will slow progression of disease or reduce wasting associated with morbidity and mortality if added to tuberculosis chemotherapy. Controlled clinical trials currently in progress in developing countries should help

provide insight into the role of micronutrient supplementation for HIV-positive and HIV-negative adults with pulmonary tuberculosis.

Conclusions

The present study shows that micronutrient malnutrition and wasting are more severe in adults with pulmonary tuberculosis who have higher HIV load. The association

between high plasma HIV load and nutrient deficiencies was strongest for the major plasma carotenoids and selenium. Further longitudinal investigations are needed to determine whether deficiencies in micronutrients are independent risk factors for increased morbidity and mortality.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

Overall guidance and initial study design was provided by RS. MvL has been in charge of the collection and analysis of data and writing of the manuscript. Provision of advice was given by AH, JK, EZ, TC and TT. All authors read and approved the final manuscript.

Acknowledgements

Supported in part by the National Institutes of Health (AI41956), the Fogarty International Center, and the United States Agency for International Development (Cooperative Agreement HRN A-0097-00015-00).

We thank Dana Totin Moncrief, Barbara Dancheck, Amanda Ray, and Michelle Ricks for their contributions and guidance in laboratory and data analyses. We thank the research team for their diligence

References

- Dolin PJ, Raviglione MC, Kochi A: **Global tuberculosis incidence and mortality during 1990-2000.** *Bull World Health Organ* 1994, **72**:213-220.
- Macallan DC: **Malnutrition in tuberculosis.** *Diagn Microbiol Infect Dis* 1999, **34**:153-157.
- Lucas SB, De Cock KM, Hounnou A, Peacock C, Diomande M, Honde M, Beaumel A, Kestens L, Kadio A: **Contributions of tuberculosis to slim disease in Africa.** *BMJ* 1994, **308**:1531-1533.
- Niyongabo T, Henzel D, Idi M, Nimubona S, Gikoro E, Melchior JC, Matheron S, Kamanfu G, Samb B, Messing B, Begue J, Aubry P, Larouze B: **Tuberculosis, human immunodeficiency virus infection, and malnutrition in Burundi.** *Nutrition* 1999, **15**:289-293.
- Lettow van M, Fawzi WW, Semba RD: **Triple trouble: the role of malnutrition in tuberculosis and human immunodeficiency virus co-infection.** *Nutr Rev* 2003, **61**:81-90.
- Markkanen T, Levanto A, Sallinen V, Virtanen S: **Folic acid and vitamin B₁₂ in tuberculosis.** *Scand J Haematol* 1967, **4**:283-291.
- Evans DIK, Attock B: **Folate deficiency in pulmonary tuberculosis: relationship to treatment and to serum vitamin A and beta-carotene.** *Tubercle* 1971, **52**:288-294.
- Cameron SJ, Horne NW: **The effect of tuberculosis and its treatment on erythropoiesis and folate activity.** *Tubercle* 1971, **52**:37-48.
- Chanarin I, Stephenson E: **Vegetarian diet and cobalamin deficiency: their association with tuberculosis.** *J Clin Pathol* 1988, **41**:759-762.
- Karyadi E, Schultink W, Nelwan RH, Gross R, Amin Z, Dolmans WM, van der Meer JW, Hautvast JG, West CE: **Poor micronutrient status of active pulmonary tuberculosis patients in Indonesia.** *J Nutr* 2000, **130**:2953-2958.
- Smurova TF, Prokopiev DI: **Vitamin A and carotene content in the blood of patients with pulmonary tuberculosis and diabetes mellitus.** *Probl Tuberk* 1969, **11**:50-55.
- Prokopiev DI: **Vitamin A content and carotene in blood plasma in pulmonary tuberculosis.** *Ter Arkh* 1966, **38**:54-59.
- Arkipova OP: **Impact of tuberculosis infection and antibacterial preparations on thiamin metabolism.** *Voprosy Meditsinskoi Khimii* 1975, **21**:449-560.
- Miasnikov VG: **Some indices of vitamin B₆ metabolism in patients with pulmonary tuberculosis in elderly and old persons.** *Vrachebnoe Delo* 1969, **3**:77-79.
- Line DH, Seitanidis B, Morgan JO, Hoffbrand AV: **The effect of chemotherapy on iron, folate and vitamin B₁₂ metabolism in tuberculosis.** *Q J Med* 1971, **40**:331-340.
- Panasiuk AV, Penenko OR, Kuz'menko IV, Suslov EI, Klimentenko MT, Kuznitsa NI, Tumanova TA, Makovetskii VP, Donchenko GV: **Vitamin E and its structural analogs in tuberculosis.** *Ukr Biokhim Zh* 1991, **63**:83-88.
- Semba RD, Tang AM: **Micronutrients and the pathogenesis of HIV infection.** *Br J Nutr* 1999, **81**:181-189.
- Bogden JD, Baker H, Frank O, Perez G, Kemp F, Bruening K, Louria D: **Micronutrient status and human immunodeficiency virus (HIV) infection.** *Ann N Y Acad Sci* 1990, **587**:189-195.
- Beach RS, Mantero-Atienza E, Shor-Posner G, Javier JJ, Szapocznik J, Morgan R, Sauberlich HE, Cornwell PE, Eisdorfer C, Baum MK: **Specific nutrient abnormalities in asymptomatic HIV-1 infection.** *AIDS* 1992, **6**:701-708.
- Ullrich R, Schneider T, Heise W, Schmidt W, Averdunk R, Riecken EO, Zeitz M: **Serum carotene deficiency in HIV-infected patients.** *AIDS* 1994, **8**:661-665.
- Baum MK, Mantero-Atienza E, Shor-Posner G: **Association of vitamin B₆ status with parameters of immune function in early HIV-1 infection.** *J Acquir Immune Defic Syndr* 1991, **4**:1122-1132.
- Ehrenpreis ED, Carlson SJ, Boorstein HL, Craig RM: **Malabsorption and deficiency of vitamin B₁₂ in HIV-infected patients with chronic diarrhea.** *Dig Dis Sci* 1994, **39**:2159-2162.
- Harriman GR, Smith PD, Horne MK, Fox CH, Koenig S, Lack EE, Lane HC, Fauci AS: **Vitamin B₁₂ malabsorption in patients with acquired immunodeficiency syndrome.** *Arch Intern Med* 1989, **149**:2039-2041.
- Shor-Posner G, Miguez MJ, Pineda LM, Rodriguez A, Ruiz P, Castillo G, Burbano X, Lecusay R, Baum M: **Impact of selenium status on the pathogenesis of mycobacterial disease in HIV-1-infected drug users during the era of highly active antiretroviral therapy.** *J Acquir Immune Defic Syndr* 2002, **29**:169-173.
- Manual of the National Tuberculosis Control Programme of Malawi, 4th edition.** Ministry of Health and Population, Malawi 1999.
- Sowell AL, Huff DL, Yeager PR, Caudill SP, Gunter EVV: **Simultaneous determination of retinol, α -tocopherol, lutein/zeaxanthin, β -cryptoxanthin, lycopene, trans- β -carotene, and four retinyl esters in serum by reverse-phase high performance liquid chromatography with multiwavelength detection.** *Clinical Chemistry* 1994, **40**:411-416.
- Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee.** *World Health Organ Tech Rep Ser* 1995, **854**:1-452.
- Sauberlich HE: *Laboratory tests for the assessment of nutritional status* Second edition. Boca Raton, CRC Press; 1999.
- Zacharia R, Spielman MP, Harries AD, Salaniponi FM: **Moderate to severe malnutrition in patients with tuberculosis is a risk factor associated with early death.** *Trans R Soc Trop Med Hyg* 2002, **96**:1-4.
- Baum MK, Shor-Posner G, Lai S, Zhang G, Lai H, Fletcher MA, Sauberlich H, Page JB: **High risk of HIV-related mortality is associated with selenium deficiency.** *J Acquir Immune Defic Syndr Hum Retrovirology* 1997, **15**:370-374.
- Burbano X, Miguez-Burbano MJ, McCollister K, Zhang G, Rodriguez A, Ruiz P, Lecusay R, Shor-Posner G: **Impact of selenium chemopreventive clinical trial on hospital admissions of HIV-infected participants.** *HIV Clin Trials* 2002, **3**:483-491.
- Institute of Medicine: **Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids.** Washington, D.C. National Academy of Sciences; 2000.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2334/4/61/prepub>